

Local Discontinuous Galerkin Method for the Diffusion Equation

We would like to solve the 1D diffusion equation with a Discontinuous Galerkin Method. The 1D diffusion equation is given as

$$q_t = q_{xx}.$$

If we were to naively apply DG methods, we could discretize the domain and consider piecewise polynomial approximation. We would then multiply by a test function and integrate by parts.

$$\begin{aligned} \int_{I_j} u_t v \, dx &= \int_{I_j} u_{xx} v \, dx \\ \int_{I_j} u_t v \, dx &= \left((\hat{u}_x v^-)_{j+1/2} - (\hat{u}_x v^+)_{j-1/2} \right) - \int_{I_j} u_x v_x \, dx \end{aligned}$$

and we use the average numerical flux

$$(\hat{u}_x)_{j+1/2} = \frac{(u_x^-)_{j+1/2} + (u_x^+)_{j+1/2}}{2}$$

This method is convergent and stable but it converges to the wrong solution.

Local Discontinuous Galerkin Method The Local Discontinuous Galerkin method proposes a different approach. First rewrite the diffusion equation as a system of first order equations.

$$r = q_x$$

$$q_t = r_x$$

The LDG method becomes the process of finding $q_h, r_h \in V_h$ in the DG solution space, such that for all test functions $v_h, w_h \in V_h$ and for all j the following equations are satisfied

$$\begin{aligned} \int_{I_j} r_h w_h \, dx &= \int_{I_j} (q_h)_x w_h \, dx \\ \int_{I_j} (q_h)_t v_h \, dx &= \int_{I_j} (r_h)_x v_h \, dx \end{aligned}$$

After integrating by parts, these equations are

$$\begin{aligned} \int_{I_j} r_h w_h \, dx &= \left((\hat{q}_h w_h^-)_{j+1/2} - (\hat{q}_h w_h^+)_{j-1/2} \right) - \int_{I_j} q_h (w_h)_x \, dx \\ \int_{I_j} (q_h)_t v_h \, dx &= \left((\hat{r}_h v_h^-)_{j+1/2} - (\hat{r}_h v_h^+)_{j-1/2} \right) - \int_{I_j} r_h (v_h)_x \, dx \end{aligned}$$

A common choice of numerical fluxes are the so-called alternating fluxes.

$$\begin{aligned} \hat{r}_h &= r_h^- \\ \hat{q}_h &= q_h^+ \end{aligned}$$

Implementation If we consider a single cell I_j , do a linear transformation from $x \in [x_{j-1/2}, x_{j+1/2}]$ to $\xi \in [-1, 1]$, and consider specifically the Legendre polynomial basis $\{\phi^k(\xi)\}$ with the following orthogonality property

$$\frac{1}{2} \int_{-1}^1 \phi^j(\xi) \phi^k(\xi) \, d\xi = \delta_{jk}$$

we can form a more concrete LDG method for implementing. The linear transformation can be expressed as

$$x = \frac{\Delta x}{2} \xi + \frac{x_{j-1/2} + x_{j+1/2}}{2}$$

or

$$\xi = \frac{2}{\Delta x} \left(x - \frac{x_{j-1/2} + x_{j+1/2}}{2} \right)$$

After this tranformation the diffusion equation become

$$q_t = \frac{4}{\Delta x^2} q_{\xi\xi}$$

on the cell I_j . We can then write this as the following system of first order equations.

$$r = \frac{2}{\Delta x} q_{\xi} q_t = \frac{2}{\Delta x} r_{\xi}$$

With the Legendre basis, the numerical solution on I_j can be written as

$$\begin{aligned} q|_{I_i} &\approx q_h|_{I_i} = \sum_{l=1}^M \left(Q_i^l \phi^l(\xi) \right) \\ r|_{I_i} &\approx r_h|_{I_i} = \sum_{l=1}^M \left(R_i^l \phi^l(\xi) \right) \end{aligned}$$

Now plugging these into the system and multiplying by a Legendre basis and integrating gives. I will use the following shorthand for numerical fluxes using one of the alternating flux options.

$$\begin{aligned} \hat{Q}_{i+1/2} &= \sum_{l=1}^M \left(Q_{i+1}^l \phi^l(-1) \right) \\ \hat{R}_{i+1/2} &= \sum_{l=1}^M \left(R_i^l \phi^l(1) \right) \end{aligned}$$

$$\begin{aligned} r &= \frac{2}{\Delta x} q_{\xi} \\ \sum_{l=1}^M \left(R_i^l \phi^l(\xi) \right) &= \frac{2}{\Delta x} \sum_{l=1}^M \left(Q_i^l \phi_{\xi}^l(\xi) \right) \\ \frac{1}{2} \int_{-1}^1 \sum_{l=1}^M \left(R_i^l \phi^l(\xi) \right) \phi^k(\xi) d\xi &= \frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(Q_i^l \phi_{\xi}^l(\xi) \right) \phi^k(\xi) d\xi \\ R_i^k &= \frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(Q_i^l \phi_{\xi}^l(\xi) \right) \phi^k(\xi) d\xi \\ R_i^k &= -\frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(Q_i^l \phi^l(\xi) \right) \phi_{\xi}^k(\xi) d\xi + \frac{1}{\Delta x} \left(\phi^k(1) \hat{Q}_{i+1/2} - \phi^k(-1) \hat{Q}_{i-1/2} \right) \end{aligned}$$

$$\begin{aligned}
q_t &= \frac{2}{\Delta x} r_\xi \\
\sum_{l=1}^M \left(\dot{Q}_i^l \phi^l(\xi) \right) &= \frac{2}{\Delta x} \sum_{l=1}^M \left(R_i^l \phi_\xi^l(\xi) \right) \\
\frac{1}{2} \int_{-1}^1 \sum_{l=1}^M \left(\dot{Q}_i^l \phi^l(\xi) \right) \phi^k(\xi) d\xi &= \frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(R_i^l \phi_\xi^l(\xi) \right) \phi^k(\xi) d\xi \\
\dot{Q}_i^k &= -\frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(U_i^l \phi_\xi^l(\xi) \right) \phi^k(\xi) d\xi + \frac{1}{\Delta x} \left(\phi^k(1) \hat{R}_{i+1/2} - \phi^k(-1) \hat{R}_{i-1/2} \right)
\end{aligned}$$

Now this is a system of ODEs, there are $M \times N$ ODEs if M is the spacial order and N is the number of cells.

Matrix Representation Some common matrices and vectors that appear in these equations are

$$\begin{aligned}
\mathbf{Q}_i &= \left[Q_i^l \right]_{l=1}^M \\
\phi(\xi) &= \left[\phi^k(\xi) \right]_{k=1}^M \\
\Phi(\xi_1, \xi_2) &= \phi(\xi_1) \phi^T(\xi_2) \\
A &= [a_{kl}]_{k,l=1}^M \\
a_{kl} &= \int_{-1}^1 \phi_\xi^k(\xi) \phi^l(\xi) d\xi
\end{aligned}$$

Also the numerical fluxes can be written as the following dot products

$$\begin{aligned}
\hat{Q}_{i+1/2} &= \sum_{l=1}^M \left(Q_{i+1}^l \phi^l(-1) \right) \\
&= \phi^T(-1) \mathbf{Q}_{i+1} \\
\hat{R}_{i+1/2} &= \sum_{l=1}^M \left(R_i^l \phi^l(1) \right) \\
&= \phi^T(1) \mathbf{R}_i
\end{aligned}$$

$$\begin{aligned}
R_i^k &= -\frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(Q_i^l \phi^l(\xi) \right) \phi_\xi^k(\xi) d\xi + \frac{1}{\Delta x} \left(\phi^k(1) \hat{Q}_{i+1/2} - \phi^k(-1) \hat{Q}_{i-1/2} \right) \\
R_i^k &= -\frac{1}{\Delta x} \sum_{l=1}^M \left(Q_i^l \int_{-1}^1 \phi^l(\xi) \phi_\xi^k(\xi) d\xi \right) + \frac{1}{\Delta x} \left(\phi^k(1) \hat{Q}_{i+1/2} - \phi^k(-1) \hat{Q}_{i-1/2} \right) \\
R_i^k &= -\frac{1}{\Delta x} (A \mathbf{Q}_i)_k + \frac{1}{\Delta x} \left(\phi^k(1) \phi^T(-1) \mathbf{Q}_{i+1} - \phi^k(-1) \phi^T(-1) \mathbf{Q}_i \right) \\
\mathbf{R}_i &= -\frac{1}{\Delta x} A \mathbf{Q}_i + \frac{1}{\Delta x} \left(\phi(1) \phi^T(-1) \mathbf{Q}_{i+1} - \phi(-1) \phi^T(-1) \mathbf{Q}_i \right) \\
\mathbf{R}_i &= -\frac{1}{\Delta x} A \mathbf{Q}_i + \frac{1}{\Delta x} (\Phi(1, -1) \mathbf{Q}_{i+1} - \Phi(-1, -1) \mathbf{Q}_i) \\
\mathbf{R}_i &= -\frac{1}{\Delta x} (A + \Phi(-1, -1)) \mathbf{Q}_i + \frac{1}{\Delta x} \Phi(1, -1) \mathbf{Q}_{i+1}
\end{aligned}$$

$$\begin{aligned}
Q_i^k &= -\frac{1}{\Delta x} \int_{-1}^1 \sum_{l=1}^M \left(R_i^l \phi^l(\xi) \right) \phi_\xi^k(\xi) d\xi + \frac{1}{\Delta x} \left(\phi^k(1) \hat{R}_{i+1/2} - \phi^k(-1) \hat{R}_{i-1/2} \right) \\
Q_i^k &= -\frac{1}{\Delta x} (A \mathbf{R}_i)_k + \frac{1}{\Delta x} \left(\phi^k(1) \phi^T(1) \mathbf{R}_i - \phi^k(-1) \phi^T(1) \mathbf{R}_{i-1} \right) \\
\mathbf{Q}_i &= -\frac{1}{\Delta x} (A - \Phi(1, 1)) \mathbf{R}_i - \frac{1}{\Delta x} \Phi(-1, 1) \mathbf{R}_{i-1}
\end{aligned}$$

Proving Stability In order to prove that this method is L^2 stable consider we sum both of the integral equations from before.

$$\begin{aligned}
& \int_{I_j} (u_h)_t v_h dx + \int_{I_j} q_h w_h dx = \left((q_h^+ v_h^-)_{j+1/2} - (q_h^+ v_h^+)_{j-1/2} \right) \\
& + \left((u_h^- w_h^-)_{j+1/2} - (u_h^- w_h^+)_{j-1/2} \right) - \int_{I_j} q_h (v_h)_x dx - \int_{I_j} u_h (w_h)_x dx
\end{aligned}$$

Consider using $v_h = u_h$ and $w_h = q_h$.

$$\begin{aligned}
& \int_{I_j} (u_h)_t u_h dx + \int_{I_j} q_h q_h dx = \left((q_h^+ u_h^-)_{j+1/2} - (q_h^+ u_h^+)_{j-1/2} \right) \\
& + \left((u_h^- q_h^-)_{j+1/2} - (u_h^- q_h^+)_{j-1/2} \right) - \int_{I_j} q_h (u_h)_x dx - \int_{I_j} u_h (q_h)_x dx
\end{aligned}$$

Consider the following shorthand notation

$$\begin{aligned}
B_j &= \int_{I_j} (u_h)_t u_h dx + \int_{I_j} q_h q_h dx \\
B_j &= \left((q_h^+ u_h^-)_{j+1/2} - (q_h^+ u_h^+)_{j-1/2} \right) + \left((u_h^- q_h^-)_{j+1/2} - (u_h^- q_h^+)_{j-1/2} \right) - \int_{I_j} q_h (u_h)_x dx - \int_{I_j} u_h (q_h)_x dx
\end{aligned}$$

This can be simplified in several ways. First simplify the left hand side.

$$\begin{aligned}
B_j &= \int_{I_j} (u_h)_t u_h dx + \int_{I_j} q_h q_h dx \\
B_j &= \frac{1}{2} \int_{I_j} \frac{d}{dt} (u_h^2) dx + \int_{I_j} q_h^2 dx \\
B_j &= \frac{1}{2} \frac{d}{dt} \int_{I_j} u_h^2 dx + \int_{I_j} q_h^2 dx \\
B_j &= \frac{1}{2} \frac{d}{dt} \|u_h\|_{L^2(I_j)}^2 + \|q_h\|_{L^2(I_j)}^2
\end{aligned}$$

Second the right hand side can be simplified.

$$\begin{aligned}
& \int_{I_j} q_h (u_h)_x dx + \int_{I_j} u_h (q_h)_x dx = \int_{I_j} q_h (u_h)_x + u_h (q_h)_x dx \\
& = \int_{I_j} (q_h u_h)_x dx \\
& = (q_h^- u_h^-)_{j+1/2} - (q_h^+ u_h^+)_{j-1/2}
\end{aligned}$$

Now

$$B_j = \left((q_h^+ u_h^-)_{j+1/2} - (q_h^+ u_h^+)_{j-1/2} \right) + \left((u_h^- q_h^-)_{j+1/2} - (u_h^- q_h^+)_{j-1/2} \right) - \left((q_h^- u_h^-)_{j+1/2} - (q_h^+ u_h^+)_{j-1/2} \right)$$

$$B_j = (q_h^+ u_h^-)_{j+1/2} - (u_h^- q_h^+)_{j-1/2}$$

Assuming periodic boundary conditions, and summing B_j over all cells

$$\begin{aligned} \sum_{j=1}^N (B_j) &= \sum_{j=1}^N \left((q_h^+ u_h^-)_{j+1/2} - (u_h^- q_h^+)_{j-1/2} \right) \\ &= - (u_h^- q_h^+)_{1/2} + \sum_{k=1}^N \left((q_h^+ u_h^-)_{k+1/2} - (u_h^- q_h^+)_{k+1/2} \right) + (q_h^+ u_h^-)_{N+1/2} \\ &= 0 \end{aligned}$$

This shows that

$$\begin{aligned} \sum_{j=1}^N (B_j) &= \sum_{j=1}^N \left(\frac{1}{2} \frac{d}{dt} \|u_h\|_{L^2(I_j)}^2 + \|q_h\|_{L^2(I_j)}^2 \right) \\ \frac{1}{2} \frac{d}{dt} \|u_h\|_{L^2}^2 + \|q_h\|_{L^2}^2 &= 0 \\ \frac{d}{dt} \|u_h\|_{L^2}^2 &\leq 0 \end{aligned}$$