

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

Discontinuous Galerkin Method for Solving Thin Film Equations

Caleb Logemann

Mathematics Department, Iowa State University

logemann@iastate.edu

December 13, 2018

Overview

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

1 Introduction

2 Method

- Convection
- Diffusion

3 Numerical Results

4 Conclusion

Motivation

Caleb Logemann

Introduction

Method

Convection


Diffusion

Numerical Results

Conclusion

References

- Aircraft Icing
- Runback



- Industrial Coating

Model Equations

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Navier-Stokes Equation

$$\nabla \cdot \mathbf{u} = 0$$

$$\partial_t \mathbf{u} + \nabla \cdot (\mathbf{u}\mathbf{u}) = -\frac{1}{\rho} \nabla p + \frac{1}{\rho} \nabla \cdot \sigma + \mathbf{g}$$

$$\partial_t h_s + (u, v)^T \cdot \nabla h_s = w$$

$$\partial_t h_b + (u, v)^T \cdot \nabla h_b = w$$

- Lubrication or reduced Reynolds number approximation
- Thin-Film Equation - 1D with q as fluid height.

$$q_t + (f(x, t)q^2 - g(x, t)q^3)_x = -(h(x, t)q^3 q_{xxx})_x$$

Operator Splitting

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Simplified Model

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Operator Splitting

$$q_t + (q^2 - q^3)_x = 0$$

$$q_t + (q^3 u_{xxx})_x = 0$$

■ Strang Splitting

$\frac{1}{2}\Delta t$ step of Convection

$$q_t + (q^2 - q^3)_x = 0$$

Δt step of Diffusion

$$q_t + (q^3 u_{xxx})_x = 0$$

$\frac{1}{2}\Delta t$ step of Convection

$$q_t + (q^2 - q^3)_x = 0$$

Convection

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Convection Equation

$$\begin{aligned}q_t + f(q)_x &= 0 & (0, T) \times \Omega \\f(q) &= q^2 - q^3\end{aligned}$$

■ Weak Form

Find q such that

$$\int_{\Omega} (q_t v - f(q) v_x) dx = 0$$

for all test functions v

Notation

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion


References

- Partition the domain, $[a, b]$ as

$$a = x_{1/2} < \cdots < x_{j-1/2} < x_{j+1/2} < \cdots < x_{N+1/2} = b$$

- $I_j = [x_{j-1/2}, x_{j+1/2}]$

- $x_j = \frac{x_{j+1/2} + x_{j-1/2}}{2}$.



Runge Kutta Discontinuous Galerkin

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Find $Q(t, x)$ such that for each time $t \in (0, T)$,
 $Q(t, \cdot) \in V_h = \left\{ v \in L^1(\Omega) : v|_{I_j} \in P^k(I_j) \right\}$

$$\begin{aligned} \int_{I_j} Q_t v \, dx &= \int_{I_j} f(Q) v_x \, dx \\ &\quad - \left(\mathcal{F}_{j+1/2} v^-(x_{j+1/2}) - \mathcal{F}_{j-1/2} v^+(x_{j-1/2}) \right) \end{aligned}$$

for all $v \in V_h$

- Rusanov/Local Lax-Friedrichs Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{1}{2} \left(f(Q_{j+1/2}^-) + f(Q_{j+1/2}^+) \right) + \max_q \{ |f'(q)| \} (Q_{j+1/2}^- - Q_{j+1/2}^+)$$

- Solve this system of ODEs with any Explicit Strong Stability Preserving (SSP) Runge-Kutta Method.

Explicit SSP Runge Kutta Methods

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Forward Euler

$$q^{n+1} = q^n + \Delta t L(q^n)$$

■ Second Order

$$q^* = q^n + \Delta t L(q^n)$$

$$q^{n+1} = \frac{1}{2}(q^n + q^*) + \frac{1}{2}\Delta t L(q^*)$$

Diffusion

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Diffusion Equation

$$q_t = -(q^3 q_{xxx})_x \quad (0, T) \times \Omega$$

■ Linearize operator at $t = t^n$, let $f(x) = q^3(t = t^n, x)$

$$q_t = -(f(x) q_{xxx})_x \quad (0, T) \times \Omega$$

Finite Difference Approach

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Let cell centers, x_i form finite difference grid.
- Finite difference space, \mathbb{R}^N .
- $Q_{DG} \in V_h \rightarrow Q_{FD} \in \mathbb{R}^N$

$$(Q_{FD})_i = \frac{1}{h} \int_{K_i} Q_{DG} \, dx$$

- $Q_{FD} \in \mathbb{R}^N \rightarrow Q_{DG} \in V_h$

$$Q_{DG}|_K \in P^1(K)$$

$$\frac{1}{h} \int_{K_i} Q_{DG} \, dx = (Q_{FD})_i$$

$$\partial_x Q_{DG}|_{K_i} = \frac{(Q_{FD})_{i+1} - (Q_{FD})_{i-1}}{2h}$$

Finite Difference Approximation

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

$$(q_{xxx})_{i+1/2} \approx \frac{-Q_{i-1} + 3Q_i - 3Q_{i+1} + Q_{i+2}}{h^3}$$

$$F_{i+1/2} = \frac{f(x_{i+1/2}^+) + f(x_{i+1/2}^-)}{2}$$

$$(-(f(x)q_{xxx})_x)_i \approx -\frac{F_{i+1/2}(q_{xxx})_{i+1/2} - F_{i-1/2}(q_{xxx})_{i-1/2}}{h}$$

Implicit L-Stable Runge Kutta

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

■ Backward Euler

$$q^{n+1} = q^n + \Delta t L(q^{n+1})$$

■ 2nd Order

$$q^* = q^n + \frac{1}{4} \Delta t (L(q^n) + L(q^*))$$

$$3q^{n+1} = 4q^* - q^n + \Delta t L(q^{n+1})$$

Riemann Problem

Caleb Logemann

Introduction

Method


Convection

Diffusion

Numerical Results

Conclusion

References



Riemann Problem

Caleb Logemann

Introduction

Method


Convection

Diffusion

Numerical Results

Conclusion

References



Square Wave

Caleb Logemann

Introduction

Method


Convection

Diffusion

Numerical Results

Conclusion

References



Square Wave

Caleb Logemann

Introduction

Method


Convection

Diffusion

Numerical Results

Conclusion

References



Future Work

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- Show second order convergence
- Runge Kutta IMEX
- Space and time dependent coefficients

Bibliography

Caleb Logemann

Introduction

Method

Convection

Diffusion

Numerical Results

Conclusion

References

- [1] Bernardo Cockburn and Chi-Wang Shu. “The local discontinuous Galerkin method for time-dependent convection-diffusion systems”. In: *SIAM Journal on Numerical Analysis* 35.6 (1998), pp. 2440–2463.
- [2] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [3] T.G. Myers and J.P.F. Charpin. “A mathematical model for atmospheric ice accretion and water flow on a cold surface”. In: *Int. J. Heat and Mass Transfer* 47.25 (2004), pp. 5483–5500.
- [4] Tim G Myers. “Thin films with high surface tension”. In: *SIAM review* 40.3 (1998), pp. 441–462.
- [5] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.
- [6] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.