

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator

Splitting

Conclusion

Discontinuous Galerkin Method for solving thin film equations

Caleb Logemann

Mathematics Department, Iowa State University

logemann@iastate.edu

September 30, 2017

Overview

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

1 Introduction

2 Convection

3 Hyper-Diffusion

4 Operator Splitting

5 Conclusion

Motivation

Caleb Logemann

Introduction

Convection


Hyper-Diffusion

Operator

Splitting

Conclusion

- Aircraft Icing
- Runback



- Industrial Coating

Model Equations

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator

Splitting

Conclusion

■ Navier-Stokes Equation

$$\rho_t + (\rho u)_x = 0$$

$$(\rho u)_t + (\rho u^2 + p)_x = \frac{4}{3Re} u_{xx}$$

$$E_t + (u(E + p))_x = \frac{1}{Re} \left(\frac{2}{3} (u^2)_{xx} + \frac{\gamma}{(\gamma - 1)Pr} \left(\frac{p}{\rho} \right)_{xx} \right)$$

■ Asymptotic Limit, $\rho \ll L$

■ Thin-Film Equation - 1D with q as fluid height.

$$q_t + (f(x, t)q^2 - g(x, t)q^3)_x = -(h(x, t)q^3 q_{xxx})_x$$

Current Model

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator

Splitting

Conclusion

■ Simplified Expression

$$q_t + (q^2 - q^3)_x = -(q^3 q_{xxx})_x$$

■ Operator Splitting

$$q_t + (q^2 - q^3)_x = 0$$

$$q_t + (q^3 q_{xxx})_x = 0$$

Introduction to Discontinuous Galerkin

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion


- Let \mathcal{T}_h partition the domain, $\Omega = [a, b]$

$$a = x_{1/2} < \cdots < x_{i-1/2} < x_{i+1/2} < \cdots < x_{N+1/2} = b$$

- $K_i \in \mathcal{T}_h = [x_{i-1/2}, x_{i+1/2}]$

- $h = x_{i+1/2} - x_{i-1/2}$

- $x_i = \frac{x_{i+1/2} + x_{i-1/2}}{2}$.



Function Spaces

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

- $P^k(K)$ - polynomials of degree less than or equal to k on $K \in \mathcal{T}_h$

-

$$V_h = \{v \in L^2(\Omega) : v|_K \in P^k(K), \quad \forall K \in \mathcal{T}_h\}$$

Numerical Solutions

Caleb Logemann

Introduction

Convection

Hyper-Diffusion


Operator

Splitting

Conclusion

- Let $\{\phi^k(\xi)\}$ be the Legendre polynomials.
- Solution of order M on each cell

$$q|_{x \in v_i} \approx Q_i = \sum_{k=1}^M Q_i^k \phi^k(\xi)$$



Convection

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

■ Convection Equation

$$q_t + \frac{2}{\Delta x} f(q)_\xi = 0$$
$$f(q) = q^2 - q^3$$

■ Weak Form

$$\int_{-1}^1 \left(q_t \phi(\xi) + \frac{2}{\Delta x} f(q)_\xi \phi(\xi) \right) d\xi = 0$$

■ Runge-Kutta Discontinuous Galerkin

$$\dot{Q}_i^\ell = \frac{1}{\Delta x} \int_{-1}^1 f(Q_i) \phi_\xi^\ell d\xi - \frac{1}{\Delta x} (\mathcal{F}_{i+1/2} - \mathcal{F}_{i-1/2})$$

■ Rusanov Numerical Flux

$$\mathcal{F}_{j+1/2} = \frac{f(Q_{i+1}(-1)) + f(Q_i(1))}{2} \phi^\ell(1)$$

- Solve this system of ODEs with any Explicit Strong Stability Preserving (SSP) Runge-Kutta Method.

Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Numerical Example - Square Wave

Caleb Logemann

Introduction


Convection

Hyper-Diffusion

Operator

Splitting

Conclusion



Hyper-Diffusion

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator

Splitting

Conclusion

■ Hyper-Diffusion Equation

$$u_t + \frac{16}{\Delta x^4} (u^3 u_{\xi\xi\xi})_{\xi} = 0$$

■ Local Discontinuous Galerkin (LDG)

$$q = \frac{2}{\Delta x} u_{\xi}$$

$$r = \frac{2}{\Delta x} q_{\xi}$$

$$s = \frac{2}{\Delta x} u^3 r_{\xi}$$

$$u_t = -\frac{2}{\Delta x} s_{\xi}$$

Local Discontinuous Galerkin

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

$$\eta(\xi) = (U_i^n)^3$$

$$Q_i^\ell = -\frac{1}{\Delta x} \left(\int_{-1}^1 U_i \phi_\xi^\ell d\xi - \mathcal{F}(U)_{i+1/2}^\ell + \mathcal{F}(U)_{i-1/2}^\ell \right)$$

$$R_i^\ell = -\frac{1}{\Delta x} \left(\int_{-1}^1 Q_i \phi_\xi^\ell d\xi - \mathcal{F}(Q)_{i+1/2}^\ell + \mathcal{F}(Q)_{i-1/2}^\ell \right)$$

$$S_i^\ell = \frac{1}{\Delta x} \left(\int_{-1}^1 (R_i)_\xi \eta(\xi) \phi^\ell d\xi \right) \\ + \frac{1}{\Delta x} \left(\mathcal{F}(\eta)_{i+1/2} \mathcal{F}(R)_{i+1/2}^\ell - \mathcal{F}(\eta)_{i-1/2} \mathcal{F}(R)_{i-1/2}^\ell \right)$$

$$\dot{U}_i^\ell = \frac{1}{\Delta x} \left(\int_{-1}^1 S_i \phi_\xi^\ell d\xi - \mathcal{F}(S)_{i+1/2}^\ell + \mathcal{F}(S)_{i-1/2}^\ell \right)$$

Local Discontinuous Galerkin

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator


Splitting

Conclusion

$$\mathcal{F}(\eta)_{i+1/2} = \frac{1}{2}(\eta_{i+1}(-1) - \eta_i(1))$$

$$\mathcal{F}(\eta)_{i-1/2} = \frac{1}{2}(\eta_{i-1}(1) - \eta_i(-1))$$

$$\mathcal{F}(\ast)_{i+1/2}^\ell = \phi^\ell(1) \ast_{i+1/2}$$



Local Discontinuous Galerkin

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator

Splitting

Conclusion

- Explicit SSP Runge Kutta
 - Severe time step restriction
 - $\Delta t \sim \Delta x^4$
 - $\Delta x = .1 \rightarrow \Delta t \approx 10^{-4}$
 - $\Delta x = .01 \rightarrow \Delta t \approx 10^{-8}$
- Implicit SSP Runge Kutta
 - Linear System Solver
 - Stabilized BiConjugate Gradient
 - MultiGrid Solver

Operator Splitting

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

- Strang Splitting
 - 1 time step
 - $1/2$ time step for convection
 - 1 time step for hyper-diffusion
 - $1/2$ time step for convection
 - Second order splitting

Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion



Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion



Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion



Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion



Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion



Numerical Results - Riemann Problem

Caleb Logemann


Introduction

Convection

Hyper-Diffusion

**Operator
Splitting**

Conclusion



Future Work

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

- Higher dimensions
- Curved surfaces
- Space and time dependent coefficients
- Incorporation with air flow models
- Runge Kutta IMEX

Conclusion

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

- Thanks
 - James Rossmannith
 - Alric Rothmayer
- Questions?

Bibliography

Caleb Logemann

Introduction

Convection

Hyper-Diffusion

Operator
Splitting

Conclusion

- [1] Y. Ha, Y.-J. Kim, and T.G. Myers. “On the numerical solution of a driven thin film equation”. In: *J. Comp. Phys.* 227.15 (2008), pp. 7246–7263.
- [2] NASA. URL: http://icebox.grc.nasa.gov/gallery/images/C95_03918.html.
- [3] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.