

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Nonconservative Discontinuous Galerkin Method for Generalized Shallow Water Equations

Caleb Logemann James Rossmanith

Mathematics Department,
Iowa State University

logemann@iastate.edu

March 05, 2021

Overview

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

1 Generalized Shallow Water Equations

2 Nonconservative Products

3 Nonconservative DG Formulation

4 Results

Generalized Shallow Water

Caleb Logemann,
James
Rossmanith


Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References



Navier Stokes Equations with a free surface

$$\nabla \cdot \mathbf{u} = 0$$

$$\mathbf{u}_t + \nabla \cdot (\mathbf{u}\mathbf{u}) = -\frac{1}{\rho}\nabla p + \frac{1}{\rho}\nabla \cdot \sigma + \mathbf{g}$$

$$(h_s)_t + [u(t, x, y, h_s), v(t, x, y, h_s)]^T \cdot \nabla h_s = w(t, x, y, h_s)$$

$$(h_b)_t + [u(t, x, y, h_b), v(t, x, y, h_b)]^T \cdot \nabla h_b = w(t, x, y, h_b)$$

Polynomial Ansatz

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

$$\begin{aligned}\tilde{u}(t, x, y, \zeta) &= u_m(t, x, y) + u_d(t, x, y, \zeta) \\ &= u_m(t, x, y) + \sum_{j=1}^N (\alpha_j(t, x, y) \phi_j(\zeta))\end{aligned}$$

$$\begin{aligned}\tilde{v}(t, x, y, \zeta) &= v_m(t, x, y) + v_d(t, x, y, \zeta) \\ &= v_m(t, x, y) + \sum_{j=1}^N (\beta_j(t, x, y) \phi_j(\zeta))\end{aligned}$$

Orthogonality Condition

$$\int_0^1 \phi_j(\zeta) \phi_i(\zeta) d\zeta = 0 \quad \text{for } j \neq i$$

$$\phi_0(\zeta) = 1, \quad \phi_1(\zeta) = 1 - 2\zeta, \quad \phi_2(\zeta) = 1 - 6\zeta + 6\zeta^2$$

Constant Moments

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Continuity Equation

$$h_t + (hu_m)_x + (hv_m)_y = 0$$

Conservation of Momentum Equations

$$\begin{aligned} & (hu_m)_t + \left(h \left(u_m^2 + \sum_{j=1}^N \frac{\alpha_j^2}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_x \\ & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_y = -\frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \alpha_j \right) + hg(e_x - e_z(h_b)_x) \\ & (hv_m)_t + \left(h \left(v_m^2 + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) + \frac{1}{2} g e_z h^2 \right)_y \\ & + \left(h \left(u_m v_m + \sum_{j=1}^N \frac{\alpha_j \beta_j}{2j+1} \right) \right)_x = -\frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \beta_j \right) + hg(e_y - e_z(h_b)_y) \end{aligned}$$

Higher Order Moments

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

$$\begin{aligned}
 (h\alpha_i)_t + \left(2hu_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\alpha_k \right)_x + \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_y \\
 = u_mD_i - \sum_{j,k=1}^N B_{ijk}D_j\alpha_k - (2i+1)\frac{\nu}{\lambda} \left(u_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h}C_{ij} \right) \alpha_j \right) \\
 (h\beta_i)_t + \left(hu_m\beta_i + hv_m\alpha_i + h \sum_{j,k=1}^N A_{ijk}\alpha_j\beta_k \right)_x + \left(2hv_m\beta_i + h \sum_{j,k=1}^N A_{ijk}\beta_j\beta_k \right)_y \\
 = v_mD_i - \sum_{j,k=1}^N B_{ijk}D_j\beta_k - (2i+1)\frac{\nu}{\lambda} \left(v_m + \sum_{j=1}^N \left(1 + \frac{\lambda}{h}C_{ij} \right) \beta_j \right)
 \end{aligned}$$

Example Systems

1D model with h_b constant, $e_x = e_y = 0$, and $e_z = 1$
Constant System

$$\begin{bmatrix} h \\ hu_m \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 \end{bmatrix}_x = -\frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m \end{bmatrix}$$

Flux Jacobian Eigenvalues, $u_m \pm \sqrt{gh}$
Linear System, $\tilde{u} = u_m + \alpha_1 \phi_1$

$$\begin{bmatrix} h \\ hu_m \\ h\alpha_1 \end{bmatrix}_t + \begin{bmatrix} hu_m \\ hu_m^2 + \frac{1}{2}gh^2 + \frac{1}{3}h\alpha_1^2 \\ 2hu_m\alpha_1 \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu_m \\ h\alpha_1 \end{bmatrix}_x - \mathbf{s}$$

$$Q = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & u_m \end{bmatrix} \quad \mathbf{s} = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u_m + \alpha_1 \\ 3(u_m + \alpha_1 + 4\frac{\lambda}{h}\alpha_1) \end{bmatrix}$$

Quasilinear Matrix Eigenvalues, $u_m \pm \sqrt{gh + \alpha_1^2}$, u_m

Example Systems

1 dimensional with h_b constant, $e_x = e_y = 0$, and $e_z = 1$
 Quadratic Vertical Profile, $\tilde{u} = u + \alpha_1 \phi_1 + \alpha_2 \phi_2$

$$\begin{bmatrix} h \\ hu \\ h\alpha_1 \\ h\alpha_2 \end{bmatrix}_t + \begin{bmatrix} hu \\ hu^2 + \frac{1}{2}gh^2 + \frac{1}{3}h\alpha_1^2 + \frac{1}{5}h\alpha_2^2 \\ 2hu\alpha_1 + \frac{4}{5}h\alpha_1\alpha_2 \\ 2hu\alpha_2 + \frac{2}{3}h\alpha_1^2 + \frac{2}{7}h\alpha_2^2 \end{bmatrix}_x = Q \begin{bmatrix} h \\ hu \\ h\alpha_1 \\ h\alpha_2 \end{bmatrix}_x - \mathbf{s}$$

$$Q = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & u - \frac{\alpha_2}{5} & \frac{\alpha_1}{5} \\ 0 & 0 & \alpha_1 & u + \frac{\alpha_2}{7} \end{bmatrix}, P = \frac{\nu}{\lambda} \begin{bmatrix} 0 \\ u + \alpha_1 + \alpha_2 \\ 3(u + \alpha_1 + \alpha_2 + 4\frac{\lambda}{h}\alpha_1) \\ 5(u + \alpha_1 + \alpha_2 + 12\frac{\lambda}{h}\alpha_2) \end{bmatrix}$$

Quasilinear Matrix Eigenvalues, $u \pm c\sqrt{gh}$

$$c^4 - \frac{10\alpha_2}{7}c^3 - \left(1 + \frac{6\alpha_2^2}{35} + \frac{6\alpha_1^2}{5}\right)c^2 + \left(\frac{22\alpha_2^3}{35} - \frac{6\alpha_2\alpha_1^2}{35} + \frac{10\alpha_2}{7}\right)c - \frac{\alpha_2^4}{35} - \frac{6\alpha_2^2\alpha_1^2}{35} - \frac{3\alpha_2^2}{7} + \frac{\alpha_1^4}{5} + \frac{\alpha_1^2}{5} = 0$$

Nonconservative Products, (Dal Maso, Lefloch, and Murat [2])

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Model Equation

$$\mathbf{q}_t + \nabla \cdot \mathbf{f}(\mathbf{q}) + g_i(\mathbf{q})\mathbf{q}_{x_i} = \mathbf{s}(\mathbf{q}) \quad \text{for } (\mathbf{x}, t) \in \Omega \times [0, T]$$

Traditionally searching for weak solutions, find \mathbf{q} such that

$$\int_0^T \int_{\Omega} (\mathbf{q}_t + \nabla \cdot \mathbf{f}(\mathbf{q}) + g_i(\mathbf{q})\mathbf{q}_{x_i}) v \, d\mathbf{x} \, dt = \int_0^T \int_{\Omega} \mathbf{s}(\mathbf{q}) v \, d\mathbf{x} \, dt$$

for all $v \in C_0^1(\Omega \times [0, T])$

Regularization Paths

Consider Lipschitz continuous paths, $\psi : [0, 1] \times \mathbb{R}^p \times \mathbb{R}^p \rightarrow \mathbb{R}^p$, that satisfy the following properties.

- 1 $\forall \mathbf{q}_L, \mathbf{q}_R \in \mathbb{R}^p$, $\psi(0, \mathbf{q}_L, \mathbf{q}_R) = \mathbf{q}_L$ and $\psi(1, \mathbf{q}_L, \mathbf{q}_R) = \mathbf{q}_R$
- 2 $\exists k > 0$, $\forall \mathbf{q}_L, \mathbf{q}_R \in \mathbb{R}^p$, $\forall s \in [0, 1]$, $\left| \frac{\partial \psi}{\partial s}(s, \mathbf{q}_L, \mathbf{q}_R) \right| \leq k |\mathbf{q}_L - \mathbf{q}_R|$
elementwise
- 3 $\exists k > 0$, $\forall \mathbf{q}_L, \mathbf{q}_R, \mathbf{u}_L, \mathbf{u}_R \in \mathbb{R}^p$, $\forall s \in [0, 1]$, elementwise

$$\left| \frac{\partial \psi}{\partial s}(s, \mathbf{q}_L, \mathbf{q}_R) - \frac{\partial \psi}{\partial s}(s, \mathbf{u}_L, \mathbf{u}_R) \right| \leq k(|\mathbf{q}_L - \mathbf{u}_L| + |\mathbf{q}_R - \mathbf{u}_R|)$$

Let $u = u_0 + H(x - x_0)(u_1 - u_0)$, then regularize

$$u^\varepsilon(x) = \begin{cases} u_0 & x < x_0 - \varepsilon \\ \psi\left(\frac{x - x_0 + \varepsilon}{2\varepsilon}, u_0, u_1\right) & x_0 - \varepsilon < x < x_0 + \varepsilon \\ u_1 & x > x_0 + \varepsilon \end{cases}$$

Nonconservative Product Definition

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Let $\mathbf{q} \in BV([a, b] \rightarrow \mathbb{R}^p)$ and $g \in C^1(\mathbb{R}^p \rightarrow \mathbb{R}^p \times \mathbb{R}^p)$, then μ is a Borel measure.

1 If q is continuous on a Borel set $B \subset [a, b]$, then

$$\mu(B) = \int_B g(\mathbf{q}) \frac{d\mathbf{q}}{dx} dx$$

2 If q is discontinuous at a point $x_0 \in [a, b]$, then

$$\mu(x_0) = \int_0^1 g(\psi(s; \mathbf{q}(x_0^-), \mathbf{q}(x_0^+))) \frac{\partial \psi}{\partial s}(s; \mathbf{q}(x_0^-), \mathbf{q}(x_0^+)) ds$$

Define

$$\mu = \left[g(\mathbf{q}) \frac{d\mathbf{q}}{dx} \right]_{\psi}$$

Nonconservative Products

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

If there exists $\mathbf{f}(\mathbf{q})$ such that $\mathbf{f}'(\mathbf{q}) = \mathbf{g}(\mathbf{q})$, then

$$[\mathbf{g}(\mathbf{q})\mathbf{q}_x]_\psi = \mathbf{f}(\mathbf{q})_x$$

or

$$\int_0^1 \mathbf{f}'(\psi(s, \mathbf{q}_L, \mathbf{q}_R)) \frac{\partial \psi}{\partial s}(s, \mathbf{q}_L, \mathbf{q}_R) ds = \mathbf{f}(\mathbf{q}_R) - \mathbf{f}(\mathbf{q}_L)$$

Find weak solution \mathbf{q} such that

$$\begin{aligned} & \int_0^T \int_{\Omega} \mathbf{q} \mathbf{v}_t \, d\mathbf{x} \, dt + \int_0^T \int_{\Omega} \mathbf{f}(\mathbf{q}) \nabla \cdot \mathbf{v} \, d\mathbf{x} \, dt \\ & + \int_0^T \int_{\Omega} [\mathbf{g}_i(\mathbf{q})\mathbf{q}_{x_i}]_\psi \mathbf{v} \, d\mathbf{x} \, dt = \int_0^T \int_{\Omega} \mathbf{s}(\mathbf{q}) \mathbf{v} \, d\mathbf{x} \, dt \end{aligned}$$

for all $\mathbf{v} \in C_0^1(\Omega \times [0, T])$.

Nonconservative DG Formulation

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

**Nonconservative
DG Formulation**

Results

References

Manufactured Solution

Shallow Water Equations, constant vertical velocity profile

1st Order			2nd Order		3rd Order	
n	error	order	error	order	error	order
20	0.226	—	8.57×10^{-3}	—	1.67×10^{-4}	—
40	0.117	0.96	2.17×10^{-3}	1.98	2.07×10^{-5}	3.02
80	0.058	1.00	5.40×10^{-4}	2.01	2.57×10^{-6}	3.01
160	0.028	1.06	1.35×10^{-4}	2.00	3.21×10^{-7}	3.00
320	0.014	0.99	3.37×10^{-5}	2.00	4.01×10^{-8}	3.00

4th Order			5th Order	
n	error	order	error	order
20	3.172×10^{-6}	—	7.606×10^{-8}	0.00
40	1.982×10^{-7}	4.00	2.380×10^{-9}	5.00
80	1.240×10^{-8}	4.00	7.713×10^{-11}	4.95
160	7.755×10^{-10}	4.00	4.035×10^{-11}	0.93
320	4.849×10^{-11}	4.00	8.085×10^{-11}	-1.00

Manufactured Solution

One moment, linear vertical velocity profile

	1st Order		2nd Order		3rd Order	
<i>n</i>	error	order	error	order	error	order
20	2.53×10^{-1}	—	9.97×10^{-3}	—	1.71×10^{-3}	—
40	1.30×10^{-1}	0.96	2.52×10^{-3}	1.98	3.85×10^{-4}	2.15
80	6.47×10^{-2}	1.00	6.28×10^{-4}	2.00	6.13×10^{-5}	2.65
160	3.13×10^{-2}	1.05	1.57×10^{-4}	2.00	9.09×10^{-6}	2.75
320	1.58×10^{-2}	0.99	3.92×10^{-5}	2.00	1.73×10^{-6}	2.39

	4th Order		5th Order	
<i>n</i>	error	order	error	order
20	1.14×10^{-4}	—	2.68×10^{-5}	—
40	1.74×10^{-5}	2.72	8.01×10^{-7}	5.06
80	7.50×10^{-7}	4.53	1.53×10^{-8}	5.71
160	1.25×10^{-7}	2.59	4.04×10^{-10}	5.25
320	8.79×10^{-9}	3.83	8.40×10^{-11}	2.27

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Manufactured Solution

Two moments, quadratic vertical velocity profile

1st Order			2nd Order		3rd Order	
n	error	order	error	order	error	order
20	2.778×10^{-1}	—	1.141×10^{-2}	—	5.350×10^{-3}	—
40	1.424×10^{-1}	0.96	2.884×10^{-3}	1.98	6.466×10^{-4}	3.05
80	7.121×10^{-2}	1.00	7.191×10^{-4}	2.00	7.836×10^{-5}	3.04
160	3.454×10^{-2}	1.04	1.797×10^{-4}	2.00	1.270×10^{-5}	2.63
320	1.740×10^{-2}	0.99	4.493×10^{-5}	2.00	2.546×10^{-6}	2.32

4th Order			5th Order	
n	error	order	error	order
20	3.688×10^{-4}	—	5.194×10^{-5}	—
40	2.461×10^{-5}	3.91	1.121×10^{-6}	5.53
80	1.403×10^{-6}	4.13	1.934×10^{-8}	5.86
160	1.144×10^{-7}	3.62	5.859×10^{-10}	5.04
320	1.092×10^{-8}	3.39	8.791×10^{-11}	2.74

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Manufactured Solution

Three moments, cubic vertical velocity profile

1st Order			2nd Order		3rd Order	
n	error	order	error	order	error	order
20	3.024×10^{-1}	—	1.300×10^{-2}	—	7.015×10^{-3}	—
40	1.556×10^{-1}	0.96	3.283×10^{-3}	1.99	6.992×10^{-4}	3.33
80	7.808×10^{-2}	0.99	8.188×10^{-4}	2.00	1.183×10^{-4}	2.56
160	3.802×10^{-2}	1.04	2.046×10^{-4}	2.00	2.545×10^{-5}	2.22
320	1.916×10^{-2}	0.99	5.117×10^{-5}	2.00	5.110×10^{-6}	2.32

4th Order			5th Order	
n	error	order	error	order
20	3.167×10^{-4}	—	5.571×10^{-5}	—
40	2.384×10^{-5}	3.73	1.099×10^{-6}	5.66
80	2.509×10^{-6}	3.25	2.639×10^{-8}	5.38
160	3.168×10^{-7}	2.99	1.371×10^{-9}	4.27
320	4.675×10^{-8}	2.76	1.171×10^{-10}	3.55

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Effect of Higher Moments

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

Conclusions

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References



Future Work

- Two Dimensional Meshes
- Icosahedral Spherical Mesh
- Shallow Water test cases on the sphere

Bibliography I

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

- [1] Bernardo Cockburn and Chi-Wang Shu. “The Runge–Kutta discontinuous Galerkin method for conservation laws V: multidimensional systems”. In: *Journal of Computational Physics* 141.2 (1998), pp. 199–224.
- [2] Gianni Dal Maso, Philippe G Lefloch, and François Murat. “Definition and weak stability of nonconservative products”. In: *Journal de mathématiques pures et appliquées* 74.6 (1995), pp. 483–548.
- [3] Julia Kowalski and Manuel Torrilhon. “Moment Approximations and Model Cascades for Shallow Flow”. In: *arXiv preprint arXiv:1801.00046* (2017).
- [4] Philippe Le Floch. “Shock waves for nonlinear hyperbolic systems in nonconservative form”. In: (1989).
- [5] Randall J LeVeque et al. *Finite volume methods for hyperbolic problems*. Vol. 31. Cambridge university press, 2002.

Bibliography II

Caleb Logemann,
James
Rossmanith

Generalized
Shallow Water
Equations

Nonconservative
Products

Nonconservative
DG Formulation

Results

References

- [6] Sander Rhebergen, Onno Bokhove, and Jaap JW van der Vegt. “Discontinuous Galerkin finite element methods for hyperbolic nonconservative partial differential equations”. In: *Journal of Computational Physics* 227.3 (2008), pp. 1887–1922.
- [7] J.A. Rossmanith. DOGPack. Available from <http://www.dogpack-code.org/>.
- [8] James A Rossmanith. “A wave propagation method for hyperbolic systems on the sphere”. In: *Journal of Computational Physics* 213.2 (2006), pp. 629–658.
- [9] David L Williamson et al. “A standard test set for numerical approximations to the shallow water equations in spherical geometry”. In: *Journal of Computational Physics* 102.1 (1992), pp. 211–224.