figures/sheffieldlogo.png

How are mass extinctions shaping biodiversity?

Confirmation Review

Caleb Scutt
Supervisors: Thomas Guillerme (UoS), Gavin Thomas (UoS) & Natalie Cooper (NHM)
Advisor: Karl Evans (UoS)

July 2025

Contents

Part I

Abstract

While mass extinctions are recognised as significant macroevolutionary events, their effects on morphological diversity (disparity) remain poorly understood. The impact of the Cretaceous–Palaeogene (K–Pg) mass extinction on eutherian mammal disparity serves as a particularly contentious case study. Conflicting conclusions regarding the magnitude and direction of disparity changes following this extinction event highlight fundamental methodological challenges in the disparity analysis pipeline. This PhD project aims to address these challenges by developing novel methodological approaches better suited to inferring changes in disparity through time. One central aim is to assess how the handling of uncertainty in ancestral state estimation affects downstream disparity inferences. This will be tested across varying models of trait evolution, different fossil sampling levels, and both selective and non-selective extinction scenarios to identify the best way of handling uncertainty in ancestral state estimation. In addition, a new statistical framework, adapted from the Before-After-Control-Impact design in ecology, will be developed to detect non-random changes in disparity through time. This framework will account for trend and uncertainty, making it possible to distinguish true disparity shifts from stochastic artefacts. In order to infer the macroevolutionary patterns of eutherian mammals, it is necessary to have a dated phylogeny. A total-evidence fossilised birth–death approach will be used to estimate the trees under competing diversification models. This will allow exploration of how divergence timing affects disparity trajectories. Integration of these three projects will culminate in a disparity analysis pipeline capable of processing multiple tree distributions and propagating uncertainty to generate probabilistic estimates of disparity change. This pipeline can then be utilised to infer whether the K-Pg mass extinction had an effect on the morphological diversification of eutherian mammals. These advances will allow a critical re-evaluation of existing hypotheses about early placental evolution, as well as advance the methodological underpinnings of disparity analyses.

Part II

Background Review

1 Morphology, Disparity and Extinction

1.1 Mass extinctions

Extinction is a fundamental evolutionary process that, alongside speciation, continuously shapes patterns of biodiversity through deep time; over 99% of all species that have ever lived are now extinct (**?**). However, extinction rates are not constant and occur in pulses across background taxonomic turnover (**?**). *Mass extinctions* are specific events that wipe out over 75% of species and have happened five times over the last 500 million years (**?**). On a macroevolutionary scale, mass extinctions mark the most dramatic species composition transitions in Earth history (**?**). Entire branches are trimmed from the tree of life, while the evolutionary trajectories of survivors are altered in the wake of a transformed global biota (**?**).

1.2 Rebound dynamics

The aftermath of this drastic species turnover is typically described as 'recovery', yet **?** describe this as a misnomer, since this implies a return to the prior pre-extinction state. Instead, biodiversity rebounds are multifaceted and the response from the survivors of a mass extinction event may take several forms, transforming the post-extinction biota (**?**). For example, trends observed in different clades pre-mass extinction may continue unperturbed, including escalations (**?**) and declines (**?**). Alternatively, the aftermath of a mass extinction can trigger diversification as surviving clades are offered ecological opportunity, filling the vacant niches left by extinct lineages.

Rebound dynamics can differ by the metric of diversity considered. Intuitively, a diversifying clade would exhibit simultaneous taxonomic and phenotypic diversification, following a pattern of early high rates that decrease through time as niches become saturated (**?**). However, empirical evidence often finds that peak taxonomic and morphological diversity are reached at different points in a clade's history (**????**). This discordance has been observed across mass extinction events, such as in graptolites across the end-Ordovician extinction (**?**), or therapsids across the Permian-Triassic extinction (**?**). It has been suggested that functional diversity may be density-independent across mass extinction events (**?**). Therefore, understanding the rebound dynamics of clades based solely on taxonomic diversity metrics may not encapsulate the full complexity of rebound dynamics (**?**).

1.3 Disparity

This phenotypic diversity, often called *disparity*, has been widely used to investigate patterns of evolution in clades (**?**). Disparity refers to the extent of morphological diversity among taxa, typically quantified as the distribution or spread of those taxa within a morphospace — a multidimensional mathematical space configured by morphological similarity (**?**). Changes in morphospace through time are often closely tied to shifts in ecological diversity (**??**). For example, expansion in morphospace may indicate a clade expanding its niche breadth during an adaptive radiation (**???**). Alternatively, a clade may move position within the morphospace, reflecting shifts in ecological niche (**?**).

1.4 Extinction selectivity

Mass extinctions pose numerous scenarios of disparity (**??**). In adaptive scenarios, disparity may increase in clades that diversify during the period of ecological opportunity following a mass extinction due to morphospace expansion. Extinction can also be random or selective based on specific traits, or regions of the morphospace, that go extinct. **?** proposed several patterns to describe different selection processes: lateral (asymmetric selection that trims one side of the morphospace), marginal (symmetric selection that removes peripheral taxa) and non-selective ('field of bullets' / independent of the variables defining morphospace).

figures/extinctionscenarios.png

Figure 1: Three scenarios that may occur within a clade across a mass extinction (denoted by dashed line). a) adaptive radiation. b) selective extinction. c) non-selective extinction. Adapted from **?**.

1.5 Disparity metric

These different types of extinction events can be distinguished from each other by comparing changes in different disparity metrics. For example, the sum of variances may remain relatively unchanged during a random extinction, since no systematic reduction in trait variance has occurred (**??**). In contrast, a reduction in sum of variances will likely be apparent in a marginal selective extinction, where the pruning of

peripheral taxa has led to a contraction in variance. **?** highlight the flaws in using a single disparity metric for assessing morphospace changes, since changes can occur in size, position and density (**?**). Studies that have implemented multiple metrics have been able to identify nuanced changes in morphospace (**?**).

2 Eutherian disparity across K-Pg Mass Extinction

2.1 Suppression hypothesis

The most recent of the 'Big Five' mass extinctions occurred 66 million years ago (Ma) at the Cretaceous-Palaeogene (K-Pg) boundary, primarily known for the demise of the non-avian dinosaurs (**???**). The effect of the K-Pg mass extinction on eutherian mammals (placentals and extinct relatives) is one of the most well-studied yet contentious inquiries in macroevolution. Eutherian mammals prior to the K-Pg boundary have been widely considered depauperate and limited in ecological variation, which contrasts the diversity seen in extant placentals. This traditional view, known as the *suppression hypothesis*, follows that the extinction of non-avian dinosaurs released eutherian mammals from constraints of predation and competition, stimulating a radiation (**????**). The most significant morphological evidence for the suppression hypothesis in the fossil record concerns a shift in body size disparity (**?**). Mesozoic mammals generally weighed less than 100 g and rarely over 5 kg (**?**). The maximum body mass of mammals increased exponentially and by roughly three orders of magnitude by the late Eocene (**????**). It is thought that this increase is indicative of a wider expansion of niche breadth as mammals filled the vacant niches left by large dinosaurs.

2.2 Early rise hypothesis

While the suppression hypothesis is strongly supported by studies that have investigated changes in body size (**??**), body size is limited as a metric for ecological variation since it can be influenced by several factors that may reflect stochastic rather than adaptive changes (**?**). Recent studies instead focusing on morpho-functional traits (intrinsically linked to ecological function), such as jaw and dentition morphology, have instead suggested that eutherian diversity was attained through successive waves of diversification, starting far earlier than the K-Pg mass extinction (**?**). The *early rise hypothesis* suggests an ecological diversification that occurred concurrently with the diversification of angiosperms and insects during and after the Cretaceous Terrestrial Revolution (125–80 Ma) (**??**) (**??**). Using morphometric analyses of molar shape **?** found that eutherian disparity began increasing during the Late Cretaceous. **?** also found a similar pattern, as well as a decrease in molar disparity across K-Pg suggesting selective extinction. Ecological variation within Mesozoic mammals was not limited to diet; diversity in locomotor styles was reached as far back as the Jurassic in early eutherians (**?**).

2.3 Delayed rise hypothesis

Other studies have pointed towards the Paleocene-Eocene Thermal Maximum, an episode of elevated global temperatures 56 Ma that widened the latitudinal range of angiosperm-dominated forest, which in turn spurred the dispersal and radiation of eutherians (**??**) (**?**). **?** found that eutherian mammals had the most significant expansion in jaw morphospace at the onset of the Eocene. Similarly, **?** investigated endocranial volume in Cenozoic placentals and also found a delayed radiation in diversity, highlighting increases in body size during the Paleocene preceded an expansion in brain size in the Eocene, which became necessary as competition intensified in an increasingly mammal-dominated biota. While these alternative hypotheses do not necessarily refute the influence of the K-Pg mass extinction, they instead suggest the extinction accentuated evolutionary trajectories already underway.

`figures/grossnickleradiations.jpg`

Figure 2: The contrasting ecological diversification triggers and placental origin hypotheses. Adapted from **?**.

3 Disparity Analysis Methods

The conflicting findings of eutherian disparity across the K-Pg boundary raise the question: why do disparity analyses aiming to answer the same question arrive at different conclusions? Addressing this requires a critical look at the methodological underpinnings of disparity analyses, specifically phylogeny, ancestral state estimation and tests for change in disparity through time.

figures/simplepipeline.png

Figure 3: Disparity pipeline (N.B. this is specifically for discrete characters. Continuous characters would bypass the distance matrix after ASE or without ASE). As shown there are several routes to disparity, including bypassing ancestral state estimation (ASE), pre-ordination ASE, post-ordination ASE

4 Phylogenetic hypotheses

The origin and tempo of lineage diversification in placental mammals has long been a point of contention, meaning there is no undisputed phylogenetic topology for eutherian mammals. There are currently three

main phylogenetic models (**??**).

Table 1: Summary of major models for the origin and diversification of placental mammals.

Model	**Placental origin**	**Evidence**	**References**
Hard explosive	66 Ma (K-Pg boundary)	Palaeontological: no crown placental fossils before K-Pg	**????**
Soft explosive	~ 80 Ma	Molecular: removed fossil calibrations for large (slow evolving) taxa	**??**
Long fuse	$\sim 120 - 80$ Ma	Molecular: genome wide data	**?????**

Given the absence of a well-resolved eutherian phylogeny, studies investigating eutherian disparity have opted to time-bin disparity by stratigraphy (**????**), which inherently assumes a punctuated equilibrium model of morphological evolution. This assumption can be particularly problematic when assessing disparity across mass extinction events (**?**). Stratigraphic time binning, which aggregates data over broad intervals, risks obscuring critical, transient shifts in disparity that may occur within shorter durations following the extinction. Incorporating a phylogeny means a time-slicing method can be used (**?**), which removes the punctuated assumption and can reveal more temporally refined patterns.

5 Ancestral state estimation

The poor resolution of the fossil record is an inevitable limitation affecting disparity analyses, since only a small biased portion of a clade's evolutionary history is preserved. The incorporation of a phylogeny together with trait data enables a switch from simply observing a pattern (points in morphospace) to inferring a process through time, through the use of ancestral state estimation (**?**).

5.1 Application of ancestral state estimation

Ancestral state estimation is the broad method of estimating trait value(s) in a set of common ancestors given a set of observed data for tips on a phylogenetic tree (**??**). Intuitively, ancestral state estimation is appealing for those studying deep-time patterns of disparity. Incorporating a phylogeny into disparity analyses will unavoidably introduce unsampled 'ghost lineages' (**?**). A ghost lineage is the hypothetical ancestor that is not observed in the fossil record, but can be inferred to have existed due to the presence of other fossils/extant taxa. This can be best explained as the branch on a phylogeny between two tips (**??**).

figures/Ichthyosauria_phylogeny.jpg

Figure 4: Phylogeny of ichthyosaurs showing observed stratigraphic range for each species (thick lines) and inferred ghost lineages (thin lines) (**?**).

Ancestral state estimation can employed as a way of inferring the traits of these hypothetical species based on the traits of their descendents. This fills in the gaps in the fossil record and alleviates the bias associated with uneven sampling across the phylogenetic tree (**??**).

5.1.1 Likelihood methods

Likelihood methods for ancestral state estimation are based on a unifying statistical framework that treats the internal node states as parameters to be estimated (**?**). The central tenet of the approach is the likelihood function, which integrates information from the topology, which reflects the evolutionary relationships between taxa, and the model of trait evolution, which specifies how traits evolve. By systematically exploring the parameter space through numerical optimisation, the method searches for the ancestral states that maximise the likelihood function. The same general likelihood framework underpins estimation of continuous and discrete characters.

5.1.2 Continuous characters

Continuous trait evolution is typically modelled under Brownian motion (BM) (**??**). BM models trait evolution as a directionless, stochastic process under which continuous characters undergo changes along the

branches of a phylogeny. These evolutionary changes come from a Gaussian distribution with a mean of 0 and variance proportional to the product of the rate of evolution and time elapsed ($\sigma^2 t$). Therefore, the expected amount of trait divergence between any two taxa is proportional to the amount of shared evolutionary history according to the phylogeny.

Under BM, the vector ($\mathbf{x}$) of observed tip trait values follows a multivariate normal distribution (MVN) and has an expected value that is equal to the root state x_0.

$$\boldsymbol{x} \sim \mathrm{MVN}(x_0,\, \sigma^2 \boldsymbol{C})$$

$\boldsymbol{C}$ is a variance-covariance matrix for the shared ancestry of the taxa, defined by the tree height from the root to each tip and the tree height above the root of the most recent common ancestor shared by each pair of tips.

When estimating ancestral states, the $\boldsymbol{C}$ matrix is expanded to include the shared history of the internal nodes. The evolutionary rate, σ^2, and root state x_0, are unknown parameters that can be estimated using Maximum Likelihood to give the multivariate normal distribution. The trait values at internal nodes are then calculated as the conditional expectations within this distribution. This means each node's estimated trait value is effectively a weighted average of the tip values, with the weights determined by the amount of shared evolutionary history (as defined by $\boldsymbol{C}$).

5.1.3 Discrete characters

Ancestral state estimation of discrete characters are modelled using a continuous-time Markov chain process, known as the Mk model (**??**). This process is specified by a $k \times k$ matrix (k being the number of possible states), $\mathbf{Q}$, which gives the instantaneous transition rates (q) between states for each character.

$$\mathbf{Q} = \begin{bmatrix} -q_{0,1} & q_{0,1} \\ q_{1,0} & -q_{1,0} \end{bmatrix}$$

The $\mathbf{Q}$ matrix is estimated using Maximum Likelihood; the value of $\mathbf{Q}$ that maximises the probability of the observed data given the tree topology. The matrix exponentiation of $\mathbf{Q}$ converts the instantaneous rates into probabilities of transitioning between states over a given time t:

$$\mathbf{P} = \exp(\mathbf{Q}t)$$

Global marginal estimation fixes the value of $\mathbf{Q}$ and calculates the probability of each state at each node, given the observed data at the tips. These likelihoods are rescaled by their sum so that they add up to 1.0, and so represent an empirical Bayes posterior probability (**?**).

5.2 Challenges in ancestral state estimation

"Character state reconstructions can provide a powerful mechanism for studying many facets of the evolutionary process. However, the zeal with which these techniques are sometimes advocated belies the complexity of the problem" (**?**).

5.2.1 Uncertainty in estimates

A critical issue is that estimates tend to have broad uncertainty (**?**). This uncertainty typically increases towards root, where information from observed tip data becomes increasingly diffuse (**?**). For discrete characters, this uncertainty manifests as inconclusive scaled likelihoods, i.e., '0' = 0.51, '1' = 0.49. For continuous characters, uncertainty is wide confidence intervals.

How this uncertainty in estimations should be dealt with is equivocal. Despite the encouragement for uncertainty to be carried forward in comparative analyses (**?**), most computational implementations of ancestral state estimation return point estimates. For continuous characters, this will be the median value of the confidence interval and for discrete characters, it will be the most probable state according to a scaled likelihood threshold. By ignoring uncertainty, point estimates can lead to overconfidence in results and erroneous inferences in downstream analyses (such as down the disparity pipeline), all while obscuring potentially 'accurate' uncertainty (**??**).

For discrete characters, there is a level of subjectivity in how to collapse the states into the most probable based on their scaled likelihood. It is common to have a threshold implementation in ancestral state estimation software (state probabilities above a set value are collapsed to a point estimate), or left as uncertain if this threshold is not met, which is denoted as '0/1'. However, these thresholds are often arbitrarily set with no guidance of what is the 'correct' value (**??**). Higher thresholds are likely too conservative and so too many states are left uncertain, perhaps missing true signals, whereas lower thresholds may be too permissive and can lead to erroneous inferences.

5.2.2 Sensitivity to fossil sampling

Given that uncertainty is often high in ancestral state estimation, how then can estimates be improved? **?** demonstrates that ancestral state estimation is *statistically inconsistent*, meaning that even if the sample size of extant tips is infinite, estimates are not expected to converge on the true value. Instead, the inclusion of fossils can improve estimates, due to the shorter evolutionary path to the root of the tree (**??**).

By extension, inclusion of fossils can greatly influence inferences into disparity. Extant-only datasets often miss early rapid morphological diversification signals (**?**), whereas analyses incorporating fossil data often finds the reverse (**?**) For example, **?** found that analyses of bird disparity were underestimated when using ancestral state estimation of extant-only datasets.

5.2.3 Model dependency

Another key challenge is the sensitivity to violations of model assumptions. For continuous characters, this is particularly pertinent when traits have evolved under non-BM (**?**). When trait evolution deviates from BM – due to directional trends, bounded evolution, or regime shifts – ancestral state estimation can return biased estimations.

Both **?** and **?** were unable to recover accurate ancestral estimates of body size in planktonic foraminifera and mammals, respectively, due to a trend of increase through time. Trends mean a moving average through time, which violates the assumption of zero-mean change under BM. Similarly, issues estimating the evolutionary rate may arise when the trait data evolved under an Ornstein-Uhlenbeck (OU) model, whereby trait data evolves under stabilising selection (**?**). It is common for regime shifts to occur at mass extinction boundaries (**?**). For example, **?** found that mammal body size was relatively constrained (OU) up until the K-Pg mass extinction which then allowed a release of constraint and BM evolution. Such shifts could affect estimation accuracy of evolutionary rate and thus state estimates.

As discussed previously, fossil sampling is crucial for accurate estimation, especially when model assumptions are violated (**?**). Fossils assist in estimating and constraining model parameters, as well as distinguishing different evolutionary processes (**?**). For example, when estimating states under an OU model, fossils are necessary for estimating the primary optimum. Similarly, fossils would be necessary to detect a trend, such as Cope's rule that body size increases through time (**?**). A similar logic applies to discrete characters; **?** found that inclusion of fossils improved estimate accuracy, irrespective of model misspecification.

6 Statistical tests in disparity

Given a calculation of disparity, it is necessary for this value to be statistically tested in the context of the research question. A popular choice is to carry out a PERMANOVA, which can be used to assess whether two groups share the same centroid and dispersion in the morphospace (**???**). Alternatively, studies may opt to investigate change in disparity, for example through the use of a t test (**???**). Therefore, a t-test may be preferred if investigating a nuanced change in morphospace that PERMANOVA cannot capture, such as change in variance.

The way these tests are implemented depends on the research question, but questions tend to relate to differences between groups (between clades for example) or through time (across time bins). However, using a single 'average change' statistic to compare disparity through time has limitations, notably that it ignores directional trends. Discounting trends in evolution may implicitly assume a punctuated model of evolution, where abrupt phenotypic shifts are interpreted as adaptive but are actually a reflection of stochastic processes (**??**).

There have been several methods already developed for identifying non-random shifts in disparity. A common approach is to implement a null model to see if the empirical data deviates (**??**). For example, the global

rank envelope test implemented by **?** identifies non-random shifts as when the empirical Disparity-Through-Time curve (see **?**) falls within the most extreme 5% ranked curves of a null Brownian model. Similarly, **?** made use of simulations of discrete trait data as a null model based on the zero-force evolutionary law (the tendency for diversity to increase through time) (**?**), to identify points at which disparity is greater/less than expected. However, a framework is needed that allows a statistical test to be applied over large numbers of time series, when inspecting individual time series becomes impractical.

Part III

Thesis plan

7 Aims and objectives

Knowledge gap

The overarching aim of this PhD project is to understand the long term effect of mass extinctions by developping new state-of-the-art disparity methods. To achieve this I will be revisiting the critical question whether the K-PG mass extinction had an effect on eutherian disparity while focusing on improving the methodology of the disparity pipeline.

Research questions

- How robust are disparity analyses to uncertainty in ancestral state estimations?
- How can trend be accounted for when identifying non-random shifts in disparity?
- How do both selective and non-selective extinction events differ in the ability to recover accurate ancestral state estimations?
- Which of the three ecological hypotheses (**??**) for the diversification of eutherian mammals is most strongly supported by disparity analyses?

Research objectives

- Using a total-evidence fossilised birth-death phylogenetic inference approach to estimate long fuse, soft explosive and hard explosive tree distributions.
- Assess the accuracy of ancestral states estimations accuracy on disparity analyses across: 1) a range of models of trait evolution, 2) fossil sampling levels, 3) mass extinctions with different intensities and selectivities, 4) and disparity metrics.
- Develop a statistical framework, built on the BACI approach in ecology, that can identify immediate and long term trend changes in disparity at a specified time point.

- Use these methodological advances to build a pipeline capable of handling multiple tree distributions and disparity parameters that outputs a distribution of 'effect values' relating to K-Pg and eutherian disparity.

8 Chapter One: Eutherian phylogenies

Chapter aim: To estimate the phylogeny of eutherian mammals, using various calibrations for the placental node in order to recover the three common placental origin hypotheses: long fuse, soft explosive and hard explosive (**??**). The rationale of this chapter is so that these three tree distributions can be used in chapter four (**??**).

Research questions:

- When did major lineages of eutherians diversify?
- When did major shifts in evolutionary rates occur in eutherians?

Research objectives:

- I will use both molecular and morphological data for a total-evidence approach, which involves concatenating both molecular and morphological alignments into a single supermatrix
- The molecular and morphological matrices have been previously collected by my supervisor in a previous project
- Bayesian analysis will use RevBayes.
- I will implement different calibration strategies at the placental node to generate three alternative models: a long fuse model (placental node $\sim$ 100 Ma), a soft explosive scenario (placental node $\sim$ 80 Ma), and a hard explosive scenario (placental node 66 Ma).
- I will specify a Fossilized Birth-Death model in RevBayes to incorporate fossil tip ages and estimate divergence times.
- I will assess convergence and summarize the posterior distributions to obtain time-calibrated phylogenetic trees that reflect each calibration scenario.
- I will use RevGadgets to summarise the trees and plot figures.

9 Chapter Two: What is the effect of ancestral state estimations on disparity?

Chapter aim: To investigate how different approaches for treating uncertainties in ancestral state estimations affect estimates of morphological disparity under varying levels of fossil sampling. To expand this

framework to understand how random and selective mass extinction events can confound ancestral state and disparity estimation accuracy.

Research questions:

- Which method of treating uncertainties recovers the most accurate ancestral state estimations?
- How does estimation accuracy change across different fossil sampling levels?
- How does estimation accuracy of disparity differ to accuracy of ancestral state estimation?
- What are the effects of mass extinction events on ancestral state estimations?
- How do random vs selective extinction events differ in their effect on ancestral state estimations?

Research objectives:

- Develop a simulation procedure that generates a distribution of 'true' evolutionary histories (i.e., phylogeny and trait data), with both discrete and continuous trait data.
- Remove varying levels of fossil tips from these 'true' phylogenies to simulate different fossil sampling levels.
- Run ancestral state estimation on these pruned trees/matrices with three different ways of treating uncertainty in estimates.
- Compare the raw distance between true vs estimated matrices.
- Compare the difference in disparity values between true vs estimated matrices.
- Run this procedure, but incorporating a mass extinction event into the simulations. These mass extinction events will have varying intensities and selectivities.
- Benchmark the findings with empirical data - likely using a dataset with exceptional fossil preservation, such as Foraminifera.

10 Chapter Three: Measuring changes in trends in disparity through time

Chapter aim: To develop a framework that can distinguish between random and non-random shifts in disparity through time, accounting for trend.

Research questions:

- How can traditional statistical tests, such as changes in average (t tests), be misleading in inferences into temporal change in disparity?

- How to implement a BACI (Before After Control Intervention) approach into disparity through time? The BACI framework, as laid out from an ecological perspective in **?**, aims to assess how interventions (which can be loosely defined as major climatic or geological events in this context) have an effect on a variable across a time series, given an estimate of the counterfactual.
- How to generate valid simulated control data, i.e., null models, for different specific questions?
- How to deal with time lags?

Research objectives:

- Building on the BACI framework laid out by **?** I will develop a method to quantify shifts in disparity across a mass extinction (or any other fixed point in time that is suspected of having an effect on disparity).
- This framework will consider the mass extinction event to be an Intervention.
- Piecewise regression will be fit to data pre-extinction (Before) and post-extinction (After)
- Simulated data will be used as the null model (Control). The model parameters for replicating the pre-extinction trajectory will be estimated and simulations built on those parameters.
- BACI contrasts, i.e., $(\mathrm{A_I} - \mathrm{B_I}) - (\mathrm{A_C} - \mathrm{B_C})$ can be used as a quantitative measure of the effect of the Intervention (mass extinction).
- Implement a modular approach to the function that allows for time lags to be accounted for. This would require the function to be able to detect breakpoints within a certain timeframe of the Intervention.

figures/baci.png

Figure 5: Blue arrows indicate positive change and red indicate negative change. The intervention effect can be interpreted either by change in average (A, B), change in trend (C, D) and/or an immediate change (E, F). BACI comparisons show the BACI contrast. For example, in panel D, there is a positive change in trend. Figure from **?**.

11 Chapter Four: Eutherian disparity over K-Pg mass extinction

Chapter aim: Incorporate the phylogenies from chapter one, the ancestral state method from chapter two and the test for non-random shifts in disparity from chapter two to develop a pipeline that can test whether the K-Pg mass extinction had an effect eutherian disparity.

Research questions:

- Is there a shift in trend of disparity at the K-Pg boundary?
- Does this shift depend on the phylogenetic topology used?
- Do patterns differ between different disparity metrics?

Research objectives:

- Use a dataset of discrete morphofunctional characters of jaw morphology and dentition, collected by **?**.
- Develop a pipeline that iterates through different models and metrics: tree topology (long fuse/soft explosive/hard explosive), time slicing model and disparity metric.
- Use the BACI contrast function developed in chapter three to score a value of 'change in disparity' at 66 Ma to obtain a distribution of effects depending on the choice of topology, time-slicing model and disparity metric.

figures/gantt.png

Figure 6: Gantt Chart for PhD timeline

Part IV

Summary of research achievements

Note that this is chapter two findings so far presented here.

12 Methods

12.1 Simulations

Simulations were carried out using `treats` (**?**). Trees were simulated using a modified version of the birth-death process algorithm. The process begins with a single lineage and iteratively selects lineages to grow the tree forward in time. At each step, all living branches are extended by a branch length increment, drawn from an exponential distribution based on the total number of lineages and the combined speciation and extinction rates. Selected lineages then undergo either speciation or extinction based on a probabilistic decision determined by the specified rates. 100 trees were simulated with birth-death parameters as speciation, $\lambda = 1.0$, and extinction, $\mu = 0.7$. Trees were fixed to have 50 extant tips and a range of 50 - 200 fossil tips.

Across each tree, discrete and continuous trait data were simulated. Discrete trait data were simulated using three different Markov chain matrices, representing fast (0.9), medium (0.1) and slow (0.01) transition rates, for 100 characters (85% binary and 15% three-state). Continuous trait data were simulated under Brownian motion (BM) and Ornstein-Uhlenbeck (OU) models, with default parameters and for 50 characters. Trait data were simulated using the function `map.traits`, which assigns both nodes and tips trait values, essentially constructing full hypothetical evolutionary histories across each phylogeny for each trait type.

In order to simulate different levels of fossil sampling, each tree was split into four chronological 'stages' to replicate the stratigraphic nature of the fossil record. From each 'stage', fossil tips were randomly pruned from the tree and corresponding trait matrix at five levels:

- Living: only extant tips are retained
- Low: 5% fossils retained
- Medium: 15% fossils retained
- High: 50% fossils retained
- All: All fossils retained

I aim to modify this method in the future so that earlier stages have lower fossil sampling than more recent stages, to replicate the temporal gradient in completeness seen in the empirical fossil record (**?**).

12.2 Ancestral state estimation

Node states were then removed from the trait matrices leaving just tip states. Ancestral states were estimated using the `multi.ace` function in the `dispRity` package (**?**).

12.2.1 Discrete characters

For discrete state estimation, `multi.ace` is a wrapper for `castor::asr_mk_model`. Ancestral states were estimated using marginal likelihood estimation, whereby each state at each node is assigned a scaled likelihood

(**??**). Symmetrical transition rates were set *a priori* during simulations and so the SYM model was used to avoid model misspecification as a confounding factor on estimation accuracy (**?**). Three different majority-rule approaches were implemented for collapsing the state probabilities into ancestral states. The *strict majority rule* uses a threshold that selects the state with a scaled likelihood equal or greater than a designated value (in this case 0.5). The *relative majority rule* sets a threshold at a scaled likelihood greater than the maximum observed scaled likelihood minus the inverse number of possible states, k

$$\textit{Relative majority threshold} = \max(L_{\text{scaled}}) - \frac{1}{k}$$

The *no majority rule* (sampling approach) returns multiple matrices (x matrices) where ancestral states are probabilistically sampled based on their scaled likelihoods. For each node, the state assigned in each matrix is randomly drawn according to the probability distribution of the scaled likelihoods, meaning states with higher likelihoods are more likely to be selected across the sample of matrices. For example, a node with uncertain estimates (e.g., state 0 = 0.6, state 1 = 0.4) will be assigned state 0 in approximately 60% of the matrices and state 1 in 40% of the matrices. In this case, 100 matrices were returned using the no majority rule.

12.2.2 Continuous characters

Continuous ancestral states were estimated using the BM likelihood method described in **??**. Two approaches were applied to account for uncertainty in estimates. The *point estimate* approach simply returns the median of the 95% confidence interval. Alternatively, the *sampling* approach, whereby trait values are sampled across a normal distribution defined by the 95% confidence interval at each node, over a specified number of matrices (in this case 100).

12.3 Measuring raw distance

An adapted Hamming distance metric was used to measure the raw accuracy at each node and state between the true and estimated discrete matrices. Both uncertain (where no state surpasses the threshold set by the majority rules) and incorrect estimates were penalised with a score of 1, thus only correct estimates were not penalised (score of 0). To achieve fair comparison between each fossil sampling level, only the nodes present across all fossil sampling levels were included in these analyses (leaving 50 nodes). This outputted a distance matrix of dimensions N by C, whereby N is the number of nodes and C is the number of characters. The mean was taken from each matrix to output a distance score between 0–1.

For continuous characters, estimation error was quantified by calculating the Euclidean distance between the estimated and true trait values at each node. Estimation values were Z-score standardised using the scaling parameters of the corresponding true matrix prior to analysis to ensure fair comparison between BM and OU models.

12.4 Measuring error in disparity

To evaluate how ancestral state estimation influences disparity estimates, both the true and estimated trait matrices were analysed using a standard disparity pipeline (**??**). For discrete characters, the raw trait matrices were converted into distance matrices using the MORD metric (**?**), the most widely used distance metric in disparity analyses. These distance matrices were ordinated using principal coordinates analysis (PCO) to generate the multidimensional traitspaces. Disparity was quantified as the sum of variances across all PCO axes for each combination of tree, transition rate, fossil sampling level, and majority-rule method. To measure the estimation accuracy in disparity, the estimated disparity value was subtracted by the true disparity value and then divided by the true disparity value, to give a relative difference in disparity (this is necessary since true disparity values can vary greatly between different parameter combinations).

For continuous characters, the trait matrices were ordinated using principal component analysis (PCA) and then also disparity measured using the sum of variances across all PC axes. These analyses were performed using the `dispRity` package (**?**).

13 Results and Discussion

13.1 Discrete characters

Raw ancestral state estimation error across all nodes varies with the majority-rule method, transition rate and fossil sampling level (**??**). Across all methods, transition rates strongly influenced estimation accuracy. Estimation accuracy was consistently highest when transition rates were slow, with estimation accuracy decreasing as rates increased. The strict and relative majority rule methods were less accurate in estimation compared to the sample-based approach for fast and medium transition rates, but recovered similar accuracy for slow rates. Notably, fossil sampling level had a relatively negligible effect on accuracy, which contrasts previous findings (**??**). Higher fossil sampling slightly improved estimates under strict and relative majority rules at fast and medium rates. In contrast, the sample-based approach remained stable across all fossil sampling levels, highlighting its robustness.

figures/strictdistance.pdf

Figure 7: Mean distance of ancestral state estimations to true value across all nodes, plotted against fossil sampling level, for each majority rule and transition rate (fast = 0.9, medium = 0.1, slow = 0.01). The shaded area represents the upper and lower quartile range across the 100 tree replicates.

When assessing accuracy at the root node solely, similar patterns are observed, with the no majority-rule approach outperforming majority-rule methods (**??**). At the root node, fossil sampling had a greater impact, as increases in sampling increase estimation accuracy for medium and slow transition rates, but had little effect on fast rates. The importance of fossil inclusion is highlighted here, particularly at deeper nodes, when uncertainty increases due to the diffusion of informative observed data. However, the no majority-rule method appears to be robust against low fossil sampling and high accuracy was maintained.

figures/rootdistance.pdf

Figure 8: Mean distance of ancestral state estimations to true value at root node, plotted against fossil sampling level, for each majority rule and transition rate (fast = 0.9, medium = 0.1, slow = 0.01). The shaded area represents the upper and lower quartile range across 100 tree replicates.

Interestingly, the pattern observed when assessing the estimation of disparity differs to that of raw estimates (**??**). The no-majority rule approach again outperformed the majority-rule approaches, was robust against low fossil sampling and the true disparity value was recovered with high accuracy across all three transition rates. In contrast, the majority-rule approaches were more substantially affected by low fossil sampling, particularly at medium rates, which returned considerable variation in the disparity values returned. Somewhat counter-intuitively, the disparity value returned from fast transition rates were generally accurate, despite having the highest error in raw estimates.

figures/sumvardistance.pdf

Figure 9: Relative distance of disparity value from estimated ancestral states to the true disparity value across all PCO axes, plotted against fossil sampling level, for each majority rule and transition rate (fast = 0.9, medium = 0.1, slow = 0.01). The shaded area represents the upper and lower quartile range across 100 tree replicates.

From these preliminary findings it can be inferred that, even when ancestral state estimation accuracy is poor at a 'raw state' level, accurate disparity signals are often still recovered. In addition, these findings highlight that accuracy should not be sacrificed for the sake of precision with ancestral state estimation. By incorporating uncertainty down the disparity pipeline using the sample approach, as opposed to deterministic methods such as the majority-rule, higher accuracy is recovered (**?**). For instance, when a true ancestral state has a posterior probability of 0.4, threshold-based approaches may discard it entirely, whereas the sample-based approach retains that state in 40% of reconstructions. This probabilistic representation enables more

accurate recovery of trait variance and disparity, especially in cases where no single state dominates the posterior.

The incorporation of uncertainty in macroevolutionary workflows has been shown numerous times previously to yield more robust inferences (**??**), yet ancestral state and disparity workflows often default to point estimates. These results highlight the utility of posterior sampling in ancestral state estimation and the robustness of disparity analyses to recovering true signals. Propagating uncertainty through posterior sampling offers a more rigorous alternative to commonly used approaches such as bootstrapping, and should be considered the standard workflow for researchers conducting disparity analyses on discrete traits. Unlike bootstrapping, posterior sampling captures the full distribution of ancestral state estimates, leading to more robust downstream inferences.

13.2 Continuous characters

Raw estimation accuracy for continuous characters was found to vary between models of trait evolution and method of handling uncertainty, but not across fossil sampling levels (**??**). Estimation accuracy of OU traits were more accurate than accuracy of BM trait estimation. This may seem intuitive, since estimation accuracy is expected to decrease as trait variance, σ, increases (**?**), although other studies have suggested that model misspecification (in this case OU when estimating traits under BM) can lead to erroneous estimates (**?**). In an OU model, trait variance is restricted due to the stabilising selection, denoted by the α parameter. Therefore, it can be hypothesised from these results that the greater the strength of α, the more accurate the estimations (something that will be tested in future analyses). Furthermore, the fact that estimation accuracy does not improve across higher fossil sampling levels is surprising, particularly given that accurate estimation of the primary optimum/root state is thought to be only possible with fossils (**?**).

figures/contdist.pdf

Figure 10: Euclidean distance between estimated ancestral continuous trait and true trait values, across BM/OU models and different fossil sampling levels. The shaded area represents the upper and lower quartile range across 100 tree replicates

Disparity estimation accuracy from continuous traits varied markedly between models and methods (**??**). Across all fossil sampling levels, estimates of BM traits derived from point estimates (left panel) were more accurate than those derived from a sample-approach. Estimates derived from sampling resulted in wide variation in estimates, which is likely due to the high uncertainty in the confidence intervals of estimated states. A contrasting pattern was observed for OU traits, with the sample-based approach recovering higher estimation accuracy than the point estimate approach, although both estimates had very low variance. Similarly to **??**, fossil sampling had little effect on estimation accuracy across both models.

figures/sumvarcontinuous.pdf

Figure 11: Relative disparity difference between traitspace of estimated ancestral continuous trait and traitspace of true trait values, across BM/OU models and different fossil sampling levels.

It remains difficult to directly compare the accuracy of ancestral state estimation between discrete and continuous traits, due to the nature of the outputted values. However, an intriguing contrast between the two data types is that for discrete characters, the sample-based approach consistently outperforms threshold-based methods, yielding higher accuracy in both raw state estimation and recovered disparity. In contrast, for continuous traits, the sample-based approach performs worse than point estimate methods. This divergence likely reflects the nature of uncertainty in each framework. For discrete traits, uncertainty is bounded by a finite number of states, making it feasible to probabilistically incorporate less certain outcomes without dramatically increasing error. For continuous traits, however, uncertainty is often expressed as wide confidence intervals over an infinite trait space, and sampling from these broad distributions may introduce excessive

noise into the reconstructed values. These findings suggest that while uncertainty propagation is beneficial for discrete characters, more caution is warranted when applying similar approaches to continuous traits. Future research should explore whether this limitation is driven by the models of trait evolution used, such as Brownian motion versus OU, or by the methods of estimating and sampling continuous trait distributions. Additionally, refining the way uncertainty is integrated, such the refining the parameters of the distribution that samples the confidence interval, could improve estimations.

14 Future directions

14.1 Extinction selectivity

These analyses will then be repeated, but incorporating mass extinction events into the trait and tree simulation process. A mass extinction event can be incorporated into the simulation framework using the `make.events` function, where a parameter is 'targeted' (in this case taxa) with a 'modification' (in this case extinction, and the proportion of lineages that goes extinct can be set) triggered by a 'condition' (can be set at an arbitrary time or when a certain number of taxa is reached). In addition, the modification can take a condition; for example, a condition might be that all lineages with positive trait values go extinct, effectively simulating a selective extinction.

Using this framework, I will simulate random mass extinction events at 70, 80 and 90%. I will also simulate selective mass extinctions at varying levels of selectivity.

14.2 Empirical tests

The same procedure will also be carried out on empirical data to benchmark these findings. I will conduct a literature search for available datasets with an exceptional fossil record; these will likely be datasets of Foraminifera, such as the one used in **?**. These datasets will be augmented to remove observed data across fossil tips in the same procedure as with simulated, and then estimate ancestral states to discern how accurate estimates are compared to the known trait state. The same procedure will be applied to analysis of disparity.