

Lecture 12

Philosophy 109

Caley Howland

November 1, 2019

- Last time, we ended up with proposal 3 to characterize inductive strength.

- Last time, we ended up with proposal 3 to characterize inductive strength.
 - ▶ **Proposal #3** “ $P \therefore C$ ” is strong just in case (1) the probability of $\sim C$ given that P is low, and (2) P is *positively relevant* to C — i.e., the probability of $\sim C$ given that P is lower than the probability of $\sim C$.

Proposal 3 and Relevance

- The third proposal adds a positive relevance requirement.

Proposal 3 and Relevance

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.

Proposal 3 and Relevance

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.
- The probability that c is not a spade, given that c is black is $1/2$. That is, the probability that $\sim C$, given that P is $1/2$.


Proposal 3 and Relevance

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.
- The probability that c is not a spade, given that c is black is $1/2$. That is, the probability that $\sim C$, given that P is $1/2$.
- This is not low (i.e., it is not less than $1/2$). But, it is *lower* than the probability that c is not a spade (i.e., the probability of $\sim C$), which is $3/4$.

- So, in this case, P is *positively relevant to* C
 - The probability of C given that $P = 1/2$, which is greater than the probability of $C = 1/4$).

- So, in this case, P is *positively relevant* to C
 - The probability of C given that $P = 1/2$, which is greater than the probability of $C = 1/4$).
- Thus, the argument “ $P \therefore C$ ” does not come out strong on proposal #3 (because it fails the low conditional probability requirement). But this argument does satisfy the positive relevance requirement.

- So, in this case, P is *positively relevant* to C
 - The probability of C given that $P = 1/2$, which is greater than the probability of $C = 1/4$).
- Thus, the argument “ $P \therefore C$ ” does not come out strong on proposal #3 (because it fails the low conditional probability requirement). But this argument does satisfy the positive relevance requirement.
- With proposal #3 in hand, we can now explain how our “scale” of argument strength works.
 - It depends on $Pr(C|P)$ and $Pr(C|P) > Pr(C)$.



Monotonicity

- A crucial difference between validity vs strength is that deductive validity is monotonic, but inductive strength is non-monotonic.
- If an argument is valid, then it cannot be rendered invalid merely by adding additional premises to it. Let's think about why this is true.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it.
Example, (\mathcal{A}):

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it.
Example, (\mathcal{A}):

- ▶ (P) It is January.
(C) If I walk out onto Lake Mendota, I will not get wet.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it. Example, (\mathcal{A}):
 - (P) It is January.
(C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So, \mathcal{A} is strong.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it.
Example, (\mathcal{A}):
 - (P) It is January.
(C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So, \mathcal{A} is strong.
- But consider \mathcal{A}' :
 - (P) It is January.
(Q) Lake Mendota is not frozen through.
 \therefore (C) If I walk out onto Lake Mendota, I will not get wet.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it. Example, (\mathcal{A}):
 - (P) It is January.
(C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So, \mathcal{A} is strong.
- But consider \mathcal{A}' :
 - (P) It is January.
(Q) Lake Mendota is not frozen through.
 \therefore (C) If I walk out onto Lake Mendota, I will not get wet.
- \mathcal{A}' is weak. So adding a new premise—Premise (Q)—made it weak.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
 - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
 - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
 - ▶ (R) John has yellow stains on his fingers and smells like tobacco.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
 - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
 - ▶ (R) John has yellow stains on his fingers and smells like tobacco.
 - ▶ (S) John rolls cigars for a living.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
 - ▶ (P) John has lung cancer.
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
 - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
 - ▶ (R) John has yellow stains on his fingers and smells like tobacco.
 - ▶ (S) John rolls cigars for a living.
- And so on. . . This non-monotonicity of inductive strength is a key feature. Diagramming can help explain why this can happen.

Numerical Probability

- So far, we have usually been relying on *intuitions* about the “inductive probabilities” in various examples.
 - ▶ Sometimes, we’ve discussed games of chance for which there are clear, *numerical* (inductive) probabilities.

Numerical Probability

- So far, we have usually been relying on *intuitions* about the “inductive probabilities” in various examples.
 - Sometimes, we’ve discussed games of chance for which there are clear, *numerical* (inductive) probabilities.
- But, we haven’t seen a precise, general way to think about (or calculate) numerical probabilities of LSL sentences. This is our current topic.

- It turns out that numerical probabilities can be understood (and calculated) using a simple numerical generalization LSL truth-tables.

- It turns out that numerical probabilities can be understood (and calculated) using a simple numerical generalization LSL truth-tables.
- I will call these Probabilistic Truth Tables (or sometimes “Stochastic Truth Tables”).

- It turns out that numerical probabilities can be understood (and calculated) using a simple numerical generalization LSL truth-tables.
- I will call these Probabilistic Truth Tables (or sometimes “Stochastic Truth Tables”).
- A probabilistic truth-table is just a truth-table with an additional column. That column gives the inductive **probabilities of each of the possible worlds** (i.e., the inductive probabilities of each of the rows).

- It turns out that numerical probabilities can be understood (and calculated) using a simple numerical generalization LSL truth-tables.
- I will call these Probabilistic Truth Tables (or sometimes “Stochastic Truth Tables”).
- A probabilistic truth-table is just a truth-table with an additional column. That column gives the inductive **probabilities of each of the possible worlds** (i.e., the inductive probabilities of each of the rows).
- It helps to start with a simple example (involving a game of chance).

- Let's start with our example of drawing a single card (c) at random from a standard deck of playing cards. There are various properties that our card might have. Here are two examples (as atomic LSL sentences).
 - ▶ $H = c$ is a heart. $R = c$ is a red card.

Probabilistic Truth Table

- A **probabilistic truth-table** (involving H and R) will look like this.

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

Probabilistic Truth Table

- A **probabilistic truth-table** (involving H and R) will look like this.

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

- The column on the left names the interpretations, the worlds, but they are just determined by the guide columns like usual.

Probabilistic Truth Table

- A **probabilistic truth-table** (involving H and R) will look like this.

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

- The column on the left names the interpretations, the worlds, but they are just determined by the guide columns like usual.
- The new right hand column assigns probabilities to these worlds.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - Take sentences as input and give numbers as output.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - Take sentences as input and give numbers as output.
 - Such numbers are called “probabilities”. They must meet certain constraints.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - Take sentences as input and give numbers as output.
 - Such numbers are called “probabilities”. They must meet certain constraints.
 - They are always Real numbers between 0 and 1.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - ▶ Take sentences as input and give numbers as output.
 - ▶ Such numbers are called “probabilities”. They must meet certain constraints.
 - ▶ They are always Real numbers between 0 and 1.
- Real numbers include Naturals, Rationals, and Irrationals.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - Take sentences as input and give numbers as output.
 - Such numbers are called “probabilities”. They must meet certain constraints.
 - They are always Real numbers between 0 and 1.
- Real numbers include Naturals, Rationals, and Irrationals.
- Real numbers are uncountably infinite.

- Here, we are “assigning” probabilities to sentences in our language.
- Probability assignments are expressed by probability functions: $Pr(\cdot) = n$
 - ▶ Take sentences as input and give numbers as output.
 - ▶ Such numbers are called “probabilities”. They must meet certain constraints.
 - ▶ They are always Real numbers between 0 and 1.
- Real numbers include Naturals, Rationals, and Irrationals.
- Real numbers are uncountably infinite.
- We will generally round things so as not to deal with irrationals.

- What does “probability” mean?

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms
- When mathematicians say “probability”, this is what they mean.

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms
- When mathematicians say “probability”, this is what they mean.
- Kolmogorov Axioms (expressed with a function over LSL)

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms
- When mathematicians say “probability”, this is what they mean.
- Kolmogorov Axioms (expressed with a function over LSL)
 - ▶ Non-negativity: for any sentence p , $Pr(p) \geq 0$.

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms
- When mathematicians say “probability”, this is what they mean.
- Kolmogorov Axioms (expressed with a function over LSL)
 - ▶ Non-negativity: for any sentence p , $Pr(p) \geq 0$.
 - ▶ Normativity: for any logical truth of LSL \top , $Pr(\top) = 1$

- What does “probability” mean?
- One answer: a probability function is a function with certain mathematical properties: it satisfies the Kolmogorov axioms
- When mathematicians say “probability”, this is what they mean.
- Kolmogorov Axioms (expressed with a function over LSL)
 - ▶ Non-negativity: for any sentence p , $Pr(p) \geq 0$.
 - ▶ Normativity: for any logical truth of LSL \top , $Pr(\top) = 1$
 - ▶ Finite Additivity: for any mutually exclusive propositions p or q , $Pr(p \vee q) = Pr(p) + Pr(q)$.

- What does “probability” mean?

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.
 - Outcomes of chance structures: dice.

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.
 - Outcomes of chance structures: dice.
 - Frequencies

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.
 - Outcomes of chance structures: dice.
 - Frequencies
 - Objective Chances (e.g., from physics)

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.
 - Outcomes of chance structures: dice.
 - Frequencies
 - Objective Chances (e.g., from physics)
 - Degrees of belief: credences

- What does “probability” mean?
- The various things we can describe with the mathematical functions meeting those axioms.
- There is a non-settled debate about what these things are.
 - Outcomes of chance structures: dice.
 - Frequencies
 - Objective Chances (e.g., from physics)
 - Degrees of belief: credences
- This last idea, that probability functions describe beliefs, is called “personalism”, and is part of what’s called Bayesianism

- The details of these debates won't matter much to us.

- The details of these debates won't matter much to us.
- To start with, I will talk about “inductive probabilities”, which are just probabilities used to measure argument strength

- The details of these debates won't matter much to us.
- To start with, I will talk about “inductive probabilities”, which are just probabilities used to measure argument strength
 - ▶ In other words, we will stay pretty neutral.

- The details of these debates won't matter much to us.
- To start with, I will talk about “inductive probabilities”, which are just probabilities used to measure argument strength
 - ▶ In other words, we will stay pretty neutral.
- However, in the background think about probabilities as degrees of belief strength; we will come back to that.

- The details of these debates won't matter much to us.
- To start with, I will talk about “inductive probabilities”, which are just probabilities used to measure argument strength
 - ▶ In other words, we will stay pretty neutral.
- However, in the background think about probabilities as degrees of belief strength; we will come back to that.
- Plus, that is how to read Titelbaum: Titelbaum uses $cr(\cdot)$ to make it explicit that is the kind of probability he is interested in.

- An inductive probability assignment is just as assignment of world probabilities. (These are technically a kind of distribution.)

- An inductive probability assignment is just as assignment of world probabilities. (These are technically a kind of distribution.)
 - Numbers between 0 and 1 that sum to 1.

- An inductive probability assignment is just as assignment of world probabilities. (These are technically a kind of distribution.)
 - ▶ Numbers between 0 and 1 that sum to 1.
- Once we have world probabilities assigned (involving some atomic sentences), we can then calculate the probability of *any LSL statement* (involving those atomic sentences). In fact, the calculation is easy.

- An inductive probability assignment is just an assignment of world probabilities. (These are technically a kind of distribution.)
 - ▶ **Numbers between 0 and 1 that sum to 1.**
- Once we have world probabilities assigned (involving some atomic sentences), we can then calculate the probability of *any LSL statement* (involving those atomic sentences). In fact, the calculation is easy.
- The probability of an LSL sentence s is just **the sum of the world probabilities of the possible worlds in which s is true.**

Probabilistic Truth Table

- Remember our **probabilistic truth-table** (involving H and R).

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

Probabilistic Truth Table

- Remember our **probabilistic truth-table** (involving H and R).

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

- Each possible world w_i has a definite, numerical probability.

Probabilistic Truth Table

- Remember our **probabilistic truth-table** (involving H and R).

World (w_i)	H	R	$Pr(w_i)$	
w_1	T	T	$1/4$	$[= Pr(H \& R)]$
w_2	T	F	0	$[= Pr(H \& \sim R)]$
w_3	F	T	$1/4$	$[= Pr(\sim H \& R)]$
w_4	F	F	$1/2$	$[= Pr(\sim H \& \sim R)]$

- Each possible world w_i has a definite, numerical probability.
- Once these world probabilities are given, all other probabilities can be calculated.

- Example: What is $Pr(H \rightarrow R)$?

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?

World (w_i)	H	R	$Pr(w_i)$	$H \rightarrow R$
w_1	T	T	$1/4$	T
w_2	T	F	0	F
w_3	F	T	$1/4$	T
w_4	F	F	$1/2$	T

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?
- The only world in which $H \rightarrow R$ is false is w_2 . So, the probability of $H \rightarrow R$ is just the sum of the world probabilities for w_1 , w_3 , and w_4 .

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?
- The only world in which $H \rightarrow R$ is false is w_2 . So, the probability of $H \rightarrow R$ is just the sum of the world probabilities for w_1 , w_3 , and w_4 .
- Therefore, $Pr(H \rightarrow R) = 1/4 + 1/4 + 1/2 = 1$.

- Example: What is $Pr(H \rightarrow R)$?
- Well, which possible worlds are worlds in which $H \rightarrow R$ is true?
- The only world in which $H \rightarrow R$ is false is w_2 . So, the probability of $H \rightarrow R$ is just the sum of the world probabilities for w_1 , w_3 , and w_4 .
- Therefore, $Pr(H \rightarrow R) = 1/4 + 1/4 + 1/2 = 1$.
 - This makes sense, since all hearts are red cards. Thus, we can be certain that $H \rightarrow R$ is true.

- What is $Pr(R \rightarrow H)$?

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?

World (w_i)	H	R	$Pr(w_i)$	$R \rightarrow H$
w_1	T	T	$1/4$	T
w_2	T	F	0	T
w_3	F	T	$1/4$	F
w_4	F	F	$1/2$	T

- What is $Pr(R \rightarrow H)$?

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?
- The only world in which $R \rightarrow H$ is false is w_3 .

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?
- The only world in which $R \rightarrow H$ is false is w_3 .
- So, $Pr(R \rightarrow H) = 1/4 + 0 + 1/2 = 3/4$.

- What is $Pr(R \rightarrow H)$?
- Which possible worlds are worlds in which $R \rightarrow H$ is true?
- The only world in which $R \rightarrow H$ is false is w_3 .
- So, $Pr(R \rightarrow H) = 1/4 + 0 + 1/2 = 3/4$.
- According to Proposal 1 for inductive strength, the argument $R \therefore H$ is going to come out strong, since $Pr(R \rightarrow H) = 3/4$, which is high.