

# Lecture 13

Philosophy 109

Caley Howland

# Administrative Stuff

- **Required** for next time:
  - ▶ Instead of new reading, do the “Chances Worksheet” on sakai.
  - ▶ Exercises: Forallx Leeds, Chapter 10, Parts A and B.
- Schedule is updated on Sakai
- Answer key to probabilistic truth table exercises will be posted shortly.
- HW 4 will be posted soon.
- I have your blue books for the exam.

# Conditional Probability

- In order to evaluate inductive strength, we need to calculate the probability of  $p$  given  $q$ , which is denoted  $Pr(p|q)$ .

# Conditional Probability

- In order to evaluate inductive strength, we need to calculate the probability of  $p$  given  $q$ , which is denoted  $Pr(p|q)$ .

# Conditional Probability

- In order to evaluate inductive strength, we need to calculate the probability of  $p$  given  $q$ , which is denoted  $Pr(p|q)$ .

## Definition of Conditional Probability

$$Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$$

- This definition makes sense, because we want to look at how probable  $p$  is within the  $q$ -worlds. And, that's just the ratio of  $Pr(p \wedge q)$  to  $Pr(q)$ .

# Inductive Strength: Precise

## Precise Inductive Strength

An argument  $P \therefore C$  is inductively strong iff:

- (1)  $C$  is probable given  $P$ , i.e.,  $Pr(C|P) > 1/2$ , and
- (2)  $P$  is **positively relevant** to  $C$ , i.e.,  $Pr(C|P) > Pr(C)$

# Rules for Calculating $Pr(p)$ and $Pr(p|q)$

- Once you're given a probabilistic truth-table, there are only two rules for calculating numerical probabilities.
- A probabilistic truth-table specifies world-probabilities for all possible worlds involving some fixed set of TFL atomic sentences.

## Rules for Calculation

# Rules for Calculating $Pr(p)$ and $Pr(p|q)$

- Once you're given a probabilistic truth-table, there are only two rules for calculating numerical probabilities.
- A probabilistic truth-table specifies world-probabilities for all possible worlds involving some fixed set of TFL atomic sentences.

## Rules for Calculation

- 1  $Pr(p)$  is the *sum* of the world-probabilities of the worlds in which  $p$  is true. That is,  $Pr(p)$  is the sum of the  $p$ -world probabilities.



# Rules for Calculating $Pr(p)$ and $Pr(p|q)$

- Once you're given a probabilistic truth-table, there are only two rules for calculating numerical probabilities.
- A probabilistic truth-table specifies world-probabilities for all possible worlds involving some fixed set of TFL atomic sentences.

## Rules for Calculation

- ①  $Pr(p)$  is the *sum* of the world-probabilities of the worlds in which  $p$  is true. That is,  $Pr(p)$  is the sum of the  $p$ -world probabilities.
- ②  $Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$

# Rules for Calculating $Pr(p)$ and $Pr(p|q)$

- Once you're given a probabilistic truth-table, there are only two rules for calculating numerical probabilities.
- A probabilistic truth-table specifies world-probabilities for all possible worlds involving some fixed set of TFL atomic sentences.

## Rules for Calculation

- ①  $Pr(p)$  is the *sum* of the world-probabilities of the worlds in which  $p$  is true. That is,  $Pr(p)$  is the sum of the  $p$ -world probabilities.
- ②  $Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$

Note: If  $Pr(q) = 0$ , then  $Pr(p|q)$  is undefined, since you can't divide by 0. So, we always assume  $Pr(q) > 0$  when we use  $Pr(p|q)$ .

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- State Descriptions:

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- **State Descriptions:**

- ▶ A special name for each world, i.e. each row

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- State Descriptions:

- ▶ A special name for each world, i.e. each row
- ▶ Tell which atomic sentences at that world are true.

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- State Descriptions:

- ▶ A special name for each world, i.e. each row
- ▶ Tell which atomic sentences at that world are true.
- ▶ Conjunctions, with negated sentences for those atomic sentences which are false.

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- **State Descriptions:**

- ▶ A special name for each world, i.e. each row
- ▶ Tell which atomic sentences at that world are true.
- ▶ Conjunctions, with negated sentences for those atomic sentences which are false.
- ▶ Conjunctions which are only true in one world.

# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- **State Descriptions:**

- ▶ A special name for each world, i.e. each row
- ▶ Tell which atomic sentences at that world are true.
- ▶ Conjunctions, with negated sentences for those atomic sentences which are false.
- ▶ Conjunctions which are only true in one world.

- **Disjunctive Normal form:** any sentence can be expressed in this form by a disjunction of state descriptions:



# State Descriptions

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- **State Descriptions:**

- ▶ A special name for each world, i.e. each row
- ▶ Tell which atomic sentences at that world are true.
- ▶ Conjunctions, with negated sentences for those atomic sentences which are false.
- ▶ Conjunctions which are only true in one world.

- **Disjunctive Normal form:** any sentence can be expressed in this form by a disjunction of state descriptions:

- ▶  $p \Leftrightarrow (p \wedge q) \vee (p \wedge \sim q)$

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

①  $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (General Disjunction)

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- 1  $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (General Disjunction)
- 2  $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$  (Law of Total Probability)

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- ①  $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (General Disjunction)
- ②  $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$  (Law of Total Probability)
- ③  $Pr(p|q) = \frac{Pr(q|p) \cdot Pr(p)}{Pr(q)}$  (Bayes' Theorem)

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- ①  $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (General Disjunction)
- ②  $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$  (Law of Total Probability)
- ③  $Pr(p|q) = \frac{Pr(q|p) \cdot Pr(p)}{Pr(q)}$  (Bayes' Theorem)
- ④  $Pr(p|p \vee q) > Pr(p)$  if  $Pr(p) > 0$  and  $Pr(p \vee B) < 1$ .

# Some Derived Rules

- We can use generic probabilistic truth-tables to derive additional rules. Here are some examples.

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- ①  $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (General Disjunction)
- ②  $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$  (Law of Total Probability)
- ③  $Pr(p|q) = \frac{Pr(q|p) \cdot Pr(p)}{Pr(q)}$  (Bayes' Theorem)
- ④  $Pr(p|p \vee q) > Pr(p)$  if  $Pr(p) > 0$  and  $Pr(p \vee B) < 1$ .
- ⑤  $Pr(p|p \vee \neg p) = Pr(p)$

# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)



# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)
- We know:  $Pr(p \vee q) = a + b + c$ , and  $Pr(p \wedge q) = a$ .

# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)
- We know:  $Pr(p \vee q) = a + b + c$ , and  $Pr(p \wedge q) = a$ .
- And:  $Pr(p) = a + b$ , and  $Pr(q) = a + c$

# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)
- We know:  $Pr(p \vee q) = a + b + c$ , and  $Pr(p \wedge q) = a$ .
- And:  $Pr(p) = a + b$ , and  $Pr(q) = a + c$
- $a + b + c = a + b + c + \textcolor{red}{a - a}$

# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)
- We know:  $Pr(p \vee q) = a + b + c$ , and  $Pr(p \wedge q) = a$ .
- And:  $Pr(p) = a + b$ , and  $Pr(q) = a + c$
- $a + b + c = a + b + c + \textcolor{red}{a - a}$
- $= Pr(p) + Pr(q) - a$

# General Disjunction Rule

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \vee q) = Pr(p) + Pr(q) - Pr(p \wedge q)$  (Disjunction Rule)
- We know:  $Pr(p \vee q) = a + b + c$ , and  $Pr(p \wedge q) = a$ .
- And:  $Pr(p) = a + b$ , and  $Pr(q) = a + c$
- $a + b + c = a + b + c + \textcolor{red}{a - a}$
- $= Pr(p) + pr(q) - a$
- $Pr(p) + Pr(q) - Pr(p \wedge q)$ .

# Total Probability

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$

# Total Probability

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p) = Pr(p \wedge q) + Pr(p \wedge \neg q)$
- $a + b = a + b$

# General Conjunction

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \wedge q) = Pr(p|q) \times Pr(q)$



# General Conjunction

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \wedge q) = Pr(p|q) \times Pr(q)$
- $Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$

# General Conjunction

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \wedge q) = Pr(p|q) \times Pr(q)$
- $Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$
- $Pr(p|q) \times Pr(q) = Pr(p \wedge q)$

# General Conjunction

World( $w_i$ )	$p$	$q$	$Pr(w_i)$	State Desc.
$w_1$	T	T	$a$	$p \wedge q$
$w_2$	T	F	$b$	$p \wedge \neg q$
$w_3$	F	T	$c$	$\neg p \wedge q$
$w_4$	F	F	$d$	$\neg p \wedge \neg q$

- $Pr(p \wedge q) = Pr(p|q) \times Pr(q)$
- $Pr(p|q) = \frac{Pr(p \wedge q)}{Pr(q)}$
- $Pr(p|q) \times Pr(q) = Pr(p \wedge q)$
- Alternative:  $Pr(p|q) = \frac{a}{a+c} \times a + c$

# An Example from Earlier

- Recall: we compared these two arguments:
  - (i)  $A \vee B \therefore A$ , and
  - (ii)  $A \vee \neg A \therefore A$

# An Example from Earlier

- Recall: we compared these two arguments:
  - (i)  $A \vee B \therefore A$ , and
  - (ii)  $A \vee \neg A \therefore A$
- (i) is better than (ii) even though neither is valid. Let's see if probability agrees.

# An Example from Earlier

- Recall: we compared these two arguments:
  - (i)  $A \vee B \therefore A$ , and
  - (ii)  $A \vee \neg A \therefore A$
- (i) is better than (ii) even though neither is valid. Let's see if probability agrees.
- By (4) from above, we know that if  $Pr(A) > 0$  and  $Pr(A \vee B) < 1$ , we have  $Pr(A|A \vee B) > Pr(A)$ . So, (i)'s premise will usually be relevant to its conclusion.

# An Example from Earlier

- Recall: we compared these two arguments:
  - (i)  $A \vee B \therefore A$ , and
  - (ii)  $A \vee \neg A \therefore A$
- (i) is better than (ii) even though neither is valid. Let's see if probability agrees.
- By (4) from above, we know that if  $Pr(A) > 0$  and  $Pr(A \vee B) < 1$ , we have  $Pr(A|A \vee B) > Pr(A)$ . So, (i)'s premise will usually be relevant to its conclusion.
- But, by (5), we know that  $Pr(A|A \vee \neg A) = Pr(A)$ . So, (ii)'s premise is always irrelevant to its conclusion.

# An Example from Earlier

- Recall: we compared these two arguments:
  - (i)  $A \vee B \therefore A$ , and
  - (ii)  $A \vee \neg A \therefore A$
- (i) is better than (ii) even though neither is valid. Let's see if probability agrees.
- By (4) from above, we know that if  $Pr(A) > 0$  and  $Pr(A \vee B) < 1$ , we have  $Pr(A|A \vee B) > Pr(A)$ . So, (i)'s premise will usually be relevant to its conclusion.
- But, by (5), we know that  $Pr(A|A \vee \neg A) = Pr(A)$ . So, (ii)'s premise is always irrelevant to its conclusion.
- Therefore, by Proposal #3, argument (i) is stronger than argument (ii).



# Monotonicity

- A crucial difference between validity vs strength is that deductive validity is monotonic, but inductive strength is non-monotonic.
- If an argument is valid, then it cannot be rendered invalid merely by adding additional premises to it. Let's think about why this is true.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it. Example, ( $\mathcal{A}$ ):

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it.

Example, ( $\mathcal{A}$ ):

- ▶ (P) It is January.  
(C) If I walk out onto Lake Mendota, I will not get wet.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it. Example, ( $\mathcal{A}$ ):
  - (P) It is January.  
(C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So,  $\mathcal{A}$  is strong.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it.  
Example, ( $\mathcal{A}$ ):
  - ▶ (P) It is January.  
(C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So,  $\mathcal{A}$  is strong.
- But consider  $\mathcal{A}'$ :
  - ▶ (P) It is January.  
(Q) Lake Mendota is not frozen through.  
 $\therefore$  (C) If I walk out onto Lake Mendota, I will not get wet.

- On the other hand, an inductively strong argument can be rendered inductively weak merely by adding premises to it. Example, ( $\mathcal{A}$ ):
  - (P) It is January.
  - (C) If I walk out onto Lake Mendota, I will not get wet.
- Usually, Lake Mendota is frozen in January. So,  $\mathcal{A}$  is strong.
- But consider  $\mathcal{A}'$ :
  - (P) It is January.
  - (Q) Lake Mendota is not frozen through.
  - $\therefore$  (C) If I walk out onto Lake Mendota, I will not get wet.
- $\mathcal{A}'$  is weak. So adding a new premise—Premise (Q)—made it weak.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.



- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
  - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
  - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
  - ▶ (R) John has yellow stains on his fingers and smells like tobacco.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
  - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
  - ▶ (R) John has yellow stains on his fingers and smells like tobacco.
  - ▶ (S) John rolls cigars for a living.

- In fact, inductive arguments can *oscillate* in strength as new premises are added. Here is an example that illustrates the phenomenon.
  - ▶ (P) John has lung cancer.  
∴ (C) John is (or was) a smoker.
- By learning more things about John (i.e., by adding premises about John), the strength of this argument can oscillate. Consider:
  - ▶ (Q) John has a certain gene that would make him susceptible to lung cancer even if he were not a smoker.
  - ▶ (R) John has yellow stains on his fingers and smells like tobacco.
  - ▶ (S) John rolls cigars for a living.
- And so on. . . This non-monotonicity of inductive strength is a key feature. Diagramming can help explain why this can happen.