### Lecture 11

Philosophy 109

Caley Howland

March 23, 2020

1/29

 Caley Howland
 Lecture 11
 03/23/2020

#### Administrative Stuff

- Reading for Next time:
  - Skyrms Chapter 2 (Optional, but highly recommended)
  - Hacking Chapter 2
- We are starting with the new unit on inductive reasoning/probability.

### Inductive Logic

Intuitively, not all good arguments are deductively valid.

### **Boring Class**

The professor put me to sleep the first lecture.

The professor put me to sleep the second lecture.

:

The professor put me to sleep the seventeenth lecture.

... The professor will put me to sleep the eighteenth lecture.

• Some invalid arguments are stronger than others.

- Some invalid arguments are stronger than others.
  - Consider:
    - ★ The professor once put me to sleep. Therefore, the professor will put me to sleep next time.

- Some invalid arguments are stronger than others.
  - Consider:
    - ★ The professor once put me to sleep.
      Therefore, the professor will put me to sleep next time.
  - ► That argument is weaker than the argument on the last slide.

- Some invalid arguments are stronger than others.
  - Consider:
    - ★ The professor once put me to sleep. Therefore, the professor will put me to sleep next time.
  - ► That argument is weaker than the argument on the last slide.
- Inductive logic aims (in part) to study the strength to which an argument's premises support its conclusion.

- Some invalid arguments are stronger than others.
  - ► Consider:
    - ★ The professor once put me to sleep.
       Therefore, the professor will put me to sleep next time.
  - ► That argument is weaker than the argument on the last slide.
- Inductive logic aims (in part) to study the strength to which an argument's premises support its conclusion.
- It's inductive argument that's behind nearly all of science!
- It's also the kind of inference that you make most in your everyday life.

- Recall: If the premises of a valid argument are all true, the the conclusion is guaranteed to be true.
- Inductively strong arguments don't need this feature. But:
  - If the premises of an inductively strong argument are all true, then they make it probable that the conclusion is also true.

 Cates Howland
 Lecture 11
 03/23/2020
 5/29

#### The Killing

Smith has confessed to killing Jones. Dr. Zed signed a statement to the effect that he saw Smith shoot Jones. A large number of witnesses heard Jones gasp with his dying breath, "Smith did it." Therefore, Smith killed Jones.

6/29

Caley Howland Lecture 11 03/23/2020

### The Killing

Smith has confessed to killing Jones. Dr. Zed signed a statement to the effect that he saw Smith shoot Jones. A large number of witnesses heard Jones gasp with his dying breath, "Smith did it." Therefore, Smith killed Jones.

 This argument is not deductively valid. However, it's still a strong argument. Assuming its premises are true, its conclusion is probable.

 It's fairly easy to see that the argument isn't logically valid.
 However, for vividness we construct the following alternative:

- It's fairly easy to see that the argument isn't logically valid.
   However, for vividness we construct the following alternative:
  - ► Suppose that Smith wants to be a famous criminal, and that he (falsely) confessed to every murder he ever heard of, but that this fact was generally unknown because he just moved to the area. This peculiarity was, however, known to Dr. Zed who was Jones's psychiatrist. For malevolent reasons, Dr. Zed decided to kill Jones and frame Smith. He convinced Jones under hypnosis that Smith was a homicidal maniac bent on killing Jones. Then Dr. Zed shot Jones from behind a potted plant and fled.

- It's fairly easy to see that the argument isn't logically valid.
   However, for vividness we construct the following alternative:
  - ► Suppose that Smith wants to be a famous criminal, and that he (falsely) confessed to every murder he ever heard of, but that this fact was generally unknown because he just moved to the area. This peculiarity was, however, known to Dr. Zed who was Jones's psychiatrist. For malevolent reasons, Dr. Zed decided to kill Jones and frame Smith. He convinced Jones under hypnosis that Smith was a homicidal maniac bent on killing Jones. Then Dr. Zed shot Jones from behind a potted plant and fled.
- Obviously, this scenario is improbable, but it's not impossible.
  - So, it's possible that the premises of the original argument are all true but the conclusion false.

 Inductive argument strength comes in degrees of inductive probability.

8/29

Caley Howland Lecture 11 03/23/2020

- Inductive argument strength comes in degrees of inductive probability.
  - ► The type of probability that grades the inductive strength of arguments—we shall call it *inductive probability*—does not depend on the probability of the premises alone or on the probability of the conclusion alone, but on the *evidential relation between* the premises and the conclusion.

 When we judge the inductive strength of an argument, we must judge how strongly the premises evidentially support the conclusion.

- When we judge the inductive strength of an argument, we must judge how strongly the premises evidentially support the conclusion.
- This is much different than what we did with deductive validity.
  - Remember, we encountered some "strange/fringe cases" of validity that did not seem to involve any sort of evidential support at all.
  - Any argument with contradictory premises is valid, and any argument with a tautological conclusion is valid. But, these do not involve any relations of evidential support between premises and conclusion.

- When we judge the inductive strength of an argument, we must judge how strongly the premises evidentially support the conclusion.
- This is much different than what we did with deductive validity.
  - Remember, we encountered some "strange/fringe cases" of validity that did not seem to involve any sort of evidential support at all.
  - Any argument with contradictory premises is valid, and any argument with a tautological conclusion is valid. But, these do not involve any relations of evidential support between premises and conclusion.
- In the case of inductive arguments, there should be some degree of relevance between the premises and conclusion.

 For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.

- For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.
  - Deductive logic

10/29

Caley Howland Lecture 11 03/23/2020

- For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.
  - Deductive logic
    - Validity: the truth of the premises guarantees truth of conclusion.

- For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.
  - Deductive logic
    - Validity: the truth of the premises guarantees truth of conclusion.
  - ► Inductive logic

- For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.
  - Deductive logic
    - Validity: the truth of the premises guarantees truth of conclusion.
  - ► Inductive logic
    - ★ Strength: the truth of the premises makes the truth of conclusion probable.

- For both deductive and inductive logic, we are interested in the connection between the truth of premises and the truth of the conclusion.
  - ► Deductive logic
    - Validity: the truth of the premises guarantees truth of conclusion.
  - ► Inductive logic
    - ★ Strength: the truth of the premises makes the truth of conclusion probable.
- So, we really only care about when the premises are true.

 Because of this, we can always re-write arguments as having a single premise.

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

 Caley Howland
 Lecture 11
 03/23/2020
 11/29

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

P1 
$$A \rightarrow B$$

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

P1  $A \rightarrow B$ 

P2 A

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

```
P1 A \rightarrow B
```

P2 A

C B

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

P1 
$$A \rightarrow B$$

P2 A

CB

• is equivalent to this one:

 Caley Howland
 Lecture 11
 03/23/2020
 11/29

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

P1 
$$A \rightarrow B$$

P2 A

CB

• is equivalent to this one:

P1 
$$(A \rightarrow B) \land A$$

11/29

Caley Howland Lecture 11 03/23/2020

- Because of this, we can always re-write arguments as having a single premise.
- An argument with multiple premises is always equivalent to one with a single premise made up of a conjunction.
- So this argument:

P1 
$$A \rightarrow B$$
  
P2  $A$ 

CB

is equivalent to this one:

P1 
$$(A \rightarrow B) \land A$$

11/29

Some terminology:

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,

Caley Howland Lecture 11 03/23/2020 12 / 29

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,
     P1 A → B

Caley Howland Lecture 11 03/23/2020 12 / 29

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,

P1  $A \rightarrow B$ 

P2 A

Caley Howland Lecture 11 03/23/2020 12 / 29

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,

```
P1 A \rightarrow B
```

P2 A

CB

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,

P1 
$$A \rightarrow B$$

▶ Becomes 
$$(A \rightarrow B) \land A :: B$$

12/29

Caley Howland Lecture 11 03/23/2020

- Some terminology:
- And let us start writing expressing arguments on one line P :: C.
  - So our previous argument,

P1 
$$A \rightarrow B$$
  
P2  $A$   
C  $B$ 

- ▶ Becomes  $(A \rightarrow B) \land A : B$
- Let us also treat P as a variable, which stands for the conjunction of a set of premises.

12/29

 Caley Howland
 Lecture 11
 03/23/2020

- Some terminology:
- And let us start writing expressing arguments on one line P ∴ C.
  - So our previous argument,

```
P1 A \rightarrow B
P2 A
C B
```

- ▶ Becomes  $(A \rightarrow B) \land A : B$
- Let us also treat P as a variable, which stands for the conjunction of a set of premises.
- So we can express a generic argument as: P ∴ C.

(ロ) (固) (重) (重) (重) の(O)

- "The argument *P* ∴ *C* is valid" is equivalent to:
  - ▶ The claim the  $P \land \neg C$  is impossible (i.e., logically false).

Caley Howland Lecture 11 03/23/2020 13 / 29

- "The argument P : C is valid" is equivalent to:
  - ▶ The claim the  $P \land \neg C$  is impossible (i.e., logically false).
- A natural way to try to generalize this to inductive strength would be.
  - ▶ **Proposal 1**: An argument P ∴ C is inductively strong just in case the claim  $P \land \neg C$  is improbable.

Caley Howland Lecture 11 03/23/2020 13/29

- "The argument P : C is valid" is equivalent to:
  - ▶ The claim the  $P \land \neg C$  is impossible (i.e., logically false).
- A natural way to try to generalize this to inductive strength would be.
  - ▶ **Proposal 1**: An argument P : C is inductively strong just in case the claim  $P \land \neg C$  is improbable.
- **Proposal 1 won't work!** Let P="There's a 2000-year-old man in Cleveland" and C="There's a 2000-year-old man in Cleveland who has 3 heads."

Caley Howland Lecture 11 03/23/2020 13/29

- "The argument P : C is valid" is equivalent to:
  - ▶ The claim the  $P \land \neg C$  is impossible (i.e., logically false).
- A natural way to try to generalize this to inductive strength would be.
  - ▶ **Proposal 1**: An argument P : C is inductively strong just in case the claim  $P \land \neg C$  is improbable.
- **Proposal 1 won't work!** Let P="There's a 2000-year-old man in Cleveland" and C="There's a 2000-year-old man in Cleveland who has 3 heads."
  - ▶ Is  $P \land \neg C$  improbable? **Yes!**
  - Is P ∴ C actually a strong argument? No!

Caley Howland Lecture 11 03/23/2020 13/29

• Intuitively, the argument P : C is *not* strong. That's because P does not evidentially support C.

Caley Howland Lecture 11 03/23/2020 14/29

- Intuitively, the argument P : C is *not* strong. That's because P does not evidentially support C.
  - ▶ If there were a 2000 year old man in Cleveland, then it would be highly likely that he would have only one head.

 Caley Howland
 Lecture 11
 03/23/2020
 14/29

- Intuitively, the argument P : C is *not* strong. That's because P does not evidentially support C.
  - ▶ If there were a 2000 year old man in Cleveland, then it would be highly likely that he would have only one head.
- So,  $P \land \neg C$  is improbable but only because P is improbable.

Caley Howland Lecture 11 03/23/2020 14/29

- There are also examples in which  $P \land \neg C$  is improbable merely because C is probable (independently of C's relationship to P).
  - ► Let *P*="A fair coin will land heads", and *C*="No man will live to be 2000".

 Cates Howland
 Lecture 11
 03/23/2020
 15/29

- There are also examples in which  $P \land \neg C$  is improbable merely because C is probable (independently of C's relationship to P).
  - ► Let *P*="A fair coin will land heads", and *C*="No man will live to be 2000".
- Moral: Looking at the (im)probability of the claim  $P \land \neg C$  does not (in general) tell us the degree to which an argument is inductively strong.


Caley Howland Lecture 11 03/23/2020 15 / 29

• How should we understand "It is improbable that C is false while P is true"? The "while" can't mean  $\land$  here.

Caley Howland Lecture 11 03/23/2020 16 / 29


- How should we understand "It is improbable that C is false while P is true"? The "while" can't mean  $\land$  here.
- What we need: How probable is the conclusion of the argument given that its premises are true?

- How should we understand "It is improbable that C is false while P is true"? The "while" can't mean  $\land$  here.
- What we need: How probable is the conclusion of the argument given that its premises are true?
- A diagram may help:




Caley Howland Lecture 11 03/23/2020 16/29


- How should we understand "It is improbable that C is false while P is true"? The "while" can't mean ∧ here.
- What we need: How probable is the conclusion of the argument given that its premises are true?
- A diagram may help:



• In this diagram (where area is proportional to probability), the probability of  $P \land \neg C$  is low, but the probability of  $\neg C$  given that P is high. Why?




Caley Howland Lecture 11 03/23/2020 17/29




• Notice that—within the P-circle—most of the area is occupied by  $\neg C$ .

Caley Howland Lecture 11 03/23/2020 17 / 29



- Notice that—within the P-circle—most of the area is occupied by  $\neg C$ .
- Therefore,  $\neg C$  is probable, given that P is true.

 Catey Howland
 Lecture 11
 03/23/2020
 17/29




- Notice that—within the P-circle—most of the area is occupied by  $\neg C$ .
- Therefore,  $\neg C$  is probable, given that P is true.
- So, when evaluating an inductive argument P :: C, we need to think about **how probable** C **is, given that** P. The more probable P is (i.e., the less probable P is)—given that P—the stronger the argument is.


Caley Howland Lecture 11 03/23/2020 17/29

#### Two Arguments

Let's compare the strength of (1)  $P \lor Q : P$  with (2)  $P \lor \neg P : P$  using the method of the last slide.

• We can picture them as follows:





• Because Q is contingent,  $P \lor Q$  must have a smaller area/probability than  $P \lor \neg P$ , which is the entire box.

Caley Howland Lecture 11 03/23/2020 19/29

- Because Q is contingent,  $P \lor Q$  must have a smaller area/probability than  $P \lor \neg P$ , which is the entire box.
- Therefore, the proportion of P within  $P \lor Q$  must be greater than the proportion of P within  $P \lor \neg P$ .

Caley Howland Lecture 11 03/23/2020 19 / 29

- Because Q is contingent,  $P \lor Q$  must have a smaller area/probability than  $P \lor \neg P$ , which is the entire box.
- Therefore, the proportion of P within  $P \lor Q$  must be greater than the proportion of P within  $P \lor \neg P$ .
- So, (1) must be stronger than (2).

 Cately Howland
 Lecture 11
 03/23/2020
 19/29

- Next, consider this example (borrowed from Skyrms):
  - (P) There is a 2000-year-old man in Cleveland.
     ∴(C) There is a 2000-year-old man in with three heads in Cleveland.

Caley Howland Lecture 11 03/23/2020 20 / 29

- Next, consider this example (borrowed from Skyrms):
  - (P) There is a 2000-year-old man in Cleveland.
     ∴(C) There is a 2000-year-old man in with three heads in Cleveland.
- In this case, *P* and *C* are each—individually—highly improbable.

 Caley Howland
 Lecture 11
 03/23/2020
 20 / 29

- Next, consider this example (borrowed from Skyrms):
  - (P) There is a 2000-year-old man in Cleveland.
     ∴(C) There is a 2000-year-old man in with three heads in Cleveland.
- In this case, *P* and *C* are each—individually—highly improbable.
- But we want to know: How probable is C given that P is true?

 Caley Howland
 Lecture 11
 03/23/2020
 20/29

- Next, consider this example (borrowed from Skyrms):
  - (P) There is a 2000-year-old man in Cleveland.
     ∴(C) There is a 2000-year-old man in with three heads in Cleveland.
- In this case, P and C are each—individually—highly improbable.
- But we want to know: How probable is C given that P is true?
- Similarly regarding another Skyrms example:
  - (P) There is a man in Cleveland who is 1999 years and 11-months-old and in good health.
    - $\therefore$  (C) No man will live to be 2000 years old.

Caley Howland Lecture 11 03/23/2020 20/29

- Next, consider this example (borrowed from Skyrms):
  - (P) There is a 2000-year-old man in Cleveland.
     ∴(C) There is a 2000-year-old man in with three heads in Cleveland.
- In this case, P and C are each—individually—highly improbable.
- But we want to know: How probable is C given that P is true?
- Similarly regarding another Skyrms example:
  - (P) There is a man in Cleveland who is 1999 years and 11-months-old and in good health.
    - $\therefore$  (C) No man will live to be 2000 years old.
- Let's do some inferential reasoning.

 Caley Howland
 Lecture 11
 03/23/2020
 20/29

#### So Far...

- So far, we've seen two proposals for understanding the notion of inductive strength.
  - ▶ **Proposal** #1. An argument P ∴ C is inductively strong just in case the claim  $P \land \neg C$  is improbable.

 Cately Howland
 Lecture 11
 03/23/2020
 21/29

#### So Far...

- So far, we've seen two proposals for understanding the notion of inductive strength.
  - ▶ **Proposal** #1. An argument P ∴ C is inductively strong just in case the claim  $P \land \neg C$  is improbable.
- Proposal 1 doesn't work, since an argument will be judged as strong if P is improbable (or C is probable). Skyrms then gives:


Caley Howland Lecture 11 03/23/2020 21/29

#### So Far...

- So far, we've seen two proposals for understanding the notion of inductive strength.
  - ▶ **Proposal** #1. An argument P ∴ C is inductively strong just in case the claim  $P \land \neg C$  is improbable.
- Proposal 1 doesn't work, since an argument will be judged as strong if P is improbable (or C is probable). Skyrms then gives:
  - Proposal #2. An argument P ∴ C is inductively strong just in case C is probable, given that P is true, i.e., if most P-worlds are C-worlds.

◆ロ ト ◆ 昼 ト ◆ 豆 ト ◆ 豆 ・ 夕 Q ○

 Proposal #1 looks at the size of the shaded region, relative to the size of the box (of all possible worlds). Proposal #2 looks at the size of the shaded region, relative to the size of the P-circle (of just the P-worlds).



 Caley Howland
 Lecture 11
 03/23/2020
 22 / 29

 It is easiest to contrast these proposals by thinking about games of chance, where the probabilities are known (and uncontroversial). E.g.,

Caley Howland Lecture 11 03/23/2020 23/29

- It is easiest to contrast these proposals by thinking about games of chance, where the probabilities are known (and uncontroversial). E.g.,
- Suppose I'm going to pick a card c at random from a standard deck.

 Caley Howland
 Lecture 11
 03/23/2020
 23/29

- It is easiest to contrast these proposals by thinking about games of chance, where the probabilities are known (and uncontroversial). E.g.,
- Suppose I'm going to pick a card c at random from a standard deck.
  - ► Let P = "c is a black card", and C = "c is a spade".

Caley Howland Lecture 11 03/23/2020 23 / 29

• Proposal #1. Calculate the probability of  $P \land \neg C$ . If this is low (less than 1/2), then Proposal #1 says the argument "P  $\therefore$  C" is strong.

Caley Howland Lecture 11 03/23/2020 24/29

- Proposal #1. Calculate the probability of  $P \land \neg C$ . If this is low (less than 1/2), then Proposal #1 says the argument "P  $\therefore$  C" is strong.
  - The probability of P ∧ ¬C = the probability that c is black but not a spade. This is just the probability that c is a club = 1/4 < 1/2 . So, according to Proposal #1, "P ∴ C" is a strong argument.

- Proposal #1. Calculate the probability of  $P \land \neg C$ . If this is low (less than 1/2), then Proposal #1 says the argument "P  $\therefore$  C" is strong.
  - The probability of P ∧ ¬C = the probability that c is black but not a spade. This is just the probability that c is a club = 1/4 < 1/2 . So, according to Proposal #1, "P ∴ C" is a strong argument.
- Proposal #2. Calculate the probability of  $\neg C$ , given that P. If this is low (less than 1/2), then Proposal #2 says "P :: C" is strong.

Caley Howland Lecture 11 03/23/2020 24/29

- Proposal #1. Calculate the probability of  $P \land \neg C$ . If this is low (less than 1/2), then Proposal #1 says the argument "P  $\therefore$  C" is strong.
  - The probability of P ∧ ¬C = the probability that c is black but not a spade. This is just the probability that c is a club = 1/4 < 1/2 . So, according to Proposal #1, "P ∴ C" is a strong argument.
- Proposal #2. Calculate the probability of  $\neg C$ , given that P. If this is low (less than 1/2), then Proposal #2 says "P  $\therefore$  C" is strong.
  - ► The probability of  $\neg C$ , given that P = the proportion of black cards that are clubs = 1/2. So, Proposal #2 says that "P : C" is not strong.

• The improbability of  $\neg C$  given that P is a much better guide to the inductive strength of "P  $\therefore$  C" than the improbability of  $P \land \neg C$ .

Caley Howland Lecture 11 03/23/2020 25 / 29

- The improbability of  $\neg C$  given that P is a much better guide to the inductive strength of "P : C" than the improbability of  $P \land \neg C$ .
- But, there is still something lacking in Skyrms's second proposal.

Caley Howland Lecture 11 03/23/2020 25/29

- The improbability of  $\neg C$  given that P is a much better guide to the inductive strength of "P :: C" than the improbability of  $P \land \neg C$ .
- But, there is still something lacking in Skyrms's second proposal.
- This defect can be illustrated via the following inductive argument.

Caley Howland Lecture 11 03/23/2020 25/29

- The improbability of  $\neg C$  given that P is a much better guide to the inductive strength of "P  $\therefore$  C" than the improbability of  $P \land \neg C$ .
- But, there is still something lacking in Skyrms's second proposal.
- This defect can be illustrated via the following inductive argument.

### Horoscope

(P) John's horoscope is unfavorable.

Therefore, (C) John will not win the lottery.

#### Horoscope

(P) John's horoscope is unfavorable Therefore, (C) John will not win the lottery.

- The probability of  $\neg C$  given that P is very low (as is the probability that  $P \land \neg C$ ).
  - ► So, Proposal #2 (and Proposal #1) says "P : C" is strong.

Caley Howland Lecture 11 03/23/2020 26/29

### Horoscope

(P) John's horoscope is unfavorable Therefore, (C) John will not win the lottery.

- The probability of  $\neg C$  given that P is very low (as is the probability that  $P \land \neg C$ ).
  - ► So, Proposal #2 (and Proposal #1) says "P : C" is strong.
- But, intuitively, P is irrelevant to C, and so (intuitively) P does not provide evidence in favor of C. This suggests a third proposal, due to Branden Fitelson.

Caley Howland Lecture 11 03/23/2020 26 / 29

### Horoscope

(P) John's horoscope is unfavorable Therefore, (C) John will not win the lottery.

- The probability of  $\neg C$  given that P is very low (as is the probability that  $P \land \neg C$ ).
  - ► So, Proposal #2 (and Proposal #1) says "P : C" is strong.
- But, intuitively, P is irrelevant to C, and so (intuitively) P does not provide evidence in favor of C. This suggests a third proposal, due to Branden Fitelson.
  - ▶ **Proposal** #3 "P ∴ C" is strong just in case (1) the probability of  $\neg C$  given that P is low, and (2) P is *positively relevant* to C i.e., the probability of  $\neg C$  given that P is lower than the probability of  $\neg C$ .

Caley Howland Lecture 11 03/23/2020 26 / 29

• The third proposal adds a positive relevance requirement.

Caley Howland Lecture 11 03/23/2020 27 / 29

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.

Caley Howland Lecture 11 03/23/2020 27 / 29

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.
- The probability that c is a not a spade, given that c is black is 1/2. That is, the probability that  $\neg C$ , given that P is 1/2.

Caley Howland Lecture 11 03/23/2020 27 / 29

- The third proposal adds a positive relevance requirement.
- If we return to our card example, we can see how relevance works.
- The probability that c is a not a spade, given that c is black is 1/2. That is, the probability that  $\neg C$ , given that P is 1/2.
- This is not low (i.e., it is not less than 1/2). But, it is *lower* than the probability that c is not a spade (i.e., the probability of  $\neg C$ ), which is 3/4.

 Caley Howland
 Lecture 11
 03/23/2020
 27 / 29

- So, in this case, P is positively relevant to C
  - ► The probability of C given that  $P = \frac{1}{2}$ , which is greater than the probability of  $C = \frac{1}{4}$ .


ロト 4回ト 4 重ト 4 重ト 重 めのの

- So, in this case, P is positively relevant to C
  - ► The probability of C given that  $P = \frac{1}{2}$ , which is greater than the probability of  $C = \frac{1}{4}$ .
- Thus, the argument "P ∴ C" does not come out strong on proposal #3 (because it fails the low conditional probability requirement). But this argument does satisfy the positive relevance requirement.

Caley Howland Lecture 11 03/23/2020 28/29

- So, in this case, P is positively relevant to C
  - ► The probability of C given that  $P = \frac{1}{2}$ , which is greater than the probability of  $C = \frac{1}{4}$ .
- Thus, the argument "P ∴ C" does not come out strong on proposal #3 (because it fails the low conditional probability requirement). But this argument does satisfy the positive relevance requirement.
- With proposal #3 in hand, we can now explain how our "scale" of argument strength works.
  - ► It depends on Pr(C|P) and Pr(C|P) > Pr(C).

 Caley Howland
 Lecture 11
 03/23/2020
 28/29





Caley Howland Lecture 11 03/23/2020 29 / 29