

# RWorksheet\_Pama#3a

Miko Pama

2023-10-16

#1. VECTORS #A.

```
first_eleven <- LETTERS[1:11]
first_eleven
```

```
## [1] "A" "B" "C" "D" "E" "F" "G" "H" "I" "J" "K"
```

#B.

```
odd_letters <- LETTERS[seq(1,25,2)]
odd_letters
```

```
## [1] "A" "C" "E" "G" "I" "K" "M" "O" "Q" "S" "U" "W" "Y"
```

#C.

```
vowels <- c(letters[which(letters %in% c("a","e","i","o","u"))],
            LETTERS[which(LETTERS %in% c("A","E","I","O","U"))])
vowels
```

```
## [1] "a" "e" "i" "o" "u" "A" "E" "I" "O" "U"
```

#D.

```
last5letter <- tail(letters, 5)
last5letter
```

```
## [1] "v" "w" "x" "y" "z"
```

#E. {r number1e} letterbetween15to24 <- letters[15:24] letterbetween15to2

#NUMBER 2 #A.

```
city <- c("Tuguegarao City", "Manila", "Iloilo City", "Tacloban", "Samal Island", "Davao City")
city
```

```
## [1] "Tuguegarao City" "Manila"           "Iloilo City"      "Tacloban"
## [5] "Samal Island"    "Davao City"
```

#B.

```
temp <- c(42, 39, 34, 34, 30, 27)
temp
```

```
## [1] 42 39 34 34 30 27
```

#C.

```
df <- data.frame(city, temp)
df
```

```
##           city temp
## 1 Tuguegarao City  42
## 2           Manila  39
## 3      Iloilo City  34
## 4      Tacloban   34
## 5      Samal Island 30
## 6      Davao City  27
```

#D.

```
names(df) <- c("City", "Temperature")
df
```

```
##           City Temperature
## 1 Tuguegarao City        42
## 2           Manila        39
## 3      Iloilo City        34
## 4      Tacloban         34
## 5      Samal Island       30
## 6      Davao City        27
```

#E. {r number2e} str[(df)]

#F.

```
df[3:4,]
```

```
##           City Temperature
## 3 Iloilo City          34
## 4   Tacloban          34
```

#G.

```
highest_temp_city <- df$City[which.max(df$Temperature)]
lowest_temp_city  <- df$City[which.min(df$Temperature)]
highest_temp_city
```

```
## [1] "Tuguegarao City"
```

```
lowest_temp_city
```

```
## [1] "Davao City"
```

#2 MATRICES #A.

```
m <- matrix(c(1:8,11:14),ncol=4,nrow=3)
m
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    1    4    7   12
## [2,]    2    5    8   13
## [3,]    3    6   11   14
```

#B.

```
matrix(c(1:8,11:14),ncol=4,nrow=3) * 2
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    2    8   14   24
## [2,]    4   10   16   26
## [3,]    6   12   22   28
```

```
#C.
```

```
matrix(c(1:8,11:14),ncol=4,nrow=3)[2,]
```

```
## [1]  2  5  8 13
```

```
#D.
```

```
matrix(c(1:8,11:14),ncol=4,nrow=3)[1:2,c(3,4)]
```

```
##      [,1] [,2]
## [1,]    7   12
## [2,]    8   13
```

```
#E.
```

```
matrix(c(1:8,11:14),ncol=4,nrow=3)[3,c(2,3)]
```

```
## [1]  6 11
```

```
#F.
```

```
matrix(c(1:8,11:14),ncol=4,nrow=3)[,4]
```

```
## [1] 12 13 14
```

```
#G.
```

```
mat <- matrix(c(1:8,11:14)*2,ncol=4,nrow=3)
rownames(mat) <- c("isa","dalawa","tatlo")
colnames(mat) <- c("uno","dos","tres","quatro")
mat
```

```
##      uno dos tres quatro
## isa      2  8  14     24
## dalawa   4 10  16     26
## tatlo    6 12  22     28
```

```
#H.
```

```
dim(m) <- c(6,2)
m
```

```
##      [,1] [,2]
## [1,]    1    7
## [2,]    2    8
## [3,]    3   11
## [4,]    4   12
## [5,]    5   13
## [6,]    6   14
```

```
#3. ARRAYS #A.
```

```
values <- c(1, 2, 3, 6, 7, 8, 9, 0, 3, 4, 5, 1)
array_3d <- array(values, dim = c(2, 4, 3))
values
```

```
## [1] 1 2 3 6 7 8 9 0 3 4 5 1
```

```
array_3d
```

```
## , , 1
```

```
##
```

```
##      [,1] [,2] [,3] [,4]
```

```
## [1,] 1 3 7 9
## [2,] 2 6 8 0
##
## , , 2
##
##      [,1] [,2] [,3] [,4]
## [1,] 3 5 1 3
## [2,] 4 1 2 6
##
## , , 3
##
##      [,1] [,2] [,3] [,4]
## [1,] 7 9 3 5
## [2,] 8 0 4 1
```

#B.

*#B: The array has three dimensions.*

#C.

```
dimnames(array_3d) <- list(c("a", "b"), LETTERS[1:4], c("1st-Dimensional Array", "2nd-Dimensional Array", "3rd-Dimensional Array"))
array_3d
```

```
## , , 1st-Dimensional Array
##
##   A B C D
## a 1 3 7 9
## b 2 6 8 0
##
## , , 2nd-Dimensional Array
##
##   A B C D
## a 3 5 1 3
## b 4 1 2 6
##
## , , 3rd-Dimensional Array
##
##   A B C D
## a 7 9 3 5
## b 8 0 4 1
```