


a) **Initial labs** in all COVID patients: CBC with diff, retic, DIC panel*, AT3, ferritin, LDH, CMP
b) **Repeat labs**: Daily CBC and DIC panel*

* DIC panel at UNC = PT, aPTT, fibrinogen, D-dimer

COVID-19 patients: Anticoagulation Management at UNC – Regular floor or ICU

All patients receive anticoagulation, unless bleeding contraindications - **thoroughly consider and factor into the decision-making a patient's bleeding risk factors**



After hospital discharge³
(applies to Groups B and C)

COVID-19 patients discharged from hospital meeting below criteria: Consider *Xarelto*® 10 mg once daily (FDA-approved for outpatient indication), *Eliquis*® 2.5 mg bid (off-label), or prophylactic heparin product for 30 days and until mobile.
Any decision to use post-discharge VTE prophylaxis should consider an individual patient's VTE risk factors, including reduced mobility, bleeding risks, and feasibility.

- Modified IMPROVE-VTE score ≥4; or
- Modified IMPROVE-VTE score ≥2 and D-dimer level >2 times the upper limit of normal; or
- Age ≥75 years; or
- Age >60 yrs and D-dimer level >2x upper limit of normal; or
- Age 40-60 yrs, and D-dimer level >2x upper limit of normal, and previous VTE or cancer.

Modified IMPROVE VTE risk score	
Previous VTE	3
Known thrombophilia	2
Current lower limb paralysis or paresis	2
History of cancer (Last 5yr, in remission)	2
ICU/CCU stay	1
Complete immobilization, ≥ 1 d	1
Age ≥60 y	1

Pregnancy / lactation in COVID+ patients:

- Details: UNC Maternal-Fetal-Medicine guidance: <https://bit.ly/2X9YnnG>
- Preferred anticoagulants during pregnancy are heparin compounds.
- Management during labor and delivery: Collaboration MFM and Hematology.

¹ If a Group B patient's D-dimer on daily f/u testing falls to ≤ 2,500 ng/mL, consider keeping patient at intermediate-intensity dosing.
² If a Group C patient's D-dimer on daily f/u testing increases to > 2,500 ng/mL, consider repeat testing to determine a trend and moving to intermediate-intensity dosing.
³ Further guidance on discharge, modified IMPROVE risk score, or pregnancy considerations per NIH COVID-19 Consensus Anticoagulation Guidelines
If there are management concerns, a hematology consult can be obtained.