

Supplement to “Statistical Inference for High-Dimensional Models via Recursive Online-Score Estimation”

Chengchun Shi, Rui Song and Wenbin Lu
Department of Statistics, North Carolina State University
and
Runze Li
Department of Statistics, Pennsylvania State University

In this supplementary article, we present proofs of Theorem 3.1, Theorem 3.2, Theorem 3.3, Lemma A.1, Lemma A.2, Lemma A.3 and Lemma B.1.

B Proofs

B.1 Proof of Theorem 3.1

We use a shorthand and write $\widehat{\mathcal{M}}_{j_0}^{(t)} = \widehat{\mathcal{M}}_{j_0}^{(-s_n)}$ for $t = 0, \dots, s_n - 1$. Let

$$\widehat{\Sigma}^* = \frac{1}{n} \sum_{i=1}^n \mathbf{X}_i b''(\mathbf{X}_i^T \beta_0) \mathbf{X}_i^T \quad \text{and} \quad \widehat{\Psi}^{(j)} = \frac{1}{n} \sum_{i=1}^n \mathbf{X}_i b'''(\mathbf{X}_i^T \beta_0) \mathbf{X}_i^T X_{i,j},$$

for any $j \in \{1, 2, \dots, p\}$. For any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$, define

$$\begin{aligned}
\boldsymbol{\omega}_{\mathcal{M},j_0} &= \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \boldsymbol{\Sigma}_{\mathcal{M},j_0}, & \sigma_{\mathcal{M},j_0}^2 &= \boldsymbol{\Sigma}_{j_0,j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}^T \boldsymbol{\Sigma}_{\mathcal{M},j_0}, \\
\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0} &= \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{-1} \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0}, & \hat{\sigma}_{\mathcal{M},j_0}^2 &= \widehat{\boldsymbol{\Sigma}}_{j_0,j_0} - \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0}^T \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}, \\
\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^* &= \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{*-1} \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0}^*, & \hat{\sigma}_{\mathcal{M},j_0}^{*2} &= \widehat{\boldsymbol{\Sigma}}_{j_0,j_0}^* - \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0}^{*T} \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^*, \\
\widetilde{\boldsymbol{\omega}}_{\mathcal{M},j_0} &= \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^* + \sum_{j=1}^p \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{*-1} \left(\widehat{\boldsymbol{\Psi}}_{\mathcal{M},j_0}^{(j)} + \widehat{\boldsymbol{\Psi}}_{\mathcal{M},\mathcal{M}}^{(j)} \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^* \right) (\widetilde{\beta}_j - \beta_{0,j}), \\
\widehat{Z}_{t+1,j_0}^* &= X_{t+1,j_0} - \widehat{\boldsymbol{\omega}}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*T} \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}, & \widetilde{Z}_{t+1,j_0} &= X_{t+1,j_0} - \widetilde{\boldsymbol{\omega}}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}, \\
\hat{\xi}_{\mathcal{M},j_0}^{(j)} &= \widehat{\boldsymbol{\Psi}}_{j_0,j_0}^{(j)} - \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^{*T} \left(2\widehat{\boldsymbol{\Psi}}_{\mathcal{M},j_0}^{(j)} + \widehat{\boldsymbol{\Psi}}_{\mathcal{M},\mathcal{M}}^{(j)} \widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}^* \right), \\
\tilde{\sigma}_{\mathcal{M},j_0}^2 &= \hat{\sigma}_{\mathcal{M},j_0}^{*2} + \sum_{j=1}^p \hat{\xi}_{\mathcal{M},j_0}^{(j)} (\widetilde{\beta}_j - \beta_{0,j}).
\end{aligned}$$

Here, $\widetilde{\boldsymbol{\omega}}_{\mathcal{M},j_0}$ and $\tilde{\sigma}_{\mathcal{M},j_0}$ correspond to first-order approximations of $\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0}$ and $\hat{\sigma}_{\mathcal{M},j_0}$ around β_0 . We introduce the following lemmas before proving Theorem 3.1. The proof of Lemma B.1 is given in Section B.7 of the supplementary material.

Lemma B.1 *Under conditions in Theorem 3.1, we have*

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sigma_{\mathcal{M},j_0} \geq \sqrt{\bar{c}}, \quad \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \leq \sqrt{c_0/\bar{c}}, \quad (\text{B.1})$$

where \bar{c} and c_0 are defined in Condition (A2*) and (A3*). Besides, the following events hold with probability tending to 1,

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \|\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \leq \bar{c}_0 \left(\frac{\sqrt{\kappa_n \log p} + \kappa_n}{\sqrt{n}} + \eta_n \right), \quad (\text{B.2})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M},j_0} - \sigma_{\mathcal{M},j_0}| \leq \bar{c}_0 \left(\frac{\sqrt{\kappa_n \log p} + \kappa_n}{\sqrt{n}} + \eta_n \right), \quad (\text{B.3})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \|\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0} - \widetilde{\boldsymbol{\omega}}_{\mathcal{M},j_0}\|_2 \leq \bar{c}_0 \eta_n^2, \quad \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M},j_0}^2 - \tilde{\sigma}_{\mathcal{M},j_0}^2| \leq \bar{c}_0 \eta_n^2, \quad (\text{B.4})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M},j_0}^2 - \hat{\sigma}_{\mathcal{M},j_0}^{*2}| \leq \bar{c}_0 \eta_n, \quad (\text{B.5})$$

for some constant $\bar{c}_0 > 0$. Moreover, we have

$$\sum_{t=0}^{n-1} \frac{\widetilde{Z}_{t+1,j_0} \varepsilon_{t+1}}{\sqrt{n}} \left(\frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\widetilde{\beta}_j - \beta_{0,j})}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}^{*3}} \right) = \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} + o_p(1).$$

For simplicity, we only consider the case where $l = 1$. When $l > 1$, assume we've shown the asymptotic normality of $\hat{\beta}_{j_0}^{(l-1)}$. Under the given conditions, we can show $\Gamma_n^{*,(l-2)}$ is lower bounded by $\sqrt{\bar{c}}/2$, with probability tending to 1. This implies $\hat{\beta}_{j_0}^{(l-1)}$ converges to β_{0,j_0} at a rate of $O_p(n^{-1/2})$. As a result, the estimator $\hat{\beta}^{(l-1)} = \tilde{\beta} + \mathbf{e}_{j_0,p}(\hat{\beta}_{j_0}^{(l-1)} - \tilde{\beta}_{j_0})$ also satisfies the conditions in (A5*). The asymptotic normality of $\hat{\beta}_{j_0}^{(l)}$ can be similarly derived.

In the following, we omit the superscript and write $\hat{\beta}_{j_0}^{(1)}$, $\Gamma_n^{*,(0)}$ as $\hat{\beta}_{j_0}$ and Γ_n^* . Let $\varepsilon_i = Y_i - \mu(\mathbf{X}_i^T \beta_0)$ for $i = 0, 1, \dots, n$. By definition, we have

$$\sqrt{n}\Gamma_n^*(\hat{\beta}_{j_0} - \tilde{\beta}_{j_0}) = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}}} \widehat{Z}_{t+1,j_0} \left\{ Y_{t+1} - \mu \left(X_{t+1} \tilde{\beta}_{0,j_0} + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \right\} \quad (\text{B.6})$$

By Condition (A1), we can show the following events occur with probability tending to 1,

$$\mathcal{M}_{j_0} \subseteq \widehat{\mathcal{M}}_{j_0}^{(t)}, \quad |\widehat{\mathcal{M}}_{j_0}^{(t)}| \leq \kappa_n, \quad t = 0, \dots, n-1. \quad (\text{B.7})$$

Besides, similar to (28) and (31), we can show

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M},j_0}^{*2} - \sigma_{\mathcal{M},j_0}^2| \leq \bar{c}_0 \left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n}{\sqrt{n}} \right), \quad (\text{B.8})$$

for some constant $\bar{c}_0 > 0$, and

$$\min_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \hat{\sigma}_{\mathcal{M},j_0} \geq \sqrt{\bar{c}}/2 \quad \text{and} \quad \min_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \hat{\sigma}_{\mathcal{M},j_0}^* \geq \sqrt{\bar{c}}/2, \quad (\text{B.9})$$

with probability tending to 1.

Under the events defined in (B.7), we have for $t = 0, 1, \dots, n-1$,

$$\mathbf{X}_{t+1}^T \beta_0 = X_{t+1,j_0} \beta_{0,j_0} + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \beta_{0,\widehat{\mathcal{M}}_{j_0}^{(t)}}.$$

Hence, using a second order Taylor expansion, we have

$$\begin{aligned} \mu(\mathbf{X}_{t+1}^T \beta_0) &= \mu \left(X_{t+1,j_0} \beta_{0,j_0} + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \beta_{0,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \\ &= \mu \left(X_{t+1,j_0} \tilde{\beta}_{j_0} + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) + b'' \left(X_{t+1,j_0} \tilde{\beta}_{j_0} + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \\ &\times \left(X_{t+1,j_0} (\beta_{0,j_0} - \tilde{\beta}_{j_0}) + \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0,\widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}}) \right) \\ &+ \frac{1}{2} b''' \left(\mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}^* \right) \left(\mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}) \right)^2, \end{aligned}$$

for some $\tilde{\beta}_t^* \in \mathbb{R}^{1+|\widehat{\mathcal{M}}_{j_0}^{(t)}|}$ lying on the line segment joining $\beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}$ and $\tilde{\beta}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}$. Let R_t^* be the second order Remainder term. Under the events defined in (B.7), we have

$$\begin{aligned} & \left| \mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_t^* \right| \leq \left| \mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right| + \left| \mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_t^*) \right| \\ &= \left| \mathbf{X}_{t+1}^T \beta_0 \right| + \left| \mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_t^*) \right| \leq \bar{\omega} + \omega_0 \left\| \beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_t^* \right\|_1 \\ &\leq \bar{\omega} + \omega_0 \left\| \beta_0 - \tilde{\beta} \right\|_1, \end{aligned}$$

where the second inequality is due to Condition (A4*). By Condition (A5*), we have with probability tending to 1,

$$\omega_0 \|\tilde{\beta} - \beta_0\|_1 \leq \omega_0 \eta_n \leq \bar{\omega}.$$

Since $b'''(\cdot)$ is continuous, $\sup_{|z| \leq 2\bar{\omega}} |b'''(z)|$ is upper bounded by some constant $c_* > 0$. Therefore, we have with probability tending to 1 that

$$\max_{t=0,\dots,n-1} \left| b''' \left(\mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_t^* \right) \right| \leq c_*. \quad (\text{B.10})$$

By Condition (A4*) and (A5*), this further implies that we have with probability tending to 1,

$$\begin{aligned} \max_{t \in \{0,\dots,n-1\}} |R_t^*| &\leq \frac{c_*}{2} \max_{t \in \{0,\dots,n-1\}} \left| \mathbf{X}_{t+1,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0,\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_t^*) \right|^2 \\ &\leq \frac{c_* \omega_0^2}{2} \left\| \beta_0 - \tilde{\beta} \right\|_1^2 \leq \frac{c_* \omega_0^2 \eta_n^2}{2}. \end{aligned} \quad (\text{B.11})$$

We now prove

$$\Pr \left\{ \max_{t \in [0,\dots,n-1]} \left| \widehat{Z}_{t+1,j_0} \right| \leq \omega_0 \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}} \right) \right\} \rightarrow 1. \quad (\text{B.12})$$

Note that

$$\max_{t \in [0,\dots,n-1]} \left| \widehat{Z}_{t+1,j_0} \right| \leq \max_{t \in [0,\dots,n-1]} |X_{t+1,j_0}| + \max_{t \in [0,\dots,n-1]} \left\| \widehat{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right\|_2 \left\| \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right\|_2.$$

By Condition (A1*) and (A4*), we have almost surely,

$$\max_{t \in [0,\dots,n-1]} \left| \widehat{Z}_{t+1,j_0} \right| \leq \omega_0 + \sqrt{\kappa_n} \omega_0 \max_{\substack{\mathcal{M} \subseteq \mathbb{I} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \widehat{\omega}_{\mathcal{M},j_0} \right\|_2. \quad (\text{B.13})$$

By (B.1) and (B.2), we have with probability tending to 1,

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \|\hat{\omega}_{\mathcal{M}, j_0}\|_2 \leq 2\sqrt{c_0/\bar{c}}, \quad (\text{B.14})$$

for sufficiently large n . Combining (B.14) together with (B.13) yields (B.12). Under the events defined in (B.7), (B.9), (B.11) and (B.12), we have

$$\begin{aligned} & \left| \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} \widehat{Z}_{t+1, j_0} \left\{ Y_{t+1} - \mu \left(X_{t+1} \tilde{\beta}_{0, j_0} + \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \right\} \right. \\ & - \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0} \varepsilon_{t+1}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} - \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} b'' \left(\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \\ & \times \left. \left(X_{t+1, j_0} (\beta_{0, j_0} - \tilde{\beta}_{j_0}) + \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T (\beta_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} - \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}}) \right) \right| \\ & \leq \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} |R_t^*| |\widehat{Z}_{t+1, j_0}| \leq \frac{\sqrt{n} c_*}{\sqrt{\bar{c}}} \omega_0^3 \eta_n^2 \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}} \right). \end{aligned}$$

By Condition (A5*), we have $\sqrt{n} \kappa_n \eta_n^2 = o(1)$. Hence, we've shown

$$\begin{aligned} & \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} \widehat{Z}_{t+1, j_0} \left\{ Y_{t+1} - \mu \left(X_{t+1} \tilde{\beta}_{0, j_0} + \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \right\} \\ & = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} b'' \left(\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \left(\tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \beta_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \\ & + \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0} \varepsilon_{t+1}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} + \sqrt{n} \Gamma_n^* (\beta_{0, j_0} - \tilde{\beta}_{j_0}) + o_p(1). \end{aligned}$$

In view of (B.6), we have

$$\begin{aligned} & \sqrt{n} \Gamma_n^* (\hat{\beta}_{j_0} - \beta_{0, j_0}) = o_p(1) + \underbrace{\frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0} \varepsilon_{t+1}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}}}_{I_1} \\ & + \underbrace{\frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} b'' \left(\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\beta}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \left(\tilde{\beta}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \beta_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right)}_{I_2}. \end{aligned}$$

In the following, we break the proof into two steps. In the first step, we prove $I_2 = o_p(1)$. In the second step, we show $I_1 \xrightarrow{d} N(0, \phi_0)$. This implies $\sqrt{n} \Gamma_n^* (\hat{\beta}_{j_0} - \beta_{0, j_0}) \xrightarrow{d} N(0, \phi_0)$. By

Condition (A7*), $\hat{\phi}$ is consistent to ϕ_0 . It follows from Slutsky's theorem that

$$\frac{\sqrt{n}\Gamma_n^*(\hat{\beta}_{j_0} - \beta_{0,j_0})}{\hat{\phi}^{1/2}} \xrightarrow{d} N(0, 1).$$

The proof is hence completed.

Step 1: Under the events defined in (B.7), using a first order Taylor expansion, we have

$$\begin{aligned} & b'' \left(\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\boldsymbol{\beta}}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) = b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \\ & + \underbrace{\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \left(\tilde{\boldsymbol{\beta}}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) b''' \left(\mathbf{X}_{t+1, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \tilde{\boldsymbol{\beta}}_t^{**} \right)}_{R_t^{**}}, \end{aligned}$$

for some $\tilde{\boldsymbol{\beta}}_t^{**} \in \mathbb{R}^{1+|\widehat{\mathcal{M}}_{j_0}^{(t)}|}$ lying on the line segment joining $\boldsymbol{\beta}_{0, \{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}$ and $\tilde{\boldsymbol{\beta}}_{\{j_0\} \cup \widehat{\mathcal{M}}_{j_0}^{(t)}}$. Using similar arguments in (B.10) and (B.11), we can show the remainder term R_t^{**} satisfies

$$\max_{t \in [0, \dots, n-1]} |R_t^{**}| \leq c_{**} \omega_0 \eta_n, \quad (\text{B.15})$$

for some constant $c_{**} > 0$, with probability tending to 1. Let

$$I_2^* = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0}}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \left(\tilde{\boldsymbol{\beta}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right),$$

we have

$$|I_2 - I_2^*| \leq \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{|\widehat{Z}_{t+1, j_0}|}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} |R_t^{**}| \left\| \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right\|_{\infty} \left\| \tilde{\boldsymbol{\beta}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right\|_1.$$

By (B.9), (B.12), (B.15), Condition (A4*) and (A5*), we have with probability tending to 1,

$$|I_2 - I_2^*| \leq \sqrt{n} \frac{2\omega_0}{\sqrt{\bar{c}}} \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}} \right) c_{**} \omega_0 \eta_n^2 = o(1).$$

This implies $I_2 = I_2^* + o_p(1)$. It suffices to show $I_2^* = o_p(1)$.

Similar to the proof of Theorem 2.1, by (B.2), (B.7) and (B.9), we can show the following event occurs with probability tending to 1,

$$\begin{aligned} & \left| \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1, j_0} - Z_{t+1, j_0}}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}} b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \left(\tilde{\boldsymbol{\beta}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0, \widehat{\mathcal{M}}_{j_0}^{(t)}} \right) \right| \quad (\text{B.16}) \\ & \leq c_{***} \frac{2\sqrt{n}}{\sqrt{\bar{c}}} \left(\frac{\sqrt{\kappa_n \log p} + \kappa_n}{\sqrt{n}} + \eta_n \right) \sqrt{\kappa_n} \omega_0^2 \eta_n, \end{aligned}$$

for some constant $c_{***} > 0$, where $Z_{t+1,j_0} = X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}}$.

Under the conditions $(\kappa_n \sqrt{\log p} + \kappa_n^{3/2})\eta_n \rightarrow 0$ and $\sqrt{n\kappa_n}\eta_n^2 \rightarrow 0$ in (A5*), (B.16) is $o_p(1)$. Similarly, we can show

$$\frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \left(\frac{Z_{t+1,j_0}}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} - \frac{Z_{t+1,j_0}}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \right) b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \left(\tilde{\boldsymbol{\beta}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) = o_p(1).$$

This together with (B.16) implies $I_2^* = I_2^{**} + o_p(1)$, where

$$I_2^{**} = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{Z_{t+1,j_0}}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \left(\tilde{\boldsymbol{\beta}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}} - \boldsymbol{\beta}_{0,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right).$$

Note that I_2^{**} can be further bounded from above by $\max_j |I_{2,j}| \|\tilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|_1$ where

$$I_{2,j} = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \frac{Z_{t+1,j_0}}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} b''(\mathbf{X}_{t+1}^T \boldsymbol{\beta}_0) \mathbf{X}_{t+1,j} I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}).$$

Similar to Lemma A.3, we can show $\max_j |I_{2,j}| = O_p(\log p)$. This together with Condition (A5*) implies $\max_j |I_{2,j}| \|\tilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|_1 = o_p(1)$ and hence $I_2^{**} = o_p(1)$. This proves $I_2 = o_p(1)$.

Step 2: By Taylor's theorem, we have for any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$,

$$\frac{1}{\hat{\sigma}_{\mathcal{M},j_0}} - \frac{1}{\hat{\sigma}_{\mathcal{M},j_0}^*} + \frac{\hat{\sigma}_{\mathcal{M},j_0}^2 - \hat{\sigma}_{\mathcal{M},j_0}^{*2}}{\hat{\sigma}_{\mathcal{M},j_0}^{*3}} = \frac{(\hat{\sigma}_{\mathcal{M},j_0}^2 - \hat{\sigma}_{\mathcal{M},j_0}^{*2})^2}{2\{\rho_{\mathcal{M}}\hat{\sigma}_{\mathcal{M},j_0} + (1 - \rho_{\mathcal{M}})\hat{\sigma}_{\mathcal{M},j_0}^*\}^5},$$

for some $0 < \rho_{\mathcal{M}} < 1$. By (B.5) and (B.9), the second-order remainder term satisfies

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left| \frac{(\hat{\sigma}_{\mathcal{M},j_0}^2 - \hat{\sigma}_{\mathcal{M},j_0}^{*2})^2}{2\{\rho_{\mathcal{M}}\hat{\sigma}_{\mathcal{M},j_0} + (1 - \rho_{\mathcal{M}})\hat{\sigma}_{\mathcal{M},j_0}^*\}^5} \right| \leq \frac{16\bar{c}_0^2\eta_n^2}{\bar{c}^{5/2}}, \quad (\text{B.17})$$

with probability tending to 1.

Besides, it follows from (B.4) and (B.9) that

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left| \frac{\hat{\sigma}_{\mathcal{M},j_0}^2 - \hat{\sigma}_{\mathcal{M},j_0}^{*2} - \sum_j \hat{\xi}_{\mathcal{M},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\hat{\sigma}_{\mathcal{M},j_0}^{*3}} \right| = \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left| \frac{\hat{\sigma}_{\mathcal{M},j_0}^2 - \tilde{\sigma}_{\mathcal{M},j_0}^2}{\hat{\sigma}_{\mathcal{M},j_0}^{*3}} \right| \leq \frac{8\bar{c}_0\eta_n^2}{\bar{c}^{3/2}},$$

with probability tending to 1. Combining this together with (B.17) yields

$$\Pr \left(\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left| \frac{1}{\hat{\sigma}_{\mathcal{M},j_0}} - \frac{1}{\hat{\sigma}_{\mathcal{M},j_0}^*} + \frac{\sum_j \hat{\xi}_{\mathcal{M},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\hat{\sigma}_{\mathcal{M},j_0}^{*3}} \right| \leq \bar{c}_1\eta_n^2 \right) \rightarrow 1,$$

for some constant $\bar{c}_1 > 0$. By Condition (A1*) and (B.12), we have

$$\begin{aligned} & \frac{1}{\sqrt{n}} \left| \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1,j_0} \varepsilon_{t+1}}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} - \sum_{t=0}^{n-1} \widehat{Z}_{t+1,j_0} \varepsilon_{t+1} \left(\frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} \right) \right| \\ & \leq \sqrt{n} \bar{c}_1 \eta_n^2 \max_t |\widehat{Z}_{t+1,j_0}| \frac{1}{n} \sum_{t=0}^{n-1} |\varepsilon_{t+1}| \leq \sqrt{n} \bar{c}_1 \eta_n^2 \omega_0 \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}} \right) \frac{1}{n} \sum_{t=0}^{n-1} |\varepsilon_{t+1}|, \end{aligned}$$

with probability tending to 1. By Condition (A6*) and Hölder's inequality, we have

$$\mathbb{E} \left(\frac{1}{n} \sum_{t=0}^{n-1} |\varepsilon_{t+1}| \right) = \mathbb{E} |\varepsilon_0| \leq (\mathbb{E} |\varepsilon_0|^3)^{1/3} = O(1).$$

Hence, it follows from Markov's inequality that $\sum_{t=0}^{n-1} |\varepsilon_{t+1}|/n = O_p(1)$. As a result, we have

$$\left| \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1,j_0} \varepsilon_{t+1}}{\sqrt{n}} \left(\frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} - \frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} + \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} \right) \right| = O_p(\sqrt{n} \kappa_n \eta_n^2) = o_p(1). \quad (\text{B.18})$$

where the last equality is due to Condition (A5*).

By (B.4), (B.8), (B.9) and Condition (A1*), (A4*), (A5*), (A6*), we can similarly show

$$\frac{1}{\sqrt{n}} \left| \sum_{t=0}^{n-1} (\widehat{Z}_{t+1,j_0} - \widetilde{Z}_{t+1,j_0}) \varepsilon_{t+1} \left(\frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} \right) \right| = o_p(1).$$

This together with (B.18) yields

$$\left| \sum_{t=0}^{n-1} \left\{ \frac{\widehat{Z}_{t+1,j_0} \varepsilon_{t+1}}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} - \widetilde{Z}_{t+1,j_0} \varepsilon_{t+1} \left(\frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} \right) \right\} \right| = o_p(\sqrt{n}).$$

Therefore, we've shown $I_1 = I_1^* + o_p(1)$ where

$$I_1^* = \frac{1}{\sqrt{n}} \sum_{t=0}^{n-1} \widetilde{Z}_{t+1,j_0} \varepsilon_{t+1} \left(\frac{1}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}^{*3}} \right).$$

In Lemma B.1, we further show I_1^* is equivalent to

$$I_1^{**} \equiv \sqrt{n} \sum_{t=0}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}}{\widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*}.$$

Hence, we have $I_1 = I_1^{**} + o_p(1)$. Unlike \tilde{Z}_{t+1,j_0} and $\tilde{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}$, \widehat{Z}_{t+1,j_0}^* and $\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*$ didn't depend on the initial estimator $\tilde{\beta}$. As a result, \widehat{Z}_{t+1,j_0}^* and $\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*$ are fixed given $\{\mathbf{X}_1, \dots, \mathbf{X}_n\}$ and $\widehat{\mathcal{M}}_{j_0}^{(t)}$. Following the arguments in the proof of Theorem 2.1, we can show

$$I_1^{**} = \frac{1}{\sqrt{n}} \sum_{t=s_n}^{n-1} \frac{Z_{t+1,j_0} \varepsilon_{t+1}}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} + o_p(1) \quad \text{and} \quad \frac{1}{\sqrt{n}} \sum_{t=s_n}^{n-1} \frac{Z_{t+1,j_0} \varepsilon_{t+1}}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \xrightarrow{d} N(0, \phi_0).$$

By Slutsky's theorem, we have $I_1 \xrightarrow{d} N(0, \phi_0)$. The proof is hence completed.

B.2 Proof of Theorem 3.2

Recall that $\mathbb{I} = [1, \dots, p]$ and $\mathbb{I}_{j_0} = \mathbb{I} - \{j_0\}$. By (17) and Lemma B.2, we have

$$\sqrt{n}L(\hat{\beta}_{j_0}^{DL}, \alpha) = 2z_{\frac{\alpha}{2}} \sqrt{\phi_0 \mathbf{e}_{j_0,p}^T \Sigma^{-1} \mathbf{e}_{j_0,p}} + o_p(1) = \frac{2z_{\frac{\alpha}{2}} \sqrt{\phi_0}}{\sigma_{\mathbb{I}_{j_0},j_0}} + o_p(1). \quad (\text{B.19})$$

It follows from (14) that

$$\sqrt{n}L(\hat{\beta}_{j_0}, \alpha) = \frac{2z_{\alpha/2} \sqrt{\phi_0}}{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}/n} + o_p(1). \quad (\text{B.20})$$

With some calculations, we have

$$\begin{aligned} & \frac{2z_{\alpha/2} \sqrt{\phi_0}}{\sigma_{\mathbb{I}_{j_0},j_0}} - \frac{2z_{\alpha/2} \sqrt{\phi_0}}{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}/n} \\ &= 2z_{\alpha/2} \sqrt{\phi_0} \frac{s_n \{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0} - \sigma_{\mathbb{I}_{j_0},j_0}\}/n + \sum_{t=s_n}^{n-1} \{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0} - \sigma_{\mathbb{I}_{j_0},j_0}\}/n}{\sigma_{\mathbb{I}_{j_0},j_0} \{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}/n\}}. \end{aligned} \quad (\text{B.21})$$

For any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$, we have

$$\begin{aligned} \sigma_{\mathcal{M},j_0}^2 &= \mathbb{E}|X_{0,j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^2 b''(\mathbf{X}_0^T \boldsymbol{\beta}_0) = \arg \min_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbb{E}|X_{0,j_0} - \mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2 b''(\mathbf{X}_0^T \boldsymbol{\beta}_0) \\ &\geq \arg \min_{\mathbf{a} \in \mathbb{R}^{p-1}} \mathbb{E}|X_{0,j_0} - \mathbf{a}^T \mathbf{X}_{0,\mathbb{I}_{j_0}}|^2 b''(\mathbf{X}_0^T \boldsymbol{\beta}_0) = \sigma_{\mathbb{I}_{j_0},j_0}^2. \end{aligned}$$

This shows $\sigma_{\mathcal{M},j_0} \geq \sigma_{\mathbb{I}_{j_0},j_0}$ for any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$. Hence, the numerator of the RHS of (B.21) is nonnegative.

On the other hand, by Condition (A4*), we have $|\mathbf{X}_0^T \boldsymbol{\beta}_0| \leq \bar{\omega}$ and hence $b''(\mathbf{X}_0^T \boldsymbol{\beta}_0) \leq \bar{k}$. Therefore,

$$\sigma_{\mathcal{M},j_0}^2 = \arg \min_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbb{E}|X_{0,j_0} - \mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2 b''(\mathbf{X}_0^T \boldsymbol{\beta}_0) \leq \bar{k} \mathbb{E}|X_{0,j_0}^2| \leq \bar{k} \sqrt{\mathbb{E}|X_{0,j_0}^4|} = \bar{k} \sqrt{c_0} \quad (\text{B.22})$$

where the last inequality is due to Condition (A3*). This implies

$$\begin{aligned} & \frac{2z_{\alpha/2}\sqrt{\phi_0}}{\sigma_{\mathbb{I}_{j_0},j_0}} - \frac{2z_{\alpha/2}\sqrt{\phi_0}}{s_n\sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}/n + \sum_{t=s_n}^{n-1}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}/n} \\ & \geq \frac{2z_{\alpha/2}\sqrt{\phi_0}}{\bar{k}\sqrt{c_0}} \left(s_n\{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0} - \sigma_{\mathbb{I}_{j_0},j_0}\}/n + \sum_{t=s_n}^{n-1}\{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0} - \sigma_{\mathbb{I}_{j_0},j_0}\}/n \right). \end{aligned} \quad (\text{B.23})$$

Besides, it follows from (B.22) that

$$\sigma_{\mathcal{M},j_0} - \sigma_{\mathbb{I}_{j_0},j_0} = \frac{\sigma_{\mathcal{M},j_0}^2 - \sigma_{\mathbb{I}_{j_0},j_0}^2}{\sigma_{\mathcal{M},j_0} + \sigma_{\mathbb{I}_{j_0},j_0}} \geq \frac{\sigma_{\mathcal{M},j_0}^2 - \sigma_{\mathbb{I}_{j_0},j_0}^2}{2\sqrt{\bar{k}}c_0^{1/4}},$$

for any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$. This together with (B.23) gives

$$\begin{aligned} & \frac{2z_{\alpha/2}\sqrt{\phi_0}}{\sigma_{\mathbb{I}_{j_0},j_0}} - \frac{2z_{\alpha/2}\sqrt{\phi_0}}{s_n\sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}/n + \sum_{t=s_n}^{n-1}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}/n} \\ & \geq \frac{z_{\alpha/2}\sqrt{\phi_0}}{\bar{k}^{3/2}c_0^{3/4}} \left(s_n\{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}^2 - \sigma_{\mathbb{I}_{j_0},j_0}^2\}/n + \sum_{t=s_n}^{n-1}\{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^2 - \sigma_{\mathbb{I}_{j_0},j_0}^2\}/n \right). \end{aligned} \quad (\text{B.24})$$

For any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$, define

$$\Omega_{\mathcal{M},j_0} = (\Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathbb{I}_{j_0} \cap \mathcal{M}^c} - \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c})^{-1}.$$

It follows from Lemma B.2 that

$$\begin{aligned} & \begin{pmatrix} \Sigma_{\mathcal{M}, \mathcal{M}} & \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \\ \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} & \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \end{pmatrix}^{-1} - \begin{pmatrix} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \\ & = \begin{pmatrix} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \Omega_{\mathcal{M}, j_0} \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} & -\Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \Omega_{\mathcal{M}, j_0} \\ -\Omega_{\mathcal{M}, j_0} \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} & \Omega_{\mathcal{M}, j_0} \end{pmatrix}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \Sigma_{\mathbb{I}_{j_0}, j_0}^T \Sigma_{\mathbb{I}_{j_0}, \mathbb{I}_{j_0}}^{-1} \Sigma_{\mathbb{I}_{j_0}, j_0} - \Sigma_{\mathcal{M}, j_0}^T \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, j_0} = (\Sigma_{j_0, \mathcal{M}}, \Sigma_{j_0, \mathbb{I}_{j_0} \cap \mathcal{M}^c}) \\ & \times \begin{pmatrix} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \Omega_{\mathcal{M}, j_0} \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} & -\Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c} \Omega_{\mathcal{M}, j_0} \\ -\Omega_{\mathcal{M}, j_0} \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathcal{M}} \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} & \Omega_{\mathcal{M}, j_0} \end{pmatrix} \begin{pmatrix} \Sigma_{\mathcal{M}, j_0} \\ \Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, j_0} \end{pmatrix} \\ & = (\Sigma_{j_0, \mathbb{I}_{j_0} \cap \mathcal{M}^c} - \omega_{\mathcal{M}, j_0}^T \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c}) \Omega_{\mathcal{M}, j_0} (\Sigma_{j_0, \mathbb{I}_{j_0} \cap \mathcal{M}^c} - \omega_{\mathcal{M}, j_0}^T \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c})^T \\ & \geq \lambda_{\min}(\Omega_{\mathcal{M}, j_0}) \|\Sigma_{j_0, \mathbb{I}_{j_0} \cap \mathcal{M}^c} - \omega_{\mathcal{M}, j_0}^T \Sigma_{\mathcal{M}, \mathbb{I}_{j_0} \cap \mathcal{M}^c}\|_2^2 = \lambda_{\min}(\Omega_{\mathcal{M}, j_0}) \|\xi_{\mathcal{M}, j_0}\|_2^2. \end{aligned}$$

By definition, we have

$$\lambda_{\min}(\Omega_{\mathcal{M}, j_0}) \geq \lambda_{\min} \left\{ (\Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathbb{I}_{j_0} \cap \mathcal{M}^c})^{-1} \right\} = \left\{ \lambda_{\max}(\Sigma_{\mathbb{I}_{j_0} \cap \mathcal{M}^c, \mathbb{I}_{j_0} \cap \mathcal{M}^c}) \right\}^{-1} \geq \left\{ \lambda_{\max}(\Sigma) \right\}^{-1} = \frac{1}{k_0},$$

and hence

$$\Sigma_{\mathbb{I}_{j_0}, j_0}^T \Sigma_{\mathbb{I}_{j_0}, \mathbb{I}_{j_0}}^{-1} \Sigma_{\mathbb{I}_{j_0}, j_0} - \Sigma_{\mathcal{M}, j_0}^T \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, j_0} \geq \frac{1}{k_0} \|\boldsymbol{\xi}_{\mathcal{M}, j_0}\|_2^2.$$

Note that we have

$$\sigma_{\mathcal{M}, j_0}^2 - \sigma_{\mathbb{I}_{j_0}, j_0}^2 = \Sigma_{j_0, j_0} - \Sigma_{\mathcal{M}, j_0}^c \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, j_0} - (\Sigma_{j_0, j_0} - \Sigma_{\mathbb{I}_{j_0}, j_0}^c \Sigma_{\mathbb{I}_{j_0}, \mathbb{I}_{j_0}}^{-1} \Sigma_{\mathbb{I}_{j_0}, j_0}).$$

This further implies

$$\sigma_{\mathcal{M}, j_0}^2 - \sigma_{\mathbb{I}_{j_0}, j_0}^2 \geq \frac{1}{k_0} \|\boldsymbol{\xi}_{\mathcal{M}, j_0}\|_2^2,$$

for any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$. By (B.24), we have

$$\frac{2z_{\alpha/2}\sqrt{\phi_0}}{\sigma_{\mathbb{I}_{j_0}, j_0}} - \frac{2z_{\alpha/2}\sqrt{\phi_0}}{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}/n} \geq \frac{\sqrt{\phi_0} z_{\alpha/2}}{\bar{k}^{3/2} c_0^{3/4} k_0} \left(\frac{s_n}{n} \|\boldsymbol{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}\|_2^2 + \frac{1}{n} \sum_{t=s_n}^{n-1} \|\boldsymbol{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}\|_2^2 \right).$$

In view of (B.19) and (B.20), we've shown

$$\sqrt{n}L(\hat{\beta}_{j_0}^{DL}, \alpha) \geq \sqrt{n}L(\hat{\beta}_{j_0}, \alpha) + \frac{\sqrt{\phi_0} z_{\alpha/2}}{\bar{k}^{3/2} c_0^{3/4} k_0} \left(\frac{s_n}{n} \|\boldsymbol{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}\|_2^2 + \frac{1}{n} \sum_{t=s_n}^{n-1} \|\boldsymbol{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}\|_2^2 \right) + o_p(1).$$

The proof is completed by noting that $\sqrt{n}L(\hat{\beta}_{j_0}^{DL}, \alpha) = \sqrt{n}L(\hat{\beta}_{j_0}^{DS}, \alpha) + o_p(1)$.

B.3 Proof of Theorem 3.3

Under the given conditions, using similar arguments in (29), we can show the following event occurs with probability tending to 1,

$$\widehat{\mathcal{M}}_{j_0}^{(-s_n)} = \widehat{\mathcal{M}}_{j_0}^{(s_n)} = \dots = \widehat{\mathcal{M}}_{j_0}^{(n)} = \mathcal{M}_{j_0}.$$

Under these events, we have

$$\frac{2z_{\alpha/2}\sqrt{\phi_0}}{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}/n} = \frac{2z_{\alpha/2}\sqrt{\phi_0}}{\sigma_{\mathcal{M}_{j_0}, j_0}}. \quad (\text{B.25})$$

By (14) and (21), for any sufficiently small $\varepsilon_0 > 0$, the following events occur with probability tending to 1,

$$\limsup_n \left| \sqrt{n}L(\hat{\beta}_{j_0}^{(l)}, \alpha) - \frac{2z_{\alpha/2}\sqrt{\phi_0}}{s_n \sigma_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}/n + \sum_{t=s_n}^{n-1} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}/n} \right| \leq \frac{\varepsilon_0}{2}, \quad (\text{B.26})$$

$$\limsup_n \left| \sqrt{n}L(\hat{\beta}_{j_0}^{oracle}, \alpha) - \frac{2z_{\alpha/2}\sqrt{\phi_0}}{\sigma_{\mathcal{M}_{j_0}, j_0}} \right| \leq \frac{\varepsilon_0}{2}. \quad (\text{B.27})$$

Conditional on the events defined in (B.25)-(B.27), we have

$$\limsup_n \left| \sqrt{n}L(\hat{\beta}_{j_0}^{(l)}, \alpha) - \sqrt{n}L(\hat{\beta}_{j_0}^{oracle}, \alpha) \right| \leq \varepsilon_0.$$

The proof is hence completed.

B.4 Proof of Lemma A.1

We first prove (24). Condition (A2) states that

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \inf_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|+1} \\ \|\mathbf{a}\|_2 \geq 1}} \mathbf{a}^T \boldsymbol{\Sigma}_{j_0 \cup \mathcal{M}, j_0 \cup \mathcal{M}} \mathbf{a} \geq \bar{c}. \quad (\text{B.28})$$

Note that

$$\begin{aligned} \sigma_{\mathcal{M}, j_0}^2 &= \boldsymbol{\Sigma}_{j_0, j_0} - \boldsymbol{\Sigma}_{j_0, \mathcal{M}} \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} \boldsymbol{\Sigma}_{\mathcal{M}, j_0} = (1, -\boldsymbol{\omega}_{\mathcal{M}, j_0}) \begin{pmatrix} \boldsymbol{\Sigma}_{j_0, j_0} & \boldsymbol{\Sigma}_{j_0, \mathcal{M}} \\ \boldsymbol{\Sigma}_{\mathcal{M}, j_0} & \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}} \end{pmatrix} \begin{pmatrix} 1 \\ -\boldsymbol{\omega}_{\mathcal{M}, j_0}^T \end{pmatrix} \\ &\geq \inf_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|+1} \\ \|\mathbf{a}\|_2 \geq 1}} \mathbf{a}^T \boldsymbol{\Sigma}_{j_0 \cup \mathcal{M}, j_0 \cup \mathcal{M}} \mathbf{a}. \end{aligned}$$

By (B.28), this implies

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sigma_{\mathcal{M}, j_0}^2 \geq \bar{c}, \quad (\text{B.29})$$

and hence

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sigma_{\mathcal{M}, j_0} \geq \sqrt{\bar{c}}.$$

By Cauchy-Schwarz inequality, Assumption (A3) implies that

$$\boldsymbol{\Sigma}_{j_0, j_0} = \mathbb{E}X_{0, j_0}^2 \leq \sqrt{\mathbb{E}X_{0, j_0}^4} \leq c_0.$$

In view of (B.29), this further implies that

$$\boldsymbol{\Sigma}_{j_0, \mathcal{M}} \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} \boldsymbol{\Sigma}_{\mathcal{M}, j_0} = \boldsymbol{\Sigma}_{j_0, j_0} - \sigma_{\mathcal{M}, j_0}^2 \leq c_0.$$

Note that $\boldsymbol{\Sigma}_{j_0, \mathcal{M}} \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} \boldsymbol{\Sigma}_{\mathcal{M}, j_0} = \boldsymbol{\omega}_{\mathcal{M}, j_0}^T \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}} \boldsymbol{\omega}_{\mathcal{M}, j_0}$. Hence, we have

$$\|\boldsymbol{\omega}_{\mathcal{M}, j_0}\|_2^2 \leq \frac{c_0}{\lambda_{\min}(\boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}})} \leq \frac{c_0}{\bar{c}},$$

where the last inequality is due to Condition (A2). Therefore, (24) is proven.

Consider (25). For any $a, b \in \mathbb{R}$, we have $(a + b)^4 \leq 8a^4 + 8b^4$. Therefore,

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \mathbb{E} |X_{0,j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4 \leq 8\mathbb{E} X_{0,j_0}^4 + \min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} 8\mathbb{E} |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4.$$

Under (A3), we have $\max_j \mathbb{E} X_{0,j}^4 \leq c_0$, $\mathbb{E} X_{0,j_0}^4 \leq c_0$. By (B.46), we have $\mathbb{E} |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|^4 \leq c_0^3/\bar{c}^2$ for all \mathcal{M} such that $|\mathcal{M}| \leq \kappa_n$. Moreover, by (24),

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{1}{\sigma_{\mathcal{M},j_0}} \leq \frac{1}{\sqrt{\bar{c}}}.$$

Thus, we have

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{1}{\sigma_{\mathcal{M},j_0}^4} \mathbb{E} |X_{0,j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4 \leq 8 \left(c_0 + \frac{c_0^3}{\bar{c}^2} \right) \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{1}{\sigma_{\mathcal{M},j_0}^4} \leq \frac{8}{\bar{c}^2} \left(c_0 + \frac{c_0^3}{\bar{c}^2} \right).$$

For any random variable Z with $\|Z\|_{\psi_2} \leq \omega$, it follows from the definition of the Orlicz norm that $\|Z\|_{\psi_1} \leq \omega^2$. Under Condition (A5), this implies

$$\max_j \|X_{0,j}^2\|_{\psi_1} \leq \max_j (\|X_{0,j}\|_{\psi_2})^2 \leq \omega_0^2. \quad (\text{B.30})$$

For any random variable Z , we have $\mathbb{E}|Z| \leq \|Z\|_{\psi_1}$. This implies

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I} \\ |\mathcal{M}| \leq \kappa_n}} \mathbb{E} \max_{j \in \mathcal{M}} X_{0,j}^2 \leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \max_{j \in \mathcal{M}} X_{0,j}^2 \right\|_{\psi_1}.$$

By (B.30) and Lemma 2.2.2 in van der Vaart and Wellner (1996), we have

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \max_{j \in \mathcal{M}} X_{0,j}^2 \right\|_{\psi_1} \leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I} \\ |\mathcal{M}| \leq \kappa_n}} K_0 \log(1 + \kappa_n) \max_{j \in \mathcal{M}} \|X_{0,j}^2\|_{\psi_1} \leq K_0 \omega_0^2 \log(1 + \kappa_n).$$

for some constant $K_0 > 0$. Note that $\kappa_n = o(n)$, we have $\log(1 + \kappa_n) \leq \log(n)$ for sufficiently large n . This completes the proof.

B.5 Proof of Lemma A.2

We first prove (27). Note that

$$\begin{aligned} & \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\boldsymbol{\omega}}_{\mathcal{M},j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 = \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{-1} \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0} - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \boldsymbol{\Sigma}_{\mathcal{M},j_0}\|_2 \\ & \leq \underbrace{\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{-1} (\widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0} - \boldsymbol{\Sigma}_{\mathcal{M},j_0})\|_2}_{\eta_1} + \underbrace{\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^{-1} - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1}) \boldsymbol{\Sigma}_{\mathcal{M},j_0}\|_2}_{\eta_2}. \end{aligned}$$

Hence, it suffices to show that with probability tending to 1,

$$\eta_1 = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right) \quad \text{and} \quad \eta_2 = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right) + O\left(\frac{\kappa_n}{\sqrt{n}}\right).$$

Upper bound for η_1 : Since

$$\begin{aligned} & \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}(\widehat{\Sigma}_{\mathcal{M}, j_0} - \Sigma_{\mathcal{M}, j_0})\|_2 \\ & \leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}\|_2 \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, j_0} - \Sigma_{\mathcal{M}, j_0})\|_2, \end{aligned}$$

it suffices to show with probability tending to 1 that,

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}\|_2 = O(1), \quad (\text{B.31})$$

and

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, j_0} - \Sigma_{\mathcal{M}, j_0})\|_2 = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right). \quad (\text{B.32})$$

Note that $\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}$ is symmetric. To prove (B.31), it is equivalent to show that the eigenvalues of $\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}$ are uniformly bounded with probability tending to 1. Hence, it suffices to prove

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \lambda_{\min}(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}) > \frac{\bar{c}}{2}, \quad (\text{B.33})$$

with probability tending to 1.

Observe that

$$\begin{aligned} & \min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \lambda_{\min}(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}) = \inf_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \inf_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbf{a}^T \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} \mathbf{a} \\ & \geq \min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \min_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbf{a}^T \Sigma_{\mathcal{M}, \mathcal{M}} \mathbf{a} - \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \max_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \left| \mathbf{a}^T (\Sigma_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}) \mathbf{a} \right|. \end{aligned}$$

By Condition (A2), the first term on the second line is greater than or equal to \bar{c} . Since $\Sigma_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}$ is symmetric, the second term is equal to

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \left\| \Sigma_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} \right\|_2. \quad (\text{B.34})$$

For any symmetric matrix \mathbf{S} , we have $\|\mathbf{S}\|_2 \leq \sqrt{\|\mathbf{S}\|_1 \|\mathbf{S}\|_\infty} = \|\mathbf{S}\|_\infty$, where $\|\cdot\|_1$ and $\|\cdot\|_\infty$ stand for the ℓ_1 and ℓ_∞ induced matrix norms. Hence, (B.34) is upper bounded by

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \left\| \boldsymbol{\Sigma}_{\mathcal{M}, \mathcal{M}} - \widehat{\boldsymbol{\Sigma}}_{\mathcal{M}, \mathcal{M}} \right\|_\infty \leq \kappa_n \max_{j_1, j_2 \in [1, \dots, p]} |\widehat{\Sigma}_{j_1, j_2} - \Sigma_{j_1, j_2}|.$$

To summarize, we've shown

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \lambda_{\min} \left(\widehat{\boldsymbol{\Sigma}}_{\mathcal{M}, \mathcal{M}} \right) \geq \bar{c} - \kappa_n \max_{j_1, j_2 \in [1, \dots, p]} |\widehat{\Sigma}_{j_1, j_2} - \Sigma_{j_1, j_2}|. \quad (\text{B.35})$$

Recall that $\widehat{\Sigma}_{j_1, j_2} - \Sigma_{j_1, j_2} = \sum_i (X_{i, j_1} X_{i, j_2} - \mathbb{E} X_{0, j_1} X_{0, j_2})/n$. Combining (B.30) with Cauchy-Schwarz inequality, we have

$$\|X_{0, j_1} X_{0, j_2}\|_{\psi_1} \leq \frac{\|X_{0, j_1}^2 + X_{0, j_2}^2\|_{\psi_1}}{2} \leq \frac{\|X_{0, j_1}^2\|_{\psi_1}}{2} + \frac{\|X_{0, j_2}^2\|_{\psi_1}}{2} \leq \omega_0^2, \quad (\text{B.36})$$

for all $j_1, j_2 \in [1, \dots, p]$. By Jensen's inequality, we have

$$\mathbb{E} \exp \left(\frac{\mathbb{E} |X_{0, j_1} X_{0, j_2}|}{\omega_0^2} \right) \leq \mathbb{E} \exp \left(\frac{|X_{0, j_1} X_{0, j_2}|}{\omega_0^2} \right) \leq 2.$$

This implies $\|\mathbb{E} X_{0, j_0} X_{0, j_2}\|_{\psi_1} \leq \omega_0^2$, $\forall j_1, j_2$. Combining this together with (B.36) gives

$$\|X_{0, j_1} X_{0, j_2} - \mathbb{E} X_{0, j_0} X_{0, j_2}\|_{\psi_1} \leq \|X_{0, j_1} X_{0, j_2}\|_{\psi_1} + \|\mathbb{E} X_{0, j_0} X_{0, j_2}\|_{\psi_1} \leq 2\omega_0^2.$$

Therefore, it follows from Bernstein's inequality (Theorem 3.1, Klartag and Mendelson, 2005) that

$$\max_{1 \leq j_1, j_2 \leq p} \Pr \left(\left| \sum_i (X_{i, j_1} X_{i, j_2} - \Sigma_{j_1, j_2}) \right| \geq t \right) \leq 2 \exp \left(-c_1 \min \left(\frac{t^2}{4n\omega_0^2}, \frac{t}{2\omega_0} \right) \right), \quad (\text{B.37})$$

for any $t > 0$ and some constant $c_1 > 0$.

Take $t_0 = 3\sqrt{n \log p} \omega_0 / \sqrt{c_1}$. Since $\log p = o(n)$, we have for sufficiently large n ,

$$\frac{t_0^2}{4n\omega_0^2} = \frac{9 \log p}{4c_1} \ll \frac{3\sqrt{n \log p}}{2\sqrt{c_1}} = \frac{t_0}{2\omega_0}.$$

It follows from (B.37) that

$$\max_{j_1, j_2} \Pr \left(\left| \sum_i (X_{i, j_1} X_{i, j_2} - \Sigma_{j_1, j_2}) \right| \geq t_0 \right) \leq 2 \exp \left(-\frac{c_1 t_0^2}{4n\omega_0^2} \right) \leq 2 \exp \left(-\frac{9 \log p}{4} \right).$$

By Bonferroni's inequality, we have

$$\begin{aligned}
& \Pr \left(\max_{j_1, j_2 \in [1, \dots, p]} \left| \sum_i (X_{i, j_1} X_{i, j_2} - \Sigma_{j_1, j_2}) \right| \geq t_0 \right) \\
& \leq \sum_{j_1, j_2 \in [1, \dots, p]} \Pr \left(\left| \sum_i (X_{i, j_1} X_{i, j_2} - \Sigma_{j_1, j_2}) \right| \geq t_0 \right) \\
& \leq p^2 2 \exp \left(-\frac{9 \log p}{4} \right) = 2 \exp \left(-\frac{9 \log p}{4} + 2 \log p \right) = 2 \exp \left(-\frac{\log p}{4} \right) \rightarrow 0.
\end{aligned} \tag{B.38}$$

Under the given conditions, we have that $\kappa_n t_0 / n = O(\kappa_n \sqrt{\log p / n}) = o(1)$. In view of (B.35), we've shown (B.33) holds for sufficiently large n .

Moreover, under the event defined in (B.38), we have

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, j_0} - \Sigma_{\mathcal{M}, j_0})\|_2 \leq \sqrt{\kappa_n} \max_{j_1, j_2 \in [1, \dots, p]} \left| \widehat{\Sigma}_{j_1, j_2} - \Sigma_{j_1, j_2} \right| \leq \frac{\sqrt{\kappa_n} t_0}{n}.$$

This proves (B.32).

Upper bound for η_2 : Observe that

$$\begin{aligned}
& \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} - \Sigma_{\mathcal{M}, \mathcal{M}}^{-1}) \Sigma_{\mathcal{M}, j_0}\|_2 \\
& = \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} (\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \Sigma_{\mathcal{M}, \mathcal{M}}) \Sigma_{\mathcal{M}, \mathcal{M}}^{-1} \Sigma_{\mathcal{M}, j_0}\|_2 \\
& \leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}\|_2 \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \Sigma_{\mathcal{M}, \mathcal{M}}) \omega_{\mathcal{M}, j_0}\|_2.
\end{aligned}$$

By (B.31), it suffices to show

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \Sigma_{\mathcal{M}, \mathcal{M}}) \omega_{\mathcal{M}, j_0}\|_2 = O \left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} \right) + O \left(\frac{\kappa_n}{\sqrt{n}} \right), \tag{B.39}$$

with probability tending to 1.

LHS of (B.39) can be upper bounded by

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} |\mathbf{a}^T (\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \Sigma_{\mathcal{M}, \mathcal{M}}) \omega_{\mathcal{M}, j_0}|. \tag{B.40}$$

For any subset \mathcal{M} such that $j_0 \notin \mathcal{M}$, $|\mathcal{M}| \leq \kappa_n$, define the empirical process

$$T_{\mathcal{M}}(\mathbf{a}) = \frac{1}{n} \sum_{i=1}^n g_{\mathcal{M}}(\mathbf{X}_i, \mathbf{a}) = \frac{1}{n} \sum_{i=1}^n \mathbf{a}^T (\mathbf{X}_{i, \mathcal{M}} \mathbf{X}_{i, \mathcal{M}}^T - \Sigma_{\mathcal{M}, \mathcal{M}}) \omega_{\mathcal{M}, j_0}.$$

The envelope function of $|g|$ is bounded by

$$G_{\mathcal{M}}(\mathbf{X}_i) \triangleq \|\mathbf{X}_{i,\mathcal{M}}\|_2 |\mathbf{X}_{i,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2 + \|\boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}\|_2 \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|. \quad (\text{B.41})$$

Note that $\boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}$ is positive definite, we have

$$\|\boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}\|_2 = \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \mathbf{a}^T \boldsymbol{\Sigma} \mathbf{a} = \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2 \leq \sqrt{\mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^4} \leq \sqrt{c_0}, \quad (\text{B.42})$$

where the first inequality follows from Cauchy-Schwarz inequality and the last inequality is due to Condition (A3). Combing this together with (B.41) and (24), we have

$$G_{\mathcal{M}}(\mathbf{X}_i) \leq \|\mathbf{X}_{i,\mathcal{M}}\|_2 |\mathbf{X}_{i,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2 + c_0/\sqrt{\bar{c}}. \quad (\text{B.43})$$

Given \mathcal{M} , the class of functions $\{\mathbf{a}^T (x_{\mathcal{M}} x_{\mathcal{M}}^T - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}) \boldsymbol{\omega}_{\mathcal{M},j_0}\}$ has finite VC index $|\mathcal{M}| + 2 \leq \kappa_n + 2 \leq 3\kappa_n$, since $\kappa_n \geq 1$. Therefore, it follows from Lemma 2.14.1 in van der Vaart and Wellner (1996) that

$$\mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \leq K_0 \sqrt{\frac{3\kappa_n}{n}} \sqrt{\mathbb{E} G_{\mathcal{M}}^2(\mathbf{X}_0)}. \quad (\text{B.44})$$

By (B.43), we have

$$\begin{aligned} \mathbb{E} G_{\mathcal{M}}^2(\mathbf{X}_0) &\leq \mathbb{E} (\|\mathbf{X}_{0,\mathcal{M}}\|_2 |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2 + c_0/\sqrt{\bar{c}})^2 \\ &\leq 2c_0^2/\bar{c} + 2\mathbb{E} \|\mathbf{X}_{0,\mathcal{M}}\|_2^2 |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2^2 \leq 2c_0^2/\bar{c} + 2\sqrt{\mathbb{E} \|\mathbf{X}_{0,\mathcal{M}}\|_2^4} \sqrt{\mathbb{E} |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2^4}, \end{aligned} \quad (\text{B.45})$$

where the last two inequalities are due to Cauchy-Schwarz inequality.

Note that

$$\begin{aligned} \mathbb{E} \|\mathbf{X}_{0,\mathcal{M}}\|_2^4 &\leq \mathbb{E} |\mathcal{M}| \sum_{j \in \mathcal{M}} \mathbf{X}_{0,j}^4 \leq \kappa_n^2 \max_{j \in [1, \dots, p]} \mathbb{E} X_{0,j}^4 \leq \kappa_n^2 \sup_{\substack{\mathbf{a} \in \mathbb{R}^{\kappa_n} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^4, \\ \mathbb{E} |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2^4 &\leq \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2^4 \mathbb{E} \left| \mathbf{X}_{0,\mathcal{M}}^T \frac{\boldsymbol{\omega}_{\mathcal{M},j_0}}{\|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2} \right|_2^4 \leq \frac{c_0^2}{\bar{c}^2} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{\kappa_n} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^4, \end{aligned}$$

where the last inequality is due to (24). Under Condition (A3), we have that

$$\mathbb{E} \|\mathbf{X}_{0,\mathcal{M}}\|_2^4 \leq \kappa_n^2 c_0 \quad \text{and} \quad \mathbb{E} |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|_2^4 \leq \frac{c_0^3}{\bar{c}^2}. \quad (\text{B.46})$$

This together with (B.45) gives that

$$\mathbb{E} G_{\mathcal{M}}^2(\mathbf{X}_0) \leq \frac{2c_0^2}{\bar{c}} + 2\sqrt{\kappa_n^2 c_0} \sqrt{\frac{c_0^3}{\bar{c}^2}} = \frac{2c_0^2}{\bar{c}} + \frac{2\kappa_n c_0^2}{\bar{c}} \leq \frac{4\kappa_n c_0^2}{\bar{c}},$$

where the last inequality is due to that $\kappa_n \geq 1$. Hence, by (B.44),

$$\mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \leq K_0 \sqrt{\frac{3\kappa_n}{n}} \sqrt{\frac{4\kappa_n c_0^2}{\bar{c}}} \leq K_0 c_0 \kappa_n \sqrt{\frac{12}{n\bar{c}}}. \quad (\text{B.47})$$

Besides, the $\|\cdot\|_{\psi_1}$ Orlicz norm of G can be upper bounded by

$$\begin{aligned} \|G_{\mathcal{M}}(\mathbf{X}_0)\|_{\psi_1} &\leq \|c_0/\sqrt{\bar{c}}\|_{\psi_1} + \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \left\| \|\mathbf{X}_{0,\mathcal{M}}\|_2^2 \right\|_{\psi_1} \\ &\leq c_0/\sqrt{\bar{c}} + \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \left\| \sum_{j \in \mathcal{M}} X_{0,j}^2 \right\|_{\psi_1} \leq c_0/\sqrt{\bar{c}} + \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \kappa_n \max_j \|X_{0,j}^2\|_{\psi_1} \\ &\leq c_0/\sqrt{\bar{c}} + \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \kappa_n \omega_0^2 \leq c_0/\sqrt{\bar{c}} \kappa_n + \|\boldsymbol{\omega}_{\mathcal{M},j_0}\|_2 \kappa_n \omega_0^2 \leq \sqrt{\frac{c_0}{\bar{c}}} \kappa_n (\sqrt{c_0} + \omega_0^2), \end{aligned}$$

where the fourth inequality follows from (B.30), and the last inequality is due to (24).

Hence, it follows from Lemma 2.2.2 in van der Vaart and Wellner (1996) that

$$\left\| \max_{i \in [1, \dots, n]} |G_{\mathcal{M}}(\mathbf{X}_i)| \right\|_{\psi_1} \leq K_1 \log(1+n) \max_{i \in [1, \dots, n]} \|G_{\mathcal{M}}(\mathbf{X}_i)\|_{\psi_1} = K_1 \sqrt{\frac{c_0}{\bar{c}}} (\sqrt{c_0} + \omega_0^2) \kappa_n \log(1+n).$$

Let $c_2 = K_1 \sqrt{c_0/\bar{c}} (\sqrt{c_0} + \omega_0^2)$, we've shown $\|\max_{i \in [1, \dots, n]} |G_{\mathcal{M}}(\mathbf{X}_i)|\|_{\psi_1} \leq c_2 \kappa_n \log(1+n)$.

Here, the constant c_2 is independent of \mathcal{M} .

Moreover, it follows from Cauchy-Schwarz inequality that

$$\begin{aligned} \sigma_*^2 &\equiv \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} g_{\mathcal{M}}(\mathbf{X}_0, \mathbf{a})^2 \\ &\leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} |\mathbf{a}^T (\mathbf{X}_{0,\mathcal{M}} \mathbf{X}_{0,\mathcal{M}}^T - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}}) \boldsymbol{\omega}_{\mathcal{M},j_0}|^2 \\ &\leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}} \mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|^2 \\ &\leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \sqrt{\mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^4} \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sqrt{\mathbb{E} |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4} \leq \frac{c_0^2}{\bar{c}}, \end{aligned}$$

where the last inequality follows by (B.46) and Condition (A3).

Therefore, it follows from Theorem 4 in Adamczak (2008) that there exists some constant $K_2 > 0$ such that

$$\begin{aligned}
& \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \Pr \left(\sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| - \frac{3}{2} \mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \geq \frac{t}{n} \right) \\
& \leq \exp \left(-\frac{t^2}{3n\sigma_*^2} \right) + 3 \exp \left(-\frac{t}{K_2 c_2 \kappa_n \log(1+n)} \right) \\
& \leq \exp \left(-\frac{t^2 \bar{c}}{3n c_0^2} \right) + 3 \exp \left(-\frac{t}{K_2 c_2 \kappa_n \log(1+n)} \right), \quad \forall t > 0.
\end{aligned}$$

Define

$$t_0 = \max \left(\frac{2c_0 \sqrt{n \kappa_n \log p}}{\sqrt{\bar{c}}}, \frac{4}{3} K_2 c_2 \kappa_n^2 \log p \log(n+1) \right),$$

we have

$$\begin{aligned}
& \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \Pr \left(\sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| - \frac{3}{2} \mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \geq \frac{t_0}{n} \right) \\
& \leq \exp \left(-\frac{4\bar{c}c_0^2 n \kappa_n \log p}{3n \bar{c} c_0^2} \right) + 3 \exp \left(-\frac{4K_2 c_2 \kappa_n^2 \log p \log(n+1)}{3K_2 c_2 \kappa_n \log(1+n)} \right) \leq 4 \exp \left(-\frac{4}{3} \kappa_n \log p \right).
\end{aligned}$$

The number of subset \mathcal{M} with less than or equal to κ_n elements is upper bounded by $C_p^{\kappa_n} \leq p^{\kappa_n}$. Hence, it follows from Bonferroni's inequality that

$$\begin{aligned}
& \Pr \left(\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| - \frac{3}{2} \mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \geq \frac{t_0}{n} \right) \\
& \leq p^{\kappa_n} \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \Pr \left(\sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| - \frac{3}{2} \mathbb{E} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \geq \frac{t_0}{n} \right) \\
& \leq 4p^{\kappa_n} \exp \left(-\frac{4}{3} \kappa_n \log p \right) = 4 \exp \left(-\frac{4}{3} \kappa_n \log p + \kappa_n \log p \right) = 4 \exp \left(-\frac{1}{3} \kappa_n \log p \right) \rightarrow 0.
\end{aligned}$$

This together with (B.44) implies that

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} |T_{\mathcal{M}}(\mathbf{a})| \leq K_0 c_0 \kappa_n \sqrt{\frac{27}{n \bar{c}}} + \frac{t_0}{n}, \quad (\text{B.48})$$

with probability tending to 1.

Under the given conditions, we have that for sufficiently large n ,

$$\frac{2c_0\sqrt{n\kappa_n\log p}}{\sqrt{\bar{c}}} \gg \frac{4}{3}K_2c_2\kappa_n^2\log p\log(n+1),$$

and hence $t_0 = 2c_0\sqrt{n\kappa_n\log p}/\sqrt{\bar{c}}$. Under the event defined in (B.48), we have for sufficiently large n ,

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} |T_{\mathcal{M}}(\mathbf{a})| = O\left(\frac{\sqrt{\kappa_n\log p}}{\sqrt{n}}\right) + O\left(\frac{\kappa_n}{\sqrt{n}}\right).$$

This proves (B.39). The upper bound for η_2 is thus given.

Consider (28). Assume for now, we've shown

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2| = O\left(\frac{\sqrt{\kappa_n\log p}}{\sqrt{n}}\right) + O\left(\frac{\kappa_n}{\sqrt{n}}\right), \quad (\text{B.49})$$

with probability tending to 1. Then, under the event defined in (B.49), we have

$$\begin{aligned} \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M}, j_0} - \sigma_{\mathcal{M}, j_0}| &= \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{|\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2|}{|\hat{\sigma}_{\mathcal{M}, j_0} + \sigma_{\mathcal{M}, j_0}|} \\ &\leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{|\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2|}{|\sigma_{\mathcal{M}, j_0}|} \leq \frac{1}{\sqrt{\bar{c}}} \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2| = O\left(\frac{\sqrt{\kappa_n\log p}}{\sqrt{n}}\right) + O\left(\frac{\kappa_n}{\sqrt{n}}\right), \end{aligned}$$

where the last inequality follows from (24) and the last equality is due to (B.49). Hence, it suffices to show (B.49).

By definition, we have

$$\begin{aligned} |\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2| &\leq |\hat{\Sigma}_{j_0, j_0} - \Sigma_{j_0, j_0}| + |\hat{\Sigma}_{\mathcal{M}, j_0}^T \hat{\omega}_{\mathcal{M}, j_0} - \Sigma_{\mathcal{M}, j_0}^T \omega_{\mathcal{M}, j_0}| \quad (\text{B.50}) \\ &\leq |\hat{\Sigma}_{j_0, j_0} - \Sigma_{j_0, j_0}| + |\Sigma_{\mathcal{M}, j_0}^T (\hat{\omega}_{\mathcal{M}, j_0} - \omega_{\mathcal{M}, j_0})| + |(\Sigma_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0})^T \omega_{\mathcal{M}, j_0}| \\ &\quad + |(\Sigma_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0})^T (\hat{\omega}_{\mathcal{M}, j_0} - \omega_{\mathcal{M}, j_0})| \leq |\hat{\Sigma}_{j_0, j_0} - \Sigma_{j_0, j_0}| \\ &\quad + \|\Sigma_{\mathcal{M}, j_0}\|_2 \|\hat{\omega}_{\mathcal{M}, j_0} - \omega_{\mathcal{M}, j_0}\|_2 + \|\Sigma_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}\|_2 \|\omega_{\mathcal{M}, j_0}\|_2 \\ &\quad + \|\hat{\omega}_{\mathcal{M}, j_0} - \omega_{\mathcal{M}, j_0}\|_2 \|\Sigma_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}\|_2. \end{aligned}$$

It follows from (24), (27), (B.32) and (B.38) that with probability tending to 1,

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\Sigma_{\mathcal{M}, j_0} - \widehat{\Sigma}_{\mathcal{M}, j_0}\|_2 = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right), \quad (\text{B.51})$$

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\omega_{\mathcal{M}, j_0} - \widehat{\omega}_{\mathcal{M}, j_0}\|_2 = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right) + O\left(\frac{\kappa_n}{\sqrt{n}}\right), \quad (\text{B.52})$$

$$|\widehat{\Sigma}_{j_0, j_0} - \Sigma_{j_0, j_0}| = O\left(\frac{\sqrt{\log p}}{\sqrt{n}}\right), \quad \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\omega_{\mathcal{M}, j_0}\|_2 = O(1). \quad (\text{B.53})$$

By Condition (A2), $\Sigma_{\mathcal{M}, \mathcal{M}}$ is invertible for any subset \mathcal{M} such that $|\mathcal{M}| \leq \kappa_n$. Hence, it follows from (B.42) that

$$\min_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \lambda_{\min}(\Sigma_{\mathcal{M}, \mathcal{M}}^{-1}) \geq c_0^{-1/2}.$$

Using similar arguments in proving (24), we can show that

$$\max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \|\Sigma_{\mathcal{M}, j_0}\|_2^2 \leq \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} \frac{c_0}{\lambda_{\min}(\Sigma_{\mathcal{M}, \mathcal{M}}^{-1})} \leq c_0^{3/2}.$$

Under the given conditions, we have $\kappa_n \log p = o(n)$. Under the events defined in (B.50)-(B.52), we obtain that

$$\begin{aligned} \max_{\substack{\mathcal{M} \subseteq [1, \dots, p] \\ j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n}} |\hat{\sigma}_{\mathcal{M}, j_0}^2 - \sigma_{\mathcal{M}, j_0}^2| &\leq O\left(\frac{\sqrt{\log p}}{\sqrt{n}}\right) + c_0^{3/2} O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n}{\sqrt{n}}\right) \\ &+ O(1) O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right) + O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}}\right) O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n}{\sqrt{n}}\right) = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n}{\sqrt{n}}\right). \end{aligned}$$

This proves (B.49). The proof is hence completed.

B.6 Proof of Lemma A.3

We first prove

$$\max_{i \in [1, \dots, n]} \max_{j \in [1, \dots, p]} |X_{i,j}| \leq \sqrt{2\omega_0^2 \log p}. \quad (\text{B.54})$$

Note that

$$\max_{i,j} \Pr\left(|X_{i,j}| > \sqrt{2\omega_0^2 \log p}\right) \leq \frac{\mathbb{E} \exp(|X_{i,j}|^2/\omega_0^2)}{\exp(2\omega_0^2 \log p/\omega_0^2)} \leq \frac{2}{p^2},$$

where the first inequality follows from Markov's inequality and the second inequality is due to the definition of the Orlicz norm. Since $p \gg n$, it follows from Bonferroni's inequality that

$$\Pr \left(\max_{i,j} |X_{i,j}| > \sqrt{2\omega_0^2 \log p} \right) \leq np \max_{i,j} \Pr \left(|X_{i,j}| > \sqrt{2\omega_0^2 \log p} \right) \leq \frac{2n}{p} \rightarrow 0.$$

This proves (B.54).

Under Condition (A1), we have for any $t \in [s_n, \dots, n-1]$,

$$\begin{aligned} & \mathbb{E} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^2 X_{t+1,j}^2 | \mathcal{F}_t \right\} \\ & \leq \sqrt{\mathbb{E} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^4 | \mathcal{F}_t \right\}} \sqrt{\mathbb{E} (X_{t+1,j}^4 | \mathcal{F}_t)} \\ & \leq \max_{j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n} \sqrt{\mathbb{E} (X_{0,j_0} - \boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}})^4} \max_{j \in [1, \dots, p]} \sqrt{\mathbb{E} X_{0,j}^4} \\ & \leq \max_{j_0 \notin \mathcal{M}, |\mathcal{M}| \leq \kappa_n} \sqrt{8\mathbb{E} X_{0,j_0}^4 + 8\mathbb{E} |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4} \max_{j \in [1, \dots, p]} \sqrt{\mathbb{E} X_{0,j}^4} \end{aligned}$$

where the first inequality follows from Cauchy-Schwarz inequality, and the last inequality is due to that the elementary inequality that $(a+b)^4 \leq 8a^4 + 8b^4, \forall a, b \in \mathbb{R}$.

Under (A3), we have $\max_j \mathbb{E} X_{0,j}^4 \leq c_0, \mathbb{E} X_{0,j_0}^4 \leq c_0$. By (B.46), we have $\mathbb{E} |\mathbf{X}_{0,\mathcal{M}}^T \boldsymbol{\omega}_{\mathcal{M},j_0}|^4 \leq c_0^3/\bar{c}^2$. Thus,

$$\mathbb{E} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^2 X_{t+1,j}^2 | \mathcal{F}_t \right\} \leq \sqrt{8} \sqrt{1 + \frac{c_0^2}{\bar{c}^2}} c_0.$$

Combining this together with (24), we have almost surely,

$$\mathbb{E} \frac{1}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^2} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^2 X_{t+1,j}^2 | \mathcal{F}_t \right\} \leq \sqrt{8 + \frac{8c_0^2}{\bar{c}^2}} \frac{c_0}{\bar{c}}. \quad (\text{B.55})$$

Let

$$I_{2,j} = \frac{1}{\sqrt{n}} \sum_{t=s_n}^{n-1} \frac{1}{\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) X_{t+1,j} I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}).$$

Define $Z_{t+1,j}^* = \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right) X_{t+1,j}$ and

$$Z_{t+1,j}^{**} = Z_{t+1,j}^* I \left\{ |Z_{t+1,j}^*| \leq 2\omega_0^2 \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \log p \right\}.$$

Under the events $\max_{i,j} |X_{i,j}| \leq \sqrt{2\omega_0^2 \log p}$ and $|\widehat{\mathcal{M}}_{j_0}^{(t)}| \leq \kappa_n$, it follows from (24) that

$$\begin{aligned} |Z_{t+1,j}^*| &\leq \left| X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right| |X_{t+1,j}| \\ &\leq \left(|X_{t+1,j_0}| + \|\boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}\|_2 \sqrt{\kappa_n} \max_j |X_{t+1,j}| \right) |X_{t+1,j}| \\ &\leq \left\{ \sqrt{2\omega_0^2 \log p} \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \right\} \sqrt{2\omega_0^2 \log p} = 2\omega_0^2 \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \log p, \end{aligned}$$

and hence $Z_{t+1,j}^{**} = Z_{t+1,j}^*$.

By (B.54) and Condition (A1), we have with probability tending to 1,

$$I_{2,j} = \sum_{t=s_n}^{n-1} \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} Z_{t+1,j}^{**} I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}).$$

Therefore, it suffices to show with probability tending to 1,

$$\max_j \left| \sum_{t=s_n}^{n-1} \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} Z_{t+1,j}^{**} I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}) \right| \leq \bar{c}_* \log p,$$

for some constant $\bar{c}_* > 0$. Let

$$I_{2,j}^* = \sum_{t=s_n}^{n-1} \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E}(Z_{t+1,j}^{**} | \mathcal{F}_t) I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}).$$

We've shown in the proof of Theorem 2.1 that $\mathbb{E}\{Z_{t+1,j}^* I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}) | \mathcal{F}_t\} = 0$. Therefore, we have

$$\begin{aligned} &\left| \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E}(Z_{t+1,j}^{**} | \mathcal{F}_t) I(j \in \widehat{\mathcal{M}}_{j_0}^{(t)}) \right| \tag{B.56} \\ &\leq \left| \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E} \left[Z_{t+1,j}^* I \left\{ |Z_{t+1,j}^*| > 2\omega_0^2 \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \log p \right\} | \mathcal{F}_t \right] \right|. \end{aligned}$$

It follows from Cauchy-Schwarz inequality that

$$\begin{aligned} &\left| \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E} \left[Z_{t+1,j}^* I \left\{ |Z_{t+1,j}^*| > 2\omega_0^2 \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \log p \right\} | \mathcal{F}_t \right] \right| \\ &\leq \left| \frac{1}{\sqrt{n} \sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \sqrt{\mathbb{E}\{(Z_{t+1,j}^*)^2 | \mathcal{F}_t\}} \right| \left| \Pr \left\{ |Z_{t+1,j}^*| > 2\omega_0^2 \left(1 + \sqrt{\frac{c_0 \kappa_n}{\bar{c}}} \right) \log p \right\} \right| \end{aligned}$$

By (B.55), the first term on the second line is upper bounded by $(8 + 8c_0^2/\bar{c}^2)^{1/4}(c_0/\bar{c})^{1/2}/\sqrt{n}$ with probability 1. As for the second term, under the event $|\widehat{\mathcal{M}}_{j_0}^{(t)}| \leq \kappa_n$, using similar arguments in the proof of Lemma A.2, we can show that

$$\Pr \left\{ |Z_{t+1,j}| > 2\omega_0^2 \left(1 + \sqrt{\frac{c_0\kappa_n}{\bar{c}}} \right) \log p \right\} \leq \Pr \left(\max_j |X_{t+1,j}| > \sqrt{2\omega_0 \log p} \right) \leq \frac{2}{p}.$$

Hence, we have

$$\left| \frac{1}{\sqrt{n}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E} \left[Z_{t+1,j} I \left\{ |Z_{t+1,j}| > 2\omega_0^2 \left(1 + \sqrt{\frac{c_0\kappa_n}{\bar{c}}} \right) \log p \right\} | \mathcal{F}_t \right] \right| \leq \sqrt{\frac{2c_0}{\bar{c}np}} (8 + 8c_0^2/\bar{c}^2)^{1/4}.$$

Therefore, it follows from (B.56) that

$$\left| \frac{1}{\sqrt{n}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \mathbb{E}(Z_{t+1,j}^* | \mathcal{F}_t) I \left(j \in \widehat{\mathcal{M}}_{j_0}^{(t)} \right) \right| \leq \sqrt{\frac{2c_0}{\bar{c}np}} (8 + 8c_0^2/\bar{c}^2)^{1/4}.$$

Since $p \gg n$, we have

$$|I_{2,j}^*| \leq \sqrt{\frac{2c_0n}{\bar{c}p}} (8 + 8c_0^2/\bar{c}^2)^{1/4} = o(1),$$

where the $o(1)$ term is uniform in j .

Hence, it suffices to show $\Pr(\max_j |I_{2,j}^{**}| \leq \bar{c}_* \log p) \rightarrow 1$, for some constant $\bar{c}_* > 0$, where

$$I_{2,j}^{**} = \sum_{t=s_n}^{n-1} \frac{1}{\sqrt{n}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \{Z_{t+1,j}^* - \mathbb{E}(Z_{t+1,j}^* | \mathcal{F}_t)\} I \left(j \in \widehat{\mathcal{M}}_{j_0}^{(t)} \right).$$

For any $j \in [1, \dots, p]$, $I_{2,j}^{**}$ is a mean zero martingale with respect to the filtration $\{\sigma(\mathcal{F}_t)\}$.

Besides, given \mathcal{F}_t , by the definition of $Z_{t+1,j}^*$ and (24), we have that

$$\begin{aligned} & \left| \frac{1}{\sqrt{n}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \{Z_{t+1,j}^* - \mathbb{E}(Z_{t+1,j}^* | \mathcal{F}_t)\} I \left(j \in \widehat{\mathcal{M}}_{j_0}^{(t)} \right) \right| \\ & \leq \left| \frac{1}{\sqrt{n}\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}} \{Z_{t+1,j}^* - \mathbb{E}(Z_{t+1,j}^* | \mathcal{F}_t)\} \right| \leq \frac{4\omega_0^2}{\sqrt{n\bar{c}}} \left(1 + \sqrt{\frac{c_0\kappa_n}{\bar{c}}} \right) \log p, \quad a.s. \end{aligned}$$

Moreover, it follows from (B.55) that we have, almost surely,

$$\sum_{t=s_n}^{n-1} \mathbb{E} \frac{1}{n\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^2} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^2 X_{t+1,j}^2 | \mathcal{F}_t \right\} \leq \sqrt{8 + \frac{8c_0^2}{\bar{c}^2}} \frac{c_0}{\bar{c}}. \quad (\text{B.57})$$

Set $y_0 = \sqrt{8(1 + c_0^2/\bar{c}^2)}c_0/\bar{c}$ and

$$c_0^* = \frac{4\omega_0^2}{\sqrt{n\bar{c}}} \left(1 + \sqrt{\frac{c_0\kappa_n}{\bar{c}}} \right) \log p.$$

It follows from Theorem 9.12 in de la Peña et al. (2009) that

$$\begin{aligned} \Pr \left(|I_{2,j}^{**}| > z, \sum_{t=s_n}^{n-1} \mathbb{E} \frac{1}{n\sigma_{\widehat{\mathcal{M}}_{j_0}^{(t)}}^2} \left\{ \left(X_{t+1,j_0} - \boldsymbol{\omega}_{\widehat{\mathcal{M}}_{j_0}^{(t)}}^T \mathbf{X}_{t+1,\widehat{\mathcal{M}}_{j_0}^{(t)}} \right)^2 X_{t+1,j}^2 | \mathcal{F}_t \right\} \leq y_0 \right) \\ \leq 2 \exp \left(-\frac{z^2}{2(y_0 + c_0^*z)} \right), \quad \forall z > 0. \end{aligned}$$

In view of (B.57), we have

$$\Pr (|I_{2,j}^{**}| > z) \leq 2 \exp \left(-\frac{z^2}{2(y_0 + c_0^*z)} \right) \leq 2 \exp \left(-\min \left(\frac{z^2}{4y_0}, \frac{z}{2c_0^*} \right) \right).$$

Let $z_0 = 3 \max(\sqrt{y_0} \log p, c_0^* \log p)$, we have

$$\Pr (|I_{2,j}^{**}| > z_0) \leq 2 \exp \left(-\min \left(\frac{9}{4} \log^2 p, \frac{3}{2} \log p \right) \right) \leq 2 \exp \left(-\frac{3}{2} \log p \right).$$

It follows from Bonferroni's inequality that

$$\Pr \left(\max_j |I_{2,j}^{**}| > z_0 \right) \leq \sum_j \Pr (|I_{2,j}^{**}| > z_0) \leq 2p \exp \left(-\frac{3}{2} \log p \right) = 2 \exp \left(-\frac{1}{2} \log p \right) \rightarrow 0.$$

Under the given conditions, we have $\kappa_n \log^2 p = o(n)$ and hence $z_0 = 3\sqrt{y_0} \log p$ for sufficiently large n . By taking $\bar{c}_* = 3\sqrt{y_0}$, this shows $\Pr(\max_j |I_{2,j}^{**}| \leq \bar{c}_* \log p) \rightarrow 1$. The proof is hence completed.

B.7 Proof of Lemma B.1

Assertion (B.1) can be proven in a similar manner as (24). We omit its proof for brevity. To prove (B.2) and (B.3), we first show the following events occur with probability tending to 1,

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}} - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}} \right\|_2 \leq \frac{\bar{c}_* \kappa_n \sqrt{\log p}}{\sqrt{n}} + \bar{c}_* \eta_n, \quad (\text{B.58})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},j_0} - \boldsymbol{\Sigma}_{\mathcal{M},j_0} \right\|_2 \leq \frac{\bar{c}_* \sqrt{\kappa_n \log p}}{\sqrt{n}} + \bar{c}_* \eta_n, \quad (\text{B.59})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \left(\widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}} - \boldsymbol{\Sigma}_{\mathcal{M},\mathcal{M}} \right) \boldsymbol{\omega}_{\mathcal{M},j_0} \right\|_2 \leq \frac{\bar{c}_* (\sqrt{\kappa_n \log p} + \kappa_n)}{\sqrt{n}} + \bar{c}_* \eta_n. \quad (\text{B.60})$$

Using similar arguments in the proof of Lemma A.2, we can show that there exists some constant $\bar{c}_{**} > 0$ such that the following events occur with probability tending to 1,

$$\begin{aligned} \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \Sigma_{\mathcal{M}, \mathcal{M}} \right\|_2 &\leq \frac{\bar{c}_{**} \kappa_n \sqrt{\log p}}{\sqrt{n}}, \\ \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \hat{\Sigma}_{\mathcal{M}, j_0}^* - \Sigma_{\mathcal{M}, j_0} \right\|_2 &\leq \frac{\bar{c}_{**} \sqrt{\kappa_n \log p}}{\sqrt{n}}, \\ \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \left(\hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \Sigma_{\mathcal{M}, \mathcal{M}} \right) \omega_{\mathcal{M}, j_0} \right\|_2 &\leq \frac{\bar{c}_{**} \sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\bar{c}_{**} \kappa_n}{\sqrt{n}}. \end{aligned} \quad (\text{B.61})$$

Therefore, it suffices to show the following events occur with probability tending to 1,

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \left(\hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \hat{\Sigma}_{\mathcal{M}, \mathcal{M}} \right) \omega_{\mathcal{M}, j_0} \right\|_2 \leq \bar{c}_{***} \eta_n, \quad (\text{B.62})$$

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \hat{\Sigma}_{\mathcal{M}, \mathcal{M}} \right\|_2 \leq \bar{c}_{***} \eta_n, \quad (\text{B.63})$$

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \hat{\Sigma}_{\mathcal{M}, j_0}^* - \hat{\Sigma}_{\mathcal{M}, j_0} \right\|_2 \leq \bar{c}_{***} \eta_n. \quad (\text{B.64})$$

Using similar arguments in (B.15), we can show that

$$\max_{i \in [1, \dots, n]} |b''(\mathbf{X}_i^T \tilde{\beta}) - b''(\mathbf{X}_i^T \beta_0)| \leq \bar{c}_{****} \omega_0 \eta_n,$$

for some constant $\bar{c}_{****} > 0$. By Condition (A4*), we have

$$\begin{aligned} &\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \left(\hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \hat{\Sigma}_{\mathcal{M}, \mathcal{M}} \right) \omega_{\mathcal{M}, j_0} \right\|_2 \\ &\leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 \leq 1}} \left| \mathbf{a}^T \left(\hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^* - \hat{\Sigma}_{\mathcal{M}, \mathcal{M}} \right) \omega_{\mathcal{M}, j_0} \right| \\ &\leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 \leq 1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i, \mathcal{M}}| |\omega_{\mathcal{M}, j_0}^T \mathbf{X}_{i, \mathcal{M}}| |b''(\mathbf{X}_i^T \tilde{\beta}) - b''(\mathbf{X}_i^T \beta_0)| \\ &\leq \bar{c}_{****} \omega_0 \eta_n \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 \leq 1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i, \mathcal{M}}| |\omega_{\mathcal{M}, j_0}^T \mathbf{X}_{i, \mathcal{M}}|. \end{aligned} \quad (\text{B.65})$$

Note that

$$\begin{aligned}
& \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}| |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{i,\mathcal{M}}| \\
& \leq \underbrace{\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \frac{1}{n} \sum_{i=1}^n (|\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}| |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{i,\mathcal{M}}| - \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}| |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|)}_{\eta_1} \\
& + \underbrace{\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}| |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|}_{\eta_2}.
\end{aligned}$$

Using similar arguments in bounding (B.40), we can show that

$$|\eta_1| = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n}{\sqrt{n}}\right), \quad (\text{B.66})$$

with probability tending to 1. Under the given conditions, the RHS of (B.66) is $o(1)$.

Besides, by Condition (A3*), (B.1) and Cauchy-Schwarz inequality, we have

$$\begin{aligned}
|\eta_2| & \leq \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} (\mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2)^{1/2} (\mathbb{E} |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^2)^{1/2} \\
& \leq \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} (\mathbb{E} |\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^4)^{1/4} (\mathbb{E} |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}|^4)^{1/4} \leq \frac{c_0}{\sqrt{c}}.
\end{aligned}$$

Combining this together with (B.66) gives

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}| |\boldsymbol{\omega}_{\mathcal{M},j_0}^T \mathbf{X}_{0,\mathcal{M}}| \leq \frac{2c_0}{\sqrt{c}}, \quad (\text{B.67})$$

with probability tending to 1. Assertion (B.62) thus follows by combining (B.65) together with (B.67).

As for (B.63), we have with probability tending to 1,

$$\begin{aligned}
\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}}^* - \widehat{\boldsymbol{\Sigma}}_{\mathcal{M},\mathcal{M}} \right\|_2 & \leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 \leq 1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2 |b''(\mathbf{X}_i^T \widetilde{\boldsymbol{\beta}}) - b''(\mathbf{X}_i^T \boldsymbol{\beta}_0)| \\
& \leq \bar{c}_{****} \omega_0 \eta_n \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 \leq 1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2. \quad (\text{B.68})
\end{aligned}$$

Besides,

$$\begin{aligned}
& \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2 \\
& \leq \underbrace{\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \frac{1}{n} \sum_{i=1}^n (|\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2 - \mathbb{E}|\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2)}_{\eta_1^*} + \underbrace{\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \mathbb{E}|\mathbf{a}^T \mathbf{X}_{0,\mathcal{M}}|^2}_{\eta_2^*}.
\end{aligned}$$

Using similar arguments in bounding (B.40), we have with probability tending to 1,

$$|\eta_1^*| = O\left(\frac{\sqrt{\kappa_n \log p}}{\sqrt{n}} + \frac{\kappa_n^{3/2}}{\sqrt{n}}\right).$$

Moreover, by (A3*) and Cauchy-Schwarz inequality, we have $\eta_2^* \leq c_0^{1/2}$. Combining these together with (B.68), we obtain

$$\Pr\left(\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}} \right\|_2 \leq 2c_0^{1/2} \bar{c}_{****} \omega_0 \eta_n\right) \rightarrow 1.$$

This proves (B.63). Similarly, we can show (B.64) holds. This proves (B.58)-(B.60). Based on these results, following the arguments in the proof of Lemma A.2, we can show (B.2) and (B.3) hold. Besides, based on (B.62)-(B.64), we can similarly show (B.5) holds.

Now, we focus on proving (B.4). We first show

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{*-1} - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \right\|_2 \leq \bar{c}_0^* \eta_n, \quad (\text{B.69})$$

for some constant $\bar{c}_0^* > 0$, with probability tending to 1. Note that

$$\begin{aligned}
& \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{*-1} - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \right\|_2 = \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{*-1} (\widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}) \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \right\|_2 \\
& \leq \max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{*-1} \right\|_2 \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^{-1} \right\|_2 \left\| \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}} \right\|_2
\end{aligned}$$

By the condition $\kappa_n^3 \log p = o(n)$, (A2*), (A5*), (B.58) and (B.61), we can show the following events occur with probability tending to 1,

$$\min_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \lambda_{\min}(\widehat{\Sigma}_{\mathcal{M},\mathcal{M}}) \geq \frac{\bar{c}}{2}, \quad \min_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \lambda_{\min}(\widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^*) \geq \frac{\bar{c}}{2}, \quad (\text{B.70})$$

for sufficiently large n . Hence, we have

$$\Pr \left(\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1} \right\|_2 \left\| \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} \right\|_2 \leq \frac{4}{\bar{c}^2} \right) \rightarrow 1.$$

Combining this together with (B.63) yields

$$\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} \right\|_2 \leq \frac{4\bar{c}_{***}\eta_n}{\bar{c}^2}.$$

This proves (B.69).

For any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$, we have

$$\begin{aligned} \widehat{\omega}_{\mathcal{M}, j_0} - \widehat{\omega}_{\mathcal{M}, j_0}^* &= \underbrace{\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1}(\widehat{\Sigma}_{\mathcal{M}, j_0} - \widehat{\Sigma}_{\mathcal{M}, j_0}^*)}_{I_1^*} + (\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1})\widehat{\Sigma}_{\mathcal{M}, j_0}^* \\ &+ \underbrace{(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1})(\widehat{\Sigma}_{\mathcal{M}, j_0} - \widehat{\Sigma}_{\mathcal{M}, j_0}^*)}_{I_2^*} = I_1^* + \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1}(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^*)\widehat{\omega}_{\mathcal{M}, j_0}^* \\ &+ I_2^* = I_1^* + I_2^* + \underbrace{(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{-1} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1})(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^*)\widehat{\omega}_{\mathcal{M}, j_0}^*}_{I_3^*} + \underbrace{\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1}(\widehat{\Sigma}_{\mathcal{M}, \mathcal{M}} - \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^*)\widehat{\omega}_{\mathcal{M}, j_0}^*}_{I_4^*} \end{aligned}$$

By (B.64) and (B.69), it is immediate to see that $|I_2^*|$ is upper bounded by $\bar{c}_0^*\bar{c}_{***}\eta_n^2$, with probability tending to 1. Besides, similar to (B.14), we can show

$$\Pr \left(\max_{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n} \left\| \widehat{\omega}_{\mathcal{M}, j_0}^* \right\|_2 \leq 2\sqrt{c_0/\bar{c}} \right) \rightarrow 1. \quad (\text{B.71})$$

This together with (B.63) and (B.69) yields that

$$\Pr \left(|I_3^*| \leq \frac{4\bar{c}_{***}\eta_n^2}{\bar{c}^2} 2\sqrt{c_0/\bar{c}} \right) \rightarrow 1.$$

Recall that

$$\widetilde{\omega}_{\mathcal{M}, j_0} = \widehat{\omega}_{\mathcal{M}, j_0}^* + \sum_{j=1}^p \widehat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1} \left(\widehat{\Psi}_{\mathcal{M}, j_0}^{(j)} + \widehat{\Psi}_{\mathcal{M}, \mathcal{M}}^{(j)} \widehat{\omega}_{\mathcal{M}, j_0}^* \right) (\widetilde{\beta}_j - \beta_{0,j}).$$

Hence, in order to prove

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \widehat{\omega}_{\mathcal{M}, j_0} - \widetilde{\omega}_{\mathcal{M}, j_0} \right\|_2 \leq \bar{c}_0\eta_n^2, \quad (\text{B.72})$$

it suffices to show the following events occur with probability tending to 1,

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| I_1^* - \sum_{j=1}^p \hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1} \hat{\Psi}_{\mathcal{M}, j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j}) \right\|_2 \leq \bar{c}_0 \eta_n^2, \quad (\text{B.73})$$

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| I_4^* - \sum_{j=1}^p \hat{\Sigma}_{\mathcal{M}, \mathcal{M}}^{*-1} \hat{\Psi}_{\mathcal{M}, \mathcal{M}}^{(j)} \hat{\omega}_{\mathcal{M}, j_0}^* (\tilde{\beta}_j - \beta_{0,j}) \right\|_2 \leq \bar{c}_0 \eta_n^2. \quad (\text{B.74})$$

We first prove (B.73). By (B.70) and the definition of $\hat{\Psi}_{\mathcal{M}, j_0}^{(j)}$, it suffices to show

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \hat{\Sigma}_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}^* - \frac{1}{n} \sum_{i=1}^n \mathbf{X}_{i, \mathcal{M}} b'''(\mathbf{X}_i^T \beta_0) X_{i, j_0} \{ \mathbf{X}_i^T (\tilde{\beta} - \beta_0) \} \right\|_2 \leq c_1^* \eta_n^2,$$

for some constant $c_1^* > 0$, with probability tending to 1. This is equivalent to show

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \left| \mathbf{a}^T \left(\hat{\Sigma}_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}^* - \frac{1}{n} \sum_{i=1}^n \mathbf{X}_{i, \mathcal{M}} b'''(\mathbf{X}_i^T \beta_0) X_{i, j_0} \{ \mathbf{X}_i^T (\tilde{\beta} - \beta_0) \} \right) \right| \leq c_1^* \eta_n^2,$$

with probability tending to 1.

For any $\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}$, it follows from Taylor's theorem that

$$\mathbf{a}^T \left(\hat{\Sigma}_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}^* \right) = \frac{1}{n} \sum_{i=1}^n \mathbf{a}^T \mathbf{X}_{i, \mathcal{M}} b'''(\mathbf{X}_i^T \beta_a^*) X_{i, j_0} \{ \mathbf{X}_i^T (\tilde{\beta} - \beta_0) \},$$

for some β_a^* lying on the line segment joining β_0 and $\tilde{\beta}$. By (A5*), we have

$$\Pr \left(\sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \|\beta_0 - \beta_a^*\|_1 \leq \eta_n \right) \rightarrow 1. \quad (\text{B.75})$$

The function b''' is Lipschitz continuous. By (A4*), under the event defined in (B.75), we have

$$\sup_{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}} \max_{i=1, \dots, n} |b'''(\mathbf{X}_i^T \beta_a^*) - b'''(\mathbf{X}_i^T \beta_0)| \leq L_0 \max_{i=1, \dots, n} \|\mathbf{X}_i\|_\infty \|\beta_0 - \beta_a^*\|_1 \leq L_0 \omega_0 \eta_n, \quad (\text{B.76})$$

for some constant $L_0 > 0$. Therefore, we have

$$\begin{aligned} & \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \left| \mathbf{a}^T \left(\hat{\Sigma}_{\mathcal{M}, j_0} - \hat{\Sigma}_{\mathcal{M}, j_0}^* - \frac{1}{n} \sum_{i=1}^n \mathbf{X}_{i, \mathcal{M}} b'''(\mathbf{X}_i^T \beta_0) X_{i, j_0} \{ \mathbf{X}_i^T (\tilde{\beta} - \beta_0) \} \right) \right| \\ & \leq \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \left| \frac{1}{n} \sum_{i=1}^n \mathbf{a}^T \mathbf{X}_{i, \mathcal{M}} \{ b'''(\mathbf{X}_i^T \beta_0) - b'''(\mathbf{X}_i^T \beta_a^*) \} X_{i, j_0} \{ \mathbf{X}_i^T (\tilde{\beta} - \beta_0) \} \right| \\ & \leq L_0 \omega_0^3 \eta_n^2 \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i, \mathcal{M}}|, \quad (\text{B.77}) \end{aligned}$$

with probability tending to 1. Similar to (B.67), we can show

$$\Pr \left(\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}| \leq c_2^* \right) \rightarrow 1,$$

for some constant $c_2^* > 0$. This proves (B.73).

Similarly, to prove (B.74), it suffices to show with probability tending to 1, we have for any $\mathcal{M} \subseteq \mathbb{I}_{j_0}$ such that $|\mathcal{M}| \leq \kappa_n$ and any $\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}$ such that $\|\mathbf{a}\|_2 = 1$,

$$\left| \mathbf{a}^T \left(\widehat{\Sigma}_{\mathcal{M},\mathcal{M}} - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* - \frac{1}{n} \sum_{i=1}^n \mathbf{X}_{i,\mathcal{M}} b'''(\mathbf{X}_i^T \beta_0) \mathbf{X}_{i,\mathcal{M}} \{ \mathbf{X}_i^T (\widetilde{\beta} - \beta_0) \} \right) \mathbf{a} \right| \leq c_3^* \eta_n^2,$$

for some constant $c_3^* > 0$. By Taylor's theorem, we have for any $\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|}$,

$$\mathbf{a}^T \left(\widehat{\Sigma}_{\mathcal{M},\mathcal{M}} - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* \right) \mathbf{a} = \frac{1}{n} \sum_{i=1}^n \mathbf{a}^T \mathbf{X}_{i,\mathcal{M}} b'''(\mathbf{X}_i^T \beta_a^*) \mathbf{X}_{i,\mathcal{M}}^T \mathbf{a} \{ \mathbf{X}_i^T (\widetilde{\beta} - \beta_0) \},$$

for some β_a^* lying on the line segment joining β_0 and $\widetilde{\beta}$. Hence, similar to (B.76) and (B.77), we can show that with probability tending to 1,

$$\begin{aligned} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \left| \mathbf{a}^T \left(\widehat{\Sigma}_{\mathcal{M},\mathcal{M}} - \widehat{\Sigma}_{\mathcal{M},\mathcal{M}}^* - \frac{1}{n} \sum_{j=1}^p \widehat{\Psi}_{\mathcal{M},\mathcal{M}}^{(j)} (\widetilde{\beta}_j - \beta_{0,j}) \right) \mathbf{a} \right| \\ \leq c_4^* \eta_n^2 \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2, \end{aligned} \quad (\text{B.78})$$

for some constant $c_4^* > 0$. Using similar arguments in (B.40), we can show

$$\Pr \left(\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2=1}} \frac{1}{n} \sum_{i=1}^n |\mathbf{a}^T \mathbf{X}_{i,\mathcal{M}}|^2 \leq c_5^* \right) \rightarrow 1, \quad (\text{B.79})$$

for some $c_5^* > 0$. This together with (B.78) proves (B.74). Hence, (B.72) is proven.

Similarly, we can show

$$\max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \left\| \widehat{\sigma}_{\mathcal{M},j_0}^2 - \widetilde{\sigma}_{\mathcal{M},j_0}^2 \right\|_2 \leq \bar{c}_0 \eta_n^2.$$

This together with (B.72) proves (B.4).

Finally, we show

$$\sum_{t=0}^{n-1} \frac{\tilde{Z}_{t+1,j_0} \varepsilon_{t+1}}{\sqrt{n}} \left(\frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)},j_0}^{*3}} \right) = \sum_{t=0}^{n-1} \frac{\hat{Z}_{t+1,j_0}^* \varepsilon_{t+1}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} + o_p(1). \quad (\text{B.80})$$

With some calculations, we have

$$\begin{aligned} & \sum_{t=0}^{n-1} \frac{\tilde{Z}_{t+1,j_0} \varepsilon_{t+1}}{\sqrt{n}} \left(\frac{1}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} - \frac{\sum_j \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)} (\tilde{\beta}_j - \beta_{0,j})}{\hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} \right) - \sum_{t=0}^{n-1} \frac{\hat{Z}_{t+1,j_0}^* \varepsilon_{t+1}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*} \\ &= \underbrace{\sum_{j=1}^p \left(\sum_{t=0}^{n-1} \frac{\hat{Z}_{t+1,j_0}^* \varepsilon_{t+1} \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} (\tilde{\beta}_j - \beta_{0,j}) \right)}_{\eta_1^*} + \underbrace{\sum_{t=0}^{n-1} \frac{(\tilde{Z}_{t+1,j_0} - \hat{Z}_{t+1,j_0}^*) \varepsilon_{t+1}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^*}}_{\eta_2^*} \\ &+ \underbrace{\sum_{j=1}^p \left(\sum_{t=0}^{n-1} \frac{(\tilde{Z}_{t+1,j_0} - \hat{Z}_{t+1,j_0}^*) \varepsilon_{t+1} \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}} (\tilde{\beta}_j - \beta_{0,j}) \right)}_{\eta_3^*}. \end{aligned}$$

In the following, we first prove $\eta_1^* = o_p(1)$. Note that $|\eta_1^*| \leq \max_j |\eta_{1,j}^*| \|\tilde{\beta} - \beta_0\|_1$ where

$$\eta_{1,j}^* = \sum_{t=0}^{n-1} \frac{\hat{Z}_{t+1,j_0}^* \varepsilon_{t+1} \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{(j)}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)},j_0}^{*3}}.$$

By Condition (A5*), it suffices to show $\max_j |\eta_{1,j}^*| = O_p(\sqrt{\kappa_n \log p} + \log p)$.

The function $b'''(\cdot)$ is Lipschitz continuous. Hence, there exists some constant L_0 such that

$$|b'''(\mathbf{X}_i^T \beta_0) - b'''(0)| \leq L_0 |\mathbf{X}_i^T \beta_0|.$$

By Condition (A4*), we obtain

$$\max_{1 \leq i \leq n} |b'''(\mathbf{X}_i^T \beta_0)| \leq c_6^*, \quad (\text{B.81})$$

for some constant $c_6^* > 0$. By Condition (A4*), we have

$$\max_{1 \leq j \leq p} |\hat{\Psi}_{j_0,j_0}^{(j)}| = \max_{1 \leq j \leq p} \left| \frac{1}{n} \sum_i X_{i,j}^3 b'''(\mathbf{X}_i^T \beta_0) \right| \leq \omega_0^3 c_6^*. \quad (\text{B.82})$$

Besides, by (B.79), (B.81) and Condition (A4*), we have

$$\begin{aligned}
& \max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Psi}_{\mathcal{M}, \mathcal{M}}^{(j)}\|_2 = \max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} |\mathbf{a}^T \widehat{\Psi}_{\mathcal{M}, \mathcal{M}}^{(j)} \mathbf{a}| \\
&= \max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \left| \frac{1}{n} \sum_{i=1}^n (\mathbf{a}^T \mathbf{X}_{i, \mathcal{M}})^2 X_{i, j} b'''(\mathbf{X}_i^T \boldsymbol{\beta}_0) \right| \leq c_6^* \omega_0 \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \sup_{\substack{\mathbf{a} \in \mathbb{R}^{|\mathcal{M}|} \\ \|\mathbf{a}\|_2 = 1}} \left| \frac{1}{n} \sum_{i=1}^n (\mathbf{a}^T \mathbf{X}_{i, \mathcal{M}})^2 \right| \\
&\leq c_5^* c_6^* \omega_0,
\end{aligned} \tag{B.83}$$

with probability tending to 1. Similarly, we can show

$$\max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \|\widehat{\Psi}_{\mathcal{M}, j_0}^{(j)}\|_2 \leq c_2^* c_6^* \omega_0^2, \tag{B.84}$$

with probability tending to 1. This together with (B.71), (B.82) and (B.83) yields

$$\Pr \left(\max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0}, |\mathcal{M}| \leq \kappa_n}} |\hat{\xi}_{\mathcal{M}, j_0}^{(j)}| \leq c_7^* \right) \rightarrow 1, \tag{B.85}$$

for some constant $c_7^* > 0$.

Besides, similar to (B.12), we can show

$$\Pr \left\{ \max_{t \in [0, \dots, n-1]} \left| \widehat{Z}_{t+1}^* \right| \leq \omega_0 \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}} \right) \right\} \rightarrow 1. \tag{B.86}$$

Note that

$$\eta_{1, j}^* = \underbrace{\sum_{t=0}^{s_n-1} \frac{\widehat{Z}_{t+1, j_0}^* \varepsilon_{t+1} \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{(j)}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{*3}}}_{\eta_{1, j}^{**}} + \underbrace{\sum_{t=s_n}^{n-1} \frac{\widehat{Z}_{t+1, j_0}^* \varepsilon_{t+1} \hat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{(j)}}{\sqrt{n} \hat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{*3}}}_{\eta_{1, j}^{***}}.$$

We first prove $\max_j |\eta_{1, j}^{***}| = O_p(\sqrt{\kappa_n \log p} + \log p)$. Define

$$\varepsilon_t^* = \varepsilon_t I(|\varepsilon_t| \leq n^{-1/3} c_n),$$

for some sequence diverging c_n which will be specified later. By Bonferroni's inequality and Markov's inequality, we have

$$\Pr \left\{ \bigcup_{t=1}^n (\varepsilon_t^* \neq \varepsilon_t) \right\} \leq n \Pr(\varepsilon_t^* \neq \varepsilon_t) \leq n \Pr(|\varepsilon_0| > n^{-1/3} c_n) \leq n \frac{\mathbb{E}|\varepsilon_0|^3}{n c_n^3} \rightarrow 0.$$

This implies

$$\eta_{1,j}^{***} = \sum_{t=s_n}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}}, \quad (\text{B.87})$$

with probability tending to 1. Note that

$$\sum_{t=s_n}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} = \underbrace{\sum_{t=s_n}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \{\varepsilon_{t+1}^* - \mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1})\} \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}}}_{\eta_{1,j}^{***}} + \underbrace{\sum_{t=s_n}^{n-1} \frac{\widehat{Z}_{t+1,j_0}^* \mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1}) \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}}}_{\eta_{1,j}^{****}}.$$

Since $\mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1}) = 0$, we have

$$|\mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1})| = |\mathbb{E}(\varepsilon_{t+1}^* I(|\varepsilon_{t+1}^*| > n^{1/3} c_n) | \mathbf{X}_{t+1})| \leq \frac{\mathbb{E}(|\varepsilon_{t+1}^*|^3 | \mathbf{X}_{t+1})}{n^{2/3} c_n^2}.$$

By Condition (A6*), there exists some constant $c_8^* > 0$ such that

$$\max_{0 \leq t \leq n-1} |\mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1})| \leq c_8^* n^{-2/3} c_n^{-2}.$$

Combining this together with (B.9), (B.85) and (B.86) yields

$$\max_j |\eta_{1,j}^{*****}| \leq \sqrt{n} c_7^* c_8^* \omega_0 \left(1 + 2\sqrt{\frac{\kappa_n c_0}{\bar{c}}}\right) n^{-2/3} c_n^{-2} \frac{8}{\bar{c}^{3/2}},$$

with probability tending to 1. Under the given conditions, we have $\kappa_n^3 = o(n)$. Hence, we've shown

$$\max_j |\eta_{1,j}^{*****}| = o_p(1). \quad (\text{B.88})$$

Define $\sigma(\mathcal{F}_t^*) = \sigma(\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n, Y_1, Y_2, \dots, Y_t)$, $\eta_{1,j}^{***}$ corresponds to a mean-zero martingale with respect to the filtration $\{\sigma(\mathcal{F}_t^*) : t \geq s_n\}$. By Condition (A1*) and (A4*), we have for any $t = 0, \dots, n-1$,

$$|\widehat{Z}_{t+1,j_0}^*| \leq \omega_0 \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} (1 + \sqrt{\kappa_n} \|\widehat{\omega}_{\mathcal{M},j_0}^*\|_2). \quad (\text{B.89})$$

Let

$$\bar{c}_n^{(j)} \equiv c_n \omega_0 n^{-1/6} \max_{1 \leq j \leq p} \max_{\substack{\mathcal{M} \subseteq \mathbb{I}_{j_0} \\ |\mathcal{M}| \leq \kappa_n}} \frac{|(1 + \sqrt{\kappa_n} \|\widehat{\omega}_{\mathcal{M},j_0}^*\|_2)|}{|\widehat{\sigma}_{\mathcal{M},j_0}^*|} |\widehat{\xi}_{\mathcal{M},j_0}^{(j)}|.$$

By Condition (A1*), (B.89) and the definition of ε_t^* , we have for any t and $1 \leq j \leq p$,

$$\Pr \left(\left| \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right| \leq \bar{c}_n^{(j)} \middle| \mathcal{F}_t^* \right) = 1. \quad (\text{B.90})$$

Hence, by Jensen's inequality, we have

$$\Pr \left(\left| \frac{\widehat{Z}_{t+1,j_0}^* \mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1}) \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right| \leq \bar{c}_n^{(j)} \middle| \mathcal{F}_t^* \right) = 1.$$

This together with (B.90) gives

$$\Pr \left(\left| \frac{\widehat{Z}_{t+1,j_0}^* \{\varepsilon_{t+1}^* - \mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1})\} \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right| \leq 2\bar{c}_n^{(j)} \middle| \mathcal{F}_t^* \right) = 1.$$

Besides, it follows from Hölder's inequality that

$$\begin{aligned} & \sum_{t=s_n}^{n-1} \mathbb{E} \left\{ \left| \left(\frac{\widehat{Z}_{t+1,j_0}^* \{\varepsilon_{t+1}^* - \mathbb{E}(\varepsilon_{t+1}^* | \mathbf{X}_{t+1})\} \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right)^2 \right| \middle| \mathcal{F}_t^* \right\} \\ & \leq \sum_{t=s_n}^{n-1} \mathbb{E} \left\{ \left| \left(\frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right)^2 \right| \middle| \mathcal{F}_t^* \right\} \leq \sum_{t=s_n}^{n-1} \mathbb{E} \left\{ \left| \left(\frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*3}} \right)^2 \right| \middle| \mathcal{F}_t^* \right\} \\ & \leq \sum_{t=s_n}^{n-1} \frac{(\widehat{Z}_{t+1,j_0}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)})^2}{n \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*6}} \mathbb{E}(\varepsilon_{t+1}^2 | \mathbf{X}_{t+1}) \leq \underbrace{\sum_{t=s_n}^{n-1} \frac{(\widehat{Z}_{t+1,j_0}^* \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{(j)})^2}{n \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0,j_0}^{(t)}}^{*6}} \mathbb{E}(|\varepsilon_{t+1}|^3 | \mathbf{X}_{t+1})}_{V_n^{(j)}} \}^{2/3}. \end{aligned}$$

Therefore, it follows from Theorem 9.12 in de la Peña et al. (2009) that for any $1 \leq j \leq p$,

$$\Pr(|\eta_{1,j}^{****}| > z) \leq 2 \exp \left(-\frac{z^2}{2(V_n^{(j)} + \bar{c}_n^{(j)} z)} \right).$$

Take $z_0^{(j)} = 6\{(V_n^{(j)} \log p)^{1/2} + (\bar{c}_n^{(j)}) \log p\}$, we have

$$\begin{aligned} \Pr(|\eta_{1,j}^{****}| > z_0^{(j)}) & \leq 2 \exp \left(-\frac{36V_n^{(j)} \log p + 36(\bar{c}_n^{(j)})^2 \log^2 p}{2V_n^{(j)} + 12\bar{c}_n^{(j)} (V_n^{(j)} \log p)^{1/2} + 12(\bar{c}_n^{(j)})^2 \log p} \right) \\ & \leq 2 \exp \left(-\frac{36V_n^{(j)} \log p + 36(\bar{c}_n^{(j)})^2 \log^2 p}{8V_n^{(j)} + 18(\bar{c}_n^{(j)})^2 \log p} \right) \leq 2 \exp(-2 \log p) = \frac{2}{p^2}. \end{aligned}$$

It follows from Bonferroni's inequality that

$$\Pr \left(\bigcap_{j=1}^p \left\{ |\eta_{1,j}^{****}| > z_0^{(j)} \right\} \right) \leq \sum_{j=1}^p \Pr(|\eta_{1,j}^{****}| > z_0^{(j)}) = \frac{2}{p}.$$

This implies that

$$\max_{j=1}^p |\eta_{1,j}^{****}| \leq \max_{j=1}^p z_0^{(j)}. \quad (\text{B.91})$$

Set $c_n = \log^{1/3} n$. Under the given conditions, we have $n^{-1/6} \sqrt{\kappa_n} c_n = o(1)$. By Condition (A1*), (A6*), (B.9), (B.71), (B.85) and (B.86), we can show

$$\max_{j=1}^p \bar{c}_n^{(j)} = o_p(1) \quad \text{and} \quad \max_{j=1}^p V_n^{(j)} = O_p(\kappa_n).$$

In view of (B.91), we've shown $\max_j |\eta_{1,j}^{****}| = O_p(\sqrt{\kappa_n \log p} + \log p)$. This further implies together with (B.87) and (B.88) yields

$$\max_j |\eta_{1,j}^{***}| = O_p(\sqrt{\kappa_n \log p} + \log p). \quad (\text{B.92})$$

Recall that

$$\eta_{1,j}^{**} = \sum_{t=0}^{s_n-1} \frac{\widehat{Z}_{t+1,j_0}^* \varepsilon_{t+1} \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(-s_n)}, j_0}^{*3}}.$$

Given $\mathbf{X}_1, \dots, \mathbf{X}_n$ and Y_{s_n+1}, \dots, Y_n , each term in $\eta_{1,j}^{**}$ is independent of others. Using similar arguments, we can show $\max_j |\eta_{1,j}^{**}| = O_p(\sqrt{\kappa_n \log p} + \log p)$. This together with (B.92) gives $\max_j |\eta_{1,j}^*| = O_p(\sqrt{\kappa_n \log p} + \log p)$. By Condition (A5*), we obtain $|\eta_1^*| \leq \max_j \max_j |\eta_{1,j}^*| \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|_1 = o_p(1)$. Similarly, we can show $\eta_2^* = o_p(1)$. It remains to show $\eta_3^* = o_p(1)$.

Note that $|\eta_3^*|$ can be upper bounded by $\max_j |\eta_{3,j}^*| \|\widetilde{\boldsymbol{\beta}} - \boldsymbol{\beta}_0\|_1$ where

$$\eta_{3,j}^* = \sum_{t=0}^{n-1} \frac{(\widetilde{Z}_{t+1,j_0} - \widehat{Z}_{t+1,j_0}^*) \varepsilon_{t+1} \widehat{\xi}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{(j)}}{\sqrt{n} \widehat{\sigma}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{*3}}.$$

Since

$$\widehat{Z}_{t+1,j_0}^* - \widetilde{Z}_{t+1,j_0} = \sum_{j=1}^p \mathbf{X}_{t+1, \widehat{\mathcal{M}}_{j_0}^{(t)}}^T \widehat{\boldsymbol{\Sigma}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, \widehat{\mathcal{M}}_{j_0}^{(t)}}^{*-1} \left(\widehat{\boldsymbol{\Psi}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^{(j)} + \widehat{\boldsymbol{\Psi}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, \widehat{\mathcal{M}}_{j_0}^{(t)}} \widehat{\boldsymbol{\omega}}_{\widehat{\mathcal{M}}_{j_0}^{(t)}, j_0}^* \right) (\widetilde{\beta}_j - \beta_{0,j}).$$

By Condition (A1*), (A4*), (A5*), (B.70), (B.71), (B.83) and (B.84), we can show

$$\Pr \left(\max_{0 \leq t \leq n-1} |\widehat{Z}_{t+1,j_0}^* - \widetilde{Z}_{t+1,j_0}| \leq c_9^* \sqrt{\kappa_n} \eta_n \right) \rightarrow 1,$$

for some constant $c_9^* > 0$. Combining this together with Condition (A1*), (B.9) and (B.85) yields

$$\max_j |\eta_{3,j}^*| \leq c_{10}^* \sqrt{n \kappa_n} \eta_n \left(\frac{1}{n} \sum_{t=0}^{n-1} |\varepsilon_{t+1}| \right). \quad (\text{B.93})$$

By Condition (A6*) and Hölder's inequality, we have

$$\mathbb{E} \left(\frac{1}{n} \sum_{t=0}^{n-1} |\varepsilon_{t+1}| \right) = \mathbb{E} |\varepsilon_0| \leq (\mathbb{E} |\varepsilon_0|^3)^{1/3} = O(1).$$

Hence, it follows from Markov's inequality that $\sum_{t=0}^{n-1} |\varepsilon_{t+1}|/n = O_p(1)$. This together with (B.93) implies that $\max_j |\eta_{3,j}^*| = O_p(\sqrt{n \kappa_n} \eta_n)$. Therefore, we have $\eta_3^* = O_p(\sqrt{n \kappa_n} \eta_n^2)$. By Condition (A5*), we obtain $\eta_3^* = o_p(1)$. The proof is hence completed.

B.8 Technical lemmas

Lemma B.2 *For any positive definite matrix*

$$\Psi = \begin{pmatrix} \Psi_{11} & \Psi_{12} \\ \Psi_{21} & \Psi_{22} \end{pmatrix},$$

denote its inverse matrix as Ω and partition it into $\Omega_{11}, \dots, \Omega_{22}$ accordingly. Then,

$$\Omega_{11} = (\Psi_{11} - \Psi_{12} \Psi_{22}^{-1} \Psi_{21})^{-1}.$$

Besides, let $\Psi_ = \Psi_{22} - \Psi_{21} \Psi_{11}^{-1} \Psi_{12}$, we have*

$$\Omega = \begin{pmatrix} \Psi_{11}^{-1} + \Psi_{11}^{-1} \Psi_{12} \Psi_*^{-1} \Psi_{21} \Psi_{11}^{-1} & -\Psi_{11}^{-1} \Psi_{12} \Psi_*^{-1} \\ -\Psi_*^{-1} \Psi_{21} \Psi_{11}^{-1} & \Psi_*^{-1} \end{pmatrix}.$$

References

Adamczak, R. a. (2008). A tail inequality for suprema of unbounded empirical processes with applications to Markov chains. *Electron. J. Probab.* 13, no. 34, 1000–1034.

de la Peña, V. H., T. L. Lai, and Q.-M. Shao (2009). *Self-normalized processes*. Probability and its Applications (New York). Springer-Verlag, Berlin. Limit theory and statistical applications.

Klartag, B. and S. Mendelson (2005). Empirical processes and random projections. *J. Funct. Anal.* 225(1), 229–245.

van der Vaart, A. W. and J. A. Wellner (1996). *Weak Convergence and Empirical Processes*. New York: Springer.