

Linearity of Expected Value

Math 252

Transformations and Sums

Transformations

If X is a random variable and f is a function, then $f(X)$ is a new random variable such that

$$p(f(X) = b) = \sum_{\{a|f(a)=b\}} p(X = a)$$

This is actually much simpler than it looks. Basically, we just transform each possible value k that X can take into $f(k)$ and then collapse any cells in the probability table that end up having the same value.

1. Below is a probability table for X . Create probability table for X^2 and $2X + 3$.

value	-1	0	1
probability	0.2	0.5	0.3

2. Below is a probability table for X . Create probability tables for X^2 and for $4 - X$.

value	0	1	2	3
probability	0.1	0.2	0.3	0.4

3. Below is a probability table for X . Create probability tables for X^2 and for $2X$.

value	0	1
probability	0.7	0.3

Sums

Consider a process that produces multiple numerical outcomes. (Example: roll a red die and a blue die. The number on the red die is one numerical result, the number on the blue die is a second numerical result.) We can represent

$$p(X = a \wedge Y = b)$$

with a function of a and b or with a two-way table like this one:

	X = 1	X = 2	X = 3
Y = 1	0.10	0.10	0.25
Y = 2	0.10	0.05	0.10
Y = 3	0.10	0.15	0.05

Use the table above to answer the following questions.

4. What is $p(X = 3 \wedge Y = 2)$? What is $p(X = 3)$? What is $p(Y = 2)$?
5. What is $p(X = Y)$? What is $p(X > Y)$?
6. Create three probability tables, one for X , one for Y , and one for $X + Y$. [Hint: Start by determining the possible values.]
7. Now compute $E(X)$, $E(Y)$, and $E(X + Y)$.

Linearity of Expected Value

Linearity Theorem for Expected Value. Expected value is linear in the following senses:

- a. $E(aX + b) = aE(X) + b$
- b. $E(X_1 + X_2 + X_3 + \cdots + X_n) = E(X_1) + E(X_2) + E(X_3) + \cdots + E(X_n)$

Exercises

8. Are your results for #9 consistent with the theorem? Explain.
9. Let S be the sum on two fair 4-sided dice. Compute $E(S)$ two ways: (a) by creating a probability table for S , and (b) by writing $S = X + Y$ and using the Linearity Theorem.
10. What is the expected value of the sum of two fair 6-sided dice? (You only have to do this one way.)
11. Here is a more interesting example. Suppose 5 people put their names into a hat. Then each person pull one name out. Let X be the number of people who draw their own name. What is $E(X)$? How would the answer change if 10 people put their names into the hat? 20? 1? [Hint: How can you write X as a sum?]
12. Let $X \sim \text{Binom}(1, p)$. What is $E(X)$? What is $E(X^2)$?
13. Let $X \sim \text{Binom}(n, p)$. What is $E(X)$? [Hint: Use the previous problem and the Linearity Lemma. How can you write X as a sum?]
14. Can you prove the linearity theorem? [For part b, you may restrict your attention to a sum of two random variables.]
15. What is the expected number of diamonds in a 5-card hand dealt from a standard deck?