

Counting and Probability

R Pruim

2019-02-22

Contents

Preface	2
1 Some Counting Problems	1
2 Pigeonhole Principle	3

Preface

These exercises were assembled to accompany a Discrete Mathematics course at Calvin College.

1 Some Counting Problems

Each of these problems involves determining how many of something there are. For most, there will be too many to simply make a list and count them off, so you will need to “count without counting”. As you go along, make a list of important counting principles you are using.

1. A new company with just two employees, Sanchez and Patel, rents a floor of a building with 12 offices. How many ways are there to assign different offices to these two employees?
2. The chairs of an auditorium are labeled with a letter (indicating the row) and a positive integer not exceeding 100 (indicating the seat within the row). Give an upper bound on the number of seats in the auditorium.
3. There are 32 microcomputers in a computer center. Each microcomputer has 24 ports. How many different ports to a microcomputer in the center are there?
4. How many bitstrings of length 8 are there?
5. How many bitstrings of length 8 either start with a 1 or end with a 1?
6. A state issues license plates that consist of three letters followed by three numbers. How many such plates are there?
7. A graph isomorphism is a 1-1 and onto function from the vertices in one graph to the vertices in another (that additionally preserves edges). You decide to write a little program that will check each 1-1 and onto function to see if it an isomorphism.


If your graphs have n vertices, how many such functions must you check?

8. Now you modify your graph isomorphism algorithm so that it only tries bijections (1-1 and onto functions) that respect the degrees of the vertices. If the degree sequences of two graphs are both $(5, 5, 4, 4, 4, 3, 3, 2)$, how many functions must you check? How much better is this than your previous algorithm (on these two graphs)?
9. Let's use the following coding scheme to describe possible telephone numbers.
 - let X denote a digit that can take on any integer value $0 - 9$;
 - let N denote a digit that can take on values $2 - 9$
 - let Y denote a digit that can only take on values 0 or 1.
 - a. In the scheme used in the 1960's, telephone numbers had to have the form NYX-NNX-XXXX. How many such phone numbers are there?
 - b. A new scheme (*The North American Numbering Plan*) was introduced that required telephone numbers have the form NXX-NXX-XXXX. How many more phone numbers are available in the new scheme?
10. How many password of length 6, 7, or 8 are there if the passwords must be alpha-numeric, capitalization doesn't matter, and there must be at least one digit?
11. In the previous example, how many passwords are there if the first character must be a letter?
12. Professor Pruim puts the names of 30 students into a hat. He draws one out and gives that person a chocolate bar. Then he draws out another name, and gives that person a bag of M and M's. Finally, he draws out a third name and gives that person a bag of Valentine hearts he picked up cheap on February 15.
 - a. How many different ways can the prizes be distributed?
 - b. If you are one of the 30 students, how many of those ways include you getting a prize?

Answers/Solutions

1. $12 * 11 = 132$.
2. $26 * 100 = 2600$
3. $32 * 24 = 768$.
4. $2^8 = 256$
5. Two ways:
 - a. only bad thing is beginning AND ending with 0, so $2^8 - 2^6 = 192$.
 - b. Start with 1 or end with 1 but not both: $2^7 + 2^7 - 2^6 = 192$
6. $10^3 \cdot 26^3 = 17576000$
7. $n!$
8. $2 \cdot 1 \cdot 3 \cdot 2 \cdot 1 \cdot 2 \cdot 1 \cdot 1 = 24$. ($8! = 40320$, so that's a pretty big speed up.)
9. See book.
10. $36^6 - 26^6 + 36^7 - 26^7 + 36^8 - 26^8 = 2684483063360$
11. $26 * (36^5 - 26^5) + 26 * (36^6 - 26^6) + 26 * (36^7 - 26^7) = 1878468937280$
12. a. $30 * 29 * 28 = 24360$
b. $1 * 29 * 28 + 29 * 1 * 28 + 29 * 28 * 1 = 30 * 29 * 28 - 29 * 28 * 27 = 2436$

2 Pigeonhole Principle



Big Idea: It is not possible for everyone to be above average or for everyone to be below average. (It is possible for everyone to be exactly average.)

So if n things are distributed into k groups,

- The largest group will have at least $\lceil n/k \rceil$ things, and
- The smallest group will have at most $\lfloor n/k \rfloor$ things.

This is called the (generalized) pigeon hole principle based on the analogy of putting letters into mail slots (called pigeon holes).

Applying the pigeonhole principle is generally easy *once you figure out what the pigeons (letters, really) and the pigeonholes are*. Figuring out the pigeons and holes can be easy, or it can take a clever idea. (Some famous proofs center on very clever uses of the pigeonhole principle.)

1. Suppose (in some alternative reality) that Professor Pruim brought 150 cookies to a class with 34 students. At the end of the hour, all of the cookies had been eaten by the students.
 - a. What is the average number of cookies eaten by a student?
 - b. This means the at least one student had at least _____ cookies.
 - c. This means the at least one student had at most _____ cookies.

In each of the following problems, each time you use the pigeonhole principle, identify the “pigeons” and the “holes” and explain how that solves the problem. If you don’t use the pigeonhole principle directly, be sure to explain your reasoning carefully.

2. Fill in the blanks.
 - a. In a class of 38 students, there are always at least _____ students who were born in the same month.
 - b. In a class of 38 students, there are always at least _____ students who were born on the same day of the week.
3. In the 2014 Winter Olympics, 88 nations competed in 98 events. Explain how this guaranteed that some country would win more than one gold medal. (No, the answer is not “Norway was there.”)
4. At the 2014 Winter Olympics, Norway won 11 gold medals. Does that fact alone guarantee that a country won no gold medals?

5. Joe has 21 white socks and 15 black socks. (Don't ask what happened to the missing socks, no one knows where they go.)
 - a. If he pulls socks in the dark (so he can't tell what color they are), how many must he grab to make sure he has a pair of white socks?
 - b. If he pulls socks in the dark (so he can't tell what color they are), how many must he grab to make sure he has a pair of black socks?
 - c. If he pulls socks in the dark (so he can't tell what color they are), how many must he grab to make sure he has a pair of matching socks?
6. For his birthday, Joe received a gift of 8 red socks (4 pairs, and he hasn't lost any yet). Now how many socks must Joe grab to be sure that he has a matching pair?
7. Joe is going on a trip and needs 3 pairs of matching socks. He's still too lazy to turn on the light to see what socks he is grabbing. How many socks must Joe bring along to make sure he has 3 matching pairs? (He doesn't care how many pairs are red, white, or black.)
8. Checkers.
 - a. Consider a 3×9 grid (like an odd-shaped checker board). Show that no matter how you place red and black checkers on the 27 squares, there will always be a rectangle whose 4 corners have the same color checker. (The rectangles must be at least 2×2 ; 1×1 rectangles don't have four different checkers on their corners.)
 - b. Consider a 3×7 grid (like an odd-shaped checker board). Show that no matter how you place red and black checkers on the 21 squares, there will always be a rectangle whose 4 corners have the same color checker. (The rectangles must be at least 2×2 ; 1×1 rectangles don't have four different checkers on their corners.)
 - c. Show how to place red and black checkers on a 3×6 grid without having any rectangles with mono-chromatic corners.
9. Show that there is an integer that is divisible by 37 and consists only of 1's and 0's in its decimal representation.
 Hint: Consider the numbers 1, 11, 111, 1111, 11111, etc. Show that two of them are congruent mod 37. How does that help you answer the question?
10. Was there anything special about the number 37 in the previous example? For what other divisors does this work? Explain.
11. Is there anything special about the digits 1 and 0 in the previous example? What other combinations of digits would work?
12. A computer lab has 15 workstations and 10 servers. Workstations can only access servers via direct connections, and each server can only use one of these connections at a time.
 - a. If we connect every workstation to every server, how many connections is that?
 - b. Show that 60 connections suffice make it possible to use any set of 10 workstations simultaneously. (No matter what we do, we won't be able to connect 11 workstations to servers since there are only 10 servers.)
 - c. Show that 59 connections is not enough.
 That is, no matter how you do the connecting, there will always be a combination of 10 workstations that cannot simultaneously connect to servers. (Hint: on average, how many connections are there per server?)
13. The integers from 1 to 10 are distributed around a circle. Prove that there must be three neighbors whose sum is at least 17. What about 18? What about 19?