






GarmanKlass (JPM, daily 2019-10-30 to 2024-10-29)






GarmanKlass (JPM, daily 2019-10-30 to 2024-10-29)




GarmanKlass (JPM, daily 2019-10-30 to 2024-10-29)





GarmanKlass (JPM, daily 2019-10-30 to 2024-10-29)




Distribution of GarmanKlass estimator values (JPM, daily 2019-10-30 to 2024-10-29)



GarmanKlass (JPM v. XLF, daily 2019-10-30 to 2024-10-29)



GarmanKlass (Correlation of JPM v. XLF, daily 2019-10-30 to 2024-10-29)



OLS Regression Results

```

=====
Dep. Variable:          y      R-squared (uncentered):          0.993
Model:                  OLS    Adj. R-squared (uncentered):          0.993
Method:                 Least Squares    F-statistic:          1.695e+05
Date:                  Tue, 29 Oct 2024    Prob (F-statistic):          0.00
Time:                  23:53:05    Log-Likelihood:          3025.4
No. Observations:      1229    AIC:          -6049.
Df Residuals:          1228    BIC:          -6044.
Df Model:              1
Covariance Type:       nonrobust
=====

```

	coef	std err	t	P> t	[0.025	0.975]
x1	1.2911	0.003	411.672	0.000	1.285	1.297

```

=====
Omnibus:              40.009    Durbin-Watson:          0.036
Prob(Omnibus):        0.000    Jarque-Bera (JB):        77.183
Skew:                 0.214    Prob(JB):                1.74e-17
Kurtosis:             4.151    Cond. No.:               1.00
=====

```

Notes:

- [1] R^2 is computed without centering (uncentered) since the model does not contain a constant.
- [2] Standard Errors assume that the covariance matrix of the errors is correctly specified.