

Example Tables

Calvin Whealton

February 3, 2016

List of Tables

1	A table where only the table values are written, not the caption, label, and horizontal lines at the beginning and end of the table.	2
2	OLS table all from R	2
3	OLS table all from R, no Constant	3

This is an example file showing how regression tables from R can be imported into \LaTeX using the code available on https://github.com/calvinwhealton/R_tables_to_LaTeX.

1 Ordinary Least Squares

Table 1: A table where only the table values are written, not the caption, label, and horizontal lines at the beginning and end of the table.

SSE	87.9		DoF	97
SSR	2222.92		R2	0.962
SST	2310.82		Adj.R2	0.9608
RSE	0.952			
Variable	Est. Coeff.	Std. Error	t Stat.	Pr(> t)
(Intercept)	4.467e+00	2.763e-01	1.616e+01	2.834e-29
x1	5.594e-01	5.762e-02	9.708e+00	5.733e-16
x2	2.046e+00	4.889e-02	4.184e+01	7.003e-64

Table 2: A table that is entirely written from R including captions (long and short) and label.

SSE	87.9		DoF	97
SSR	2222.92		R2	0.962
SST	2310.82		Adj.R2	0.9608
RSE	0.952			
Variable	Est. Coeff.	Std. Error	t Stat.	Pr(> t)
b0	4.467e+00	2.763e-01	1.616e+01	2.834e-29
bx1	5.594e-01	5.762e-02	9.708e+00	5.733e-16
bx2	2.046e+00	4.889e-02	4.184e+01	7.003e-64

2 Nonlinear Least Squares

3 Ordinary Least Squares

Table 3: A table that is entirely written from R including captions (long and short) and label. There is no constant in this model.

SSE	324.67		DoF	98
SSR	3708.83		R2	0.8595
SST	2310.82		Adj.R2	0.8566
RSE	1.82			
Variable	Est. Coeff.	Std. Error	t Stat.	Pr(> t)
cx1	1.018e+00	9.588e-02	1.062e+01	5.537e-18
cx2	2.496e+00	7.688e-02	3.246e+01	3.078e-54