

Indirect measurement of cosmic-ray proton spectrum using γ -ray data from *Fermi* Large Area Telescope

Patomporn Payoungkhamdee[†]

under supervision of

Asst. Prof. Warit Mitthumsiri[†], Prof. David John Ruffolo[†]

[†]Mahidol University

Abstract

Cosmic rays (CRs) are high-energy particles propagating in space. It is mainly constituted by protons and the rigidity spectrum is well described by a power law. Recent measurements by PAMELA and AMS-02 indicate an abrupt change of the CR proton spectral index at about 336 GV. When protons interact with the Earth's upper atmosphere, γ -rays can be produced and detected by space-based detectors. The Earth Limb γ -ray data was collected by the *Fermi* Large Area Telescope (LAT) along with proton-air interaction models to determine the CR proton spectral indices that best fit the γ -ray data.

Keywords— Cosmic rays, gamma-rays, *Fermi*-LAT

References

- [1] M. Ackermann et al., (Fermi LAT Collaboration), Phys. Rev. Lett. 112, 151103 (2014)
- [2] M. Aguilar et al. (AMS Collaboration), Phys. Rev. Lett. 114, 171103 (2015)
- [3] J. Alcaraz et al., Physics Letters B 490, 27 (2000)
- [4] S. Haino et al., Physics Letters B 594, 35 (2004)