


Accepted Manuscript

Title: HEPATITIS E VIRUS IN HEMATOPOIETIC STEM CELL TRANSPLANT RECIPIENTS: A SYSTEMATIC REVIEW

Authors: Sara Cruz, Carla Campos, Mafalda Timóteo, Ana Tavares, Maria São José Nascimento, Rui Medeiros, Hugo Sousa



PII: S1386-6532(19)30171-4
DOI: <https://doi.org/10.1016/j.jcv.2019.08.002>
Reference: JCV 4177

To appear in: *Journal of Clinical Virology*

Received date: 15 May 2019
Revised date: 30 July 2019
Accepted date: 1 August 2019

Please cite this article as: Cruz S, Campos C, Timóteo M, Tavares A, José Nascimento MS, Medeiros R, Sousa H, HEPATITIS E VIRUS IN HEMATOPOIETIC STEM CELL TRANSPLANT RECIPIENTS: A SYSTEMATIC REVIEW, *Journal of Clinical Virology* (2019), <https://doi.org/10.1016/j.jcv.2019.08.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

TITLE:

HEPATITIS E VIRUS IN HEMATOPOIETIC STEM CELL TRANSPLANT RECIPIENTS: A SYSTEMATIC REVIEW

Running Title:

HEV in HSCT recipients: a systematic review

AUTHORS

Sara Cruz¹, Carla Campos², Mafalda Timóteo¹, Ana Tavares³, Maria São José Nascimento^{4,5}, Rui Medeiros^{1,6,7}, Hugo Sousa^{1,6,8,9}

Affiliations

1 Molecular Oncology and Viral Pathology Group (CI-IPOP), Portuguese Oncology Institute of Porto, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, PORTUGAL

2 Microbiology Service, Portuguese Oncology Institute of Porto, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, PORTUGAL

3 Pathology Department, Portuguese Oncology Institute of Porto, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, PORTUGAL

4 Microbiology Laboratory, Biological Sciences Department, Pharmacy Faculty, University of Porto, 4050-313 Porto, Portugal

5 Epidemiology Research Unit, Public Health Institute, University of Porto, 4050-313 Porto, Portugal

6 Virology Service, Portuguese Oncology Institute of Porto, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, PORTUGAL

7 Research Department, Portuguese League Against Cancer (Liga Portuguesa Contra o Cancro-Núcleo Regional do Norte), Estrada interior da Circunvalação 6657, 4200-172 Porto, Portugal.

8 Life and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

9 ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal

Correspondence to:

Hugo Sousa, MD PhD

*Virology Service – Laboratories 4th floor, Portuguese Oncology Institute of Porto FG EPE
Rua Dr. António Bernardino Almeida, 4200-072 Porto, Portugal.*

Phone: +351 22 508 4000 (ext 5410)

Fax: +351 22 508 4001

E-mail: hugo.sousa@ipopoporto.min-saude.pt

Abstract word count 229

Manuscript word count 2082

Highlights

- Hematopoietic stem cell transplant recipients are at high risk of HEV infection
- The overall prevalence of HEV infection differs according to the diagnostic method
- The overall anti-HEV IgM/IgG seroprevalance was 2.0% and 11.4%, respectively
- HEV RNA prevalence in HSCT recipients is 1.50%
- HEV infection is underestimated in HSCT recipients

Abstract: Background: In developed countries, Hepatitis E virus (HEV) infections, especially by HEV-3, are frequently associated with asymptomatic infection or self-limiting acute hepatitis, although it has been described as a cause of chronic infection, especially in immunocompromised hosts. Hematopoietic stem cell transplant (HSCT) recipients have been recognized as an important risk group for HEV infection due to their prolonged immunosuppression state.

Objectives: We aimed to perform a systematic review of published data to evaluate HEV infection prevalence among HSCT recipients.

Study Design: Literature search was performed concerning published manuscripts regarding 'hepatitis E virus AND stem cell transplantation' following the Preferred Reporting of Systematic Reviews and MetaAnalyses (PRISMA) guidelines. Statistical analysis was performed using the MetaXL software to estimate the overall prevalence of HEV infection according to the different diagnostic approaches (HEV RNA and anti-HEV IgM and/or IgG detection).

Results: A total of 7 manuscripts were included for data analysis, with 6 studies performed in Europe and 1 study in China. Regarding HEV RNA detection, the overall HEV infection prevalence was 1.50% (95% CI: 0.70- 2.60). The overall anti-HEV IgM seroprevalence was 2.00% (95% CI: 0.30- 4.50), and anti-HEV IgG was 11.4% (95% CI: 1.80-26.3).

Conclusions: This systematic review reveals that the overall prevalence of HEV infection in HSCT patients differ according to the diagnostic, thus emphasizing the need of more studies to increase the data regarding prevalence and incidence in HSCT recipients.

Keywords: Hepatitis E virus; HEV; stem cell transplantation; HSCT; prevalence; systematic review.

BACKGROUND

Hepatitis E virus (HEV) was recently recognized as the most common cause of acute viral hepatitis worldwide, with the World Health Organization (WHO) estimating 20 million infections, >3 million acute cases, and >57,000 HEV-related deaths, annually^{1,2}. In industrialized countries, hepatitis E was considered a rare disease until the discovery of the new HEV genotypes that turned this infection a concern of public health. In fact, four major HEV genotypes infect humans: genotypes 1 (HEV-1) and 2 (HEV-2), that are transmitted through the fecal-oral route via fecal contaminated water, being prevalent in areas of poor sanitation, such as in developing countries^{3,4}; and genotypes 3 (HEV-3) and 4 (HEV-4) that are zoonotic viruses transmitted to humans mainly through undercooked pork and boar meat and contact with pigs³⁻⁸, although, the transfusion of blood products has also been recently recognized as a risk factor⁹⁻¹¹. HEV-3 is today recognized as the main cause of sporadic autochthonous cases in industrialized countries^{3,4} and the numbers show that the incidence of reported cases has been increasing¹²⁻¹⁷.

In Europe, HEV infections are mainly caused by HEV-3, a genotype that causes asymptomatic infection or self-limiting acute hepatitis in healthy individuals, although it can lead to chronic infection with rapidly progressive cirrhosis in immunosuppressed patients, such as individuals with HIV, hematological malignancies, or transplant-related patients¹⁸⁻²¹. Solid organ transplant (SOT) recipients and hematopoietic stem cell transplant (HSCT) recipients are an important group at risk for HEV infection due to their prolonged immunosuppression state, that increases the risk for developing chronic infection¹⁹. Patients undergoing allogeneic-HSCT (allo-HSCT) have in general a higher risk for viral infections and higher incidence of graft-versus-host disease (GVHD) than autologous-HSCT (auto-HSCT)^{22,23}. In allo-HSCT recipients, progression to chronic infection may be favored by the severity of immunosuppression, which results in impaired immune reconstitution, including insufficient lymphocyte recovery, that are risk factors for post-transplantation infections²⁴⁻²⁶.

Moreover, HSCT recipients have a high transfusion burden, which is a problem particularly in countries that have not introduced HEV screening in blood donations^{22,27,28}.

The evidence that most of the HEV-3 infections become chronic in immunocompromised hosts, especially those after transplantation, makes these patients an important group of study^{29,30}. In SOT recipients, the reported acute HEV infection prevalence is 1-3%, with 47-83% of the patients developing chronic hepatitis³¹⁻³³, however, the prevalence and incidence of HEV in HSCT recipients is largely unknown.

OBJECTIVES

In the present study, we aim to summarize published data regarding HEV infections in HSCT recipients by performing a systematic review of the literature.

STUDY DESIGN

Literature search and study selection

Preferred Reporting of Systematic Reviews and MetaAnalyses (PRISMA) guidelines were followed in the preparation of this systematic review. Different queries, including MeSH terms, were tested and the literature search was performed with the query that obtained more representative manuscripts: ‘hepatitis E virus AND stem cell transplantation’. PubMed and Scopus databases were searched, independently by two of the authors (SC and CC), for published manuscripts on 31st January 2019 without restrictions on time period, sample size or population.

The eligibility criteria applied to studies were: 1) HEV infection (present or past) identified by the presence of HEV RNA and/or HEV-specific antibodies (IgM/IgG) in tested samples; 2) HSCT

recipients; 3) provide prevalence data. The exclusion criteria applied were: 1) duplicate data; 2) other types of manuscripts (reviews, case reports, comments or letters to the editor); 3) no access to abstracts and/or full texts; and 4) other languages rather than English, Spanish or Portuguese. Manuscript titles and abstracts were screened according to the eligibility criteria and selected manuscripts were fully reviewed for data extraction (author, publication date, country, population, age range, type of HSCT, HEV detection methods and the number of positive and negative cases). All manuscripts were reviewed independently by two of the authors (SC and CC) with disagreements mediated by the senior researcher (HS).

Statistical analysis

All data was inserted in a database that was used for prevalence analysis and comparison between studies. Prevalence analysis was performed using the *MetaXL* program version 5.3 (EpiGear International, Sunrise Beach, Queensland, Australia). The overall prevalence of HEV infection was estimated using the different approaches of diagnosis (HEV RNA or anti-HEV IgM/IgG detection) in HSCT recipients pooling the study data using the random effects model. The random effects model was used since a considerable heterogeneity among studies was expected, due largely to the different settings (populations, types of patients, age, gender, diagnostic methods) in which studies were conducted. The double arcsine transformation method was used for variance stabilization³⁴ considering a 95% confidence interval and a 5% statistical significance level ($p < 0.050$).

RESULTS

Study selection and description

The literature search retrieved a total of 73 manuscripts from both databases, and after duplicates removal, a total of 54 records were screened (Figure 1). After applying inclusion/exclusion criteria, 41

records were excluded: language (n=1), reviews (n=18), other types of articles such as Case Reports and Letter to the Editor (n=9), and studies not related to HSCT patients or HEV detection (n=13). A total of 13 full-text articles were assessed for full-reading of which 6 were excluded: 1 review, 4 case reports and 1 was not performed in HSCT patients. The bibliography of the selected manuscripts was reviewed to identify any new publications and no other article was added to the analysis.

After the full revision process, we included 7 manuscripts for data analysis^{18,21,35-39} (Table 1). Overall, these 7 studies included a total of 1178 HSCT patients from different countries: six studies were performed in Europe (United Kingdom, Netherlands, France, and Germany)^{18,21,36-39} and one study in China³⁵. These studies evaluated the prevalence of HEV infection based on the detection of HEV RNA (n=7) and/or the presence of anti-HEV IgM (n=3)/IgG (n=4). Phylogenetic analysis was performed in only three of these seven studies, revealing only HEV-3 genotype^{21,37,39}.

Here, we briefly resume the data from all included studies. In China, a study evaluated HEV infection in 177 haploidentical-HSCT recipients that presented unexplained elevated transaminases after transplant, identifying 7 patients with an acute/current HEV infection (2 positives for HEV RNA and 5 positives for anti-HEV IgM/IgG)³⁵. In the United Kingdom, two studies identified a total of 4 recipients with an acute HEV infection based in the presence of HEV RNA: in one study, 259 HSCT recipients (111 allo-HSCT, 145 auto-HSCT, and 3 CD34 top-up procedures) were evaluated and only 1 HEV RNA positive patient was identified³⁹; while the other study analyzed 144 allo-HSCT recipients and 3 patients were positive for HEV RNA²¹. In the Netherlands, two studies analysed HEV infection in HSCT: one study was performed in 130 allo-HSCT recipients with elevated alanine aminotransferase (ALT), identifying 5 HEV RNA positive patients³⁶; the other study analyzed 328 allo-HSCT recipients transplanted over a 5-year period, identifying a total of 10 patients with a current infection (8 were positive for HEV RNA and 2 were positive for anti-HEV IgM) and 41 patients were seropositive for anti-HEV IgG³⁷. In France, a study with 88 HSCT recipients (72 allo-HSCT and 16 auto-HSCT) revealed that none patient tested positive for HEV RNA, while 3 were positive for anti-

HEV IgM¹⁸. Interestingly, this study found a seroprevalence of anti-HEV IgG of 12,5% (11/88) and 36,4% (32/88) in the same group of HSCT recipients when using two different enzyme immunoassays (EIA) methods. In the study from Germany with 52 allo-HSCT recipients with elevated ALT, but without any specific cause of hepatitis, none of them tested positive for HEV RNA, while 3 were positive for anti-HEV IgG³⁸.

HEV infection prevalence analysis

We have performed an analysis of the prevalence of HEV infection in HSCT recipients according to the different approaches of diagnosis. The diagnosis of HEV positive cases is based on the identification of a patient with an HEV RNA positive sample (serum, plasma, blood or feces) or both anti-HEV IgM and IgG positive sample, nevertheless, the seroprevalence of the infection is referred to the detection of anti-HEV IgG only.

Considering the analysis of HEV RNA detection, a total of 1178 HSCT patients were studied and only 19 were positive, which gives an overall prevalence of 1.50% (95% CI: 0.70-2.60) with no significant difference observed between the different studies ($p=0.090$; Figure 2).

The analysis of anti-HEV IgM and IgG was performed separately: the detection of anti-HEV IgM as marker of acute infection was tested in a total of 593 HSCT patients with only 10 positive cases, giving an overall anti-HEV IgM prevalence of 2.00% (95% CI: 0.30-4.50; $p=0.060$; Figure 3); while the detection of anti-HEV IgG, a marker of past infection, was described in a total of 645 samples, 82 HSCT recipients were found positive to anti-HEV IgG giving an overall IgG seroprevalence of 11.4% (95% CI: 1.80-26.3), with statistically significant differences between the studies ($p<0.001$; Figure 4).

DISCUSSION

The evidence that most of the HEV-3 infections become chronic in immunocompromised hosts, especially those after transplantation, makes these patients an important group of study^{29,30}. Since the prevalence and incidence of HEV in HSCT recipients is largely unknown, we have performed a systematic review to understand the burden of HEV infection in these group of immunocompromised patients.

We found that there is a wide variation in HEV infection definition in literature, although in accordance with a recent surveillance report by the *European Centre for Disease Prevention and Control* (ECDC), concerning hepatitis E virus in Europe, a positive case is considered in a patient with an HEV RNA positive sample (serum, plasma, blood or feces) or both anti-HEV IgM and IgG positive sample¹². HEV RNA is detected usually between 2 and 8 weeks when viremia reaches its peak before strong declining, then, around 3 weeks after clinical symptoms, HEV RNA becomes undetectable, with the virus continuing to be shed in the stool for another 1 to 2 weeks. On the other hand, the anti-HEV IgM immune response remains detectable for 3–12 months and the IgG response reaches its peak four weeks later than IgM, remaining detectable for several years, although the exact duration of this response remains uncertain^{20,30}.

The fact that there is no standardized case definition, in addition to the heterogeneity in the analytical sensitivity of the commercial HEV assays, makes the comparison of studies more difficult. In this systematic review, all the seven studies performed the detection of HEV RNA, the most robust marker of acute/active infection^{18,21,35–39}, while four performed also the detection of anti-HEV IgM/IgG^{18,35,37,38}. The studies that performed HEV RNA detection as the diagnostic method for HEV infection in HSCT recipients revealed similar results between them, with an overall prevalence of HEV infection of 1.5%. Furthermore, we found that the overall prevalence of anti-HEV IgM was of 2.0%, a comparable value to the reported by the detection of HEV RNA, which is expected since both are markers of acute infection.

The analysis of anti-HEV IgG showed an overall seroprevalence of 11.4% in HSCT patients, with significant differences between the results of the different studies, mainly due to the study from France which revealed a much higher prevalence when compared to the others studies¹⁸. These results are difficult to compare due to the differences in the sensitivities and specificities of anti-HEV IgG commercial immunoassays⁴⁰⁻⁴³. Moreover, seroprevalence is greatly influenced by food habits^{5,43} and to differences in culinary practices between countries/regions⁴³. Reported anti-HEV IgG seroprevalence not only varies between countries but also within countries, with significant variance between regional areas⁴³. Indeed, even seroprevalences in Europe are very difficult to compare in consequence of these serology limitations and very different rates have been reported across the continent, ranging from 1.3-52%⁴⁴⁻⁴⁶. In Finland, anti-HEV IgG seroprevalence was found to be 27.6% in general population⁴⁷, while in Norway, Germany and Portugal reported seroprevalences were 11.4%, 16.8% and 16.3%, respectively⁴⁸⁻⁵⁰. Furthermore, several countries such as Netherlands, Spain, France, and Southwest England have performed these studies in healthy blood donors reporting an anti-HEV IgG prevalence ranging from 16.0 to 26.7%⁵¹⁻⁵⁴. These facts contribute to the assumption that the anti-HEV IgG prevalence does not reflect the spread of HEV in the HSCT population, which reinforces the importance of better characterization in this group of patients.

HSCT recipients are at higher risk of HEV infection, that could lead to chronic infection, and the diagnosis of HEV infection is highly recommended. Over the last years, some attention has been given to this group of immunocompromised patients, but there is still a small number of studies in HSCT as demonstrated in the present systematic review. Therefore, more studies are needed to increase our understanding of the epidemiology of HEV in HSCT recipients.

Conflict Of Interest

The authors declare they have no conflict of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions:

SC, acquisition of data, analysis and interpretation of data, drafting the article, final approval; CC, acquisition of data, final approval; CC and AT final approval; RM, conception and design of the study, final approval; MSJN, interpretation of data, revising the article, final approval; HS, analysis and interpretation of data, conception and design of the study, revising the article, final approval.

REFERENCES

1. World Health Organization. Hepatitis E. <https://www.who.int/news-room/fact-sheets/detail/hepatitis-e>. Published September 19, 2018. Accessed March 22, 2019.
2. World Health Organization. Hepatitis E | Immunization, Vaccines and Biologicals. WHO. <http://www.who.int/immunization/diseases/hepatitisE/en/>. Published October 19, 2015. Accessed March 22, 2019.
3. Kamar N, Bendall R, Legrand-Abravanel F, et al. Hepatitis E. *The Lancet*. 2012;379(9835):2477-2488. doi:10.1016/S0140-6736(11)61849-7
4. Meng X-J. Zoonotic and foodborne transmission of hepatitis E virus. *Semin Liver Dis*. 2013;33(1):41-49. doi:10.1055/s-0033-1338113
5. World Health Organization. The global prevalence of hepatitis E virus infection and susceptibility: A systematic review. 2010. <https://apps.who.int/iris/handle/10665/70513>. Accessed March 22, 2019.
6. Pavio N, Meng X-J, Renou C. Zoonotic hepatitis E: Animal reservoirs and emerging risks. *Vet Res*. 2010;41(6):46. doi:10.1051/vetres/2010018
7. Sonoda H, Abe M, Sugimoto T, et al. Prevalence of hepatitis E virus (HEV) Infection in wild boars and deer and genetic identification of a genotype 3 HEV from a boar in Japan. *J Clin Microbiol*. 2004;42(11):5371-5374. doi:10.1128/JCM.42.11.5371-5374.2004
8. Tei S, Kitajima N, Takahashi K, Mishiro S. Zoonotic transmission of hepatitis E virus from deer to human beings. *Lancet*. 2003;362(9381):371-373. doi:10.1016/S0140-6736(03)14025-1
9. Huzly D, Umhau M, Bettinger D, et al. Transfusion-transmitted hepatitis E in Germany, 2013. *Euro Surveill*. 2014;19(21).

10. Hauser L, Roque-Afonso A-M, Beylouné A, et al. Hepatitis E transmission by transfusion of Intercept blood system-treated plasma. *Blood*. 2014;123(5):796-797. doi:10.1182/blood-2013-09-524348
11. Hewitt PE, Ijaz S, Brailsford SR, et al. Hepatitis E virus in blood components: A prevalence and transmission study in southeast England. *Lancet*. 2014;384(9956):1766-1773. doi:10.1016/S0140-6736(14)61034-5
12. Aspinall EJ, Couturier E, Faber M, et al. Hepatitis E virus infection in Europe: Surveillance and descriptive epidemiology of confirmed cases, 2005 to 2015. *Eurosurveillance*. 2017;22(26):30561. doi:10.2807/1560-7917.ES.2017.22.26.30561
13. Koot H, Hogema BM, Koot M, Molier M, Zaaijer HL. Frequent hepatitis E in the Netherlands without traveling or immunosuppression. *Journal of Clinical Virology*. 2015;62:38-40. doi:10.1016/j.jcv.2014.11.020
14. Adlhoch C, Avellon A, Baylis SA, et al. Hepatitis E virus: Assessment of the epidemiological situation in humans in Europe, 2014/15. *Journal of Clinical Virology*. 2016;82:9-16. doi:10.1016/j.jcv.2016.06.010
15. Ijaz S, Vyse AJ, Morgan D, Pebody RG, Tedder RS, Brown D. Indigenous hepatitis E virus infection in England: More common than it seems. *Journal of Clinical Virology*. 2009;44(4):272-276. doi:10.1016/j.jcv.2009.01.005
16. Pischke S, Behrendt P, Bock C-T, Jilg W, Manns MP, Wedemeyer H. Hepatitis E in Germany – An under-reported infectious disease. *Dtsch Arztebl Int*. 2014;111(35-36):577-583. doi:10.3238/arztebl.2014.0577
17. Chalupa P, Vasickova P, Pavlik I, Holub M. Endemic Hepatitis E in the Czech Republic. *Clin Infect Dis*. 2014;58(4):509-516. doi:10.1093/cid/cit782
18. Abravanel F, Mansuy J-M, Huynh A, et al. Low risk of hepatitis E virus reactivation after

- haematopoietic stem cell transplantation. *J Clin Virol.* 2012;54(2):152-155. doi:10.1016/j.jcv.2012.02.015
19. Kamar N, Dalton HR, Abravanel F, Izopet J. Hepatitis E virus infection. *Clin Microbiol Rev.* 2014;27. doi:10.1128/CMR.00057-13
 20. Kamar N, Izopet J, Pavio N, et al. Hepatitis E virus infection. *Nat Rev Dis Primers.* 2017;3:17086. doi:10.1038/nrdp.2017.86
 21. Ankorn MJ, Ijaz S, Poh J, et al. Toward systematic screening for persistent hepatitis E virus infections in transplant patients. *Transplantation.* 2018;102(7):1139-1147. doi:10.1097/TP.0000000000002097
 22. Ljungman P, Bregni M, Brune M, et al. Allogeneic and autologous transplantation for haematological diseases, solid tumours and immune disorders: Current practice in Europe 2009. *Bone Marrow Transplant.* 2010;45(2):219-234. doi:10.1038/bmt.2009.141
 23. Galgano L, Hutt D. HSCT: How Does It Work? In: Kenyon M, Babic A, eds. *The European Blood and Marrow Transplantation Textbook for Nurses: Under the Auspices of EBMT.* Cham: Springer International Publishing; 2018:23-36. doi:10.1007/978-3-319-50026-3_2
 24. Storek J, Espino G, Dawson MA, Storer B, Flowers ME, Maloney DG. Low B-cell and monocyte counts on day 80 are associated with high infection rates between days 100 and 365 after allogeneic marrow transplantation. *Blood.* 2000;96(9):3290-3293.
 25. Wils E-J, van der Holt B, Broers AEC, et al. Insufficient recovery of thymopoiesis predicts for opportunistic infections in allogeneic hematopoietic stem cell transplant recipients. *Haematologica.* 2011;96(12):1846-1854. doi:10.3324/haematol.2011.047696
 26. Wils E-J, Cornelissen JJ. Thymopoiesis following allogeneic stem cell transplantation: New possibilities for improvement. *Blood Rev.* 2005;19(2):89-98. doi:10.1016/j.blre.2004.04.001
 27. Leahy MF, Trentino KM, May C, Swain SG, Chuah H, Farmer SL. Blood use in patients

receiving intensive chemotherapy for acute leukemia or hematopoietic stem cell transplantation: The impact of a health system-wide patient blood management program: IMPACT OF PBM PROGRAM IN HEMATOLOGY. *Transfusion*. 2017;57(9):2189-2196. doi:10.1111/trf.14191

28. Warner MA, Jambhekar NS, Saadeh S, et al. Implementation of a patient blood management program in hematopoietic stem cell transplantation: PBM IN STEM CELL TRANSPLANTATION. *Transfusion*. June 2019. doi:10.1111/trf.15414

29. Singh A, Seth R, Gupta A, et al. Chronic hepatitis E – an emerging disease in an immunocompromised host. *Gastroenterol Rep (Oxf)*. 2018;6(2):152-155. doi:10.1093/gastro/gow024

30. W. Webb G, Dalton HR. Hepatitis E: An underestimated emerging threat. *Therapeutic Advances in Infectious Disease*. 2019;6:204993611983716. doi:10.1177/2049936119837162

31. Pas SD, de Man RA, Mulders C, et al. Hepatitis E virus infection among solid organ transplant recipients, the Netherlands. *Emerging Infect Dis*. 2012;18(5):869-872. doi:10.3201/eid1805.111712

32. Kamar N, Selves J, Mansuy J-M, et al. Hepatitis E virus and chronic hepatitis in organ-transplant recipients. *N Engl J Med*. 2008;358(8):811-817. doi:10.1056/NEJMoa0706992

33. Haagsma EB, van den Berg AP, Porte RJ, et al. Chronic hepatitis E virus infection in liver transplant recipients. *Liver Transpl*. 2008;14(4):547-553. doi:10.1002/lt.21480

34. Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta-analysis of prevalence. *J Epidemiol Community Health*. 2013;67(11):974-978. doi:10.1136/jech-2013-203104

35. Tang F-F, Mo X-D, Wang Y, et al. Hepatitis E virus infection after haploidentical haematopoietic stem cell transplantation: Incidence and clinical course. *Br J Haematol*. 2019;184(5):788-796. doi:10.1111/bjh.15672

36. Willemse SB, Bezuur DL, Blom P, et al. Hepatitis E virus infection and hepatic GvHD in allogeneic hematopoietic stem cell transplantation recipients. *Bone Marrow Transplant*. 2017;52(4):622-624. doi:10.1038/bmt.2016.327

37. Versluis J, Pas SD, Agteresch HJ, et al. Hepatitis E virus: An underestimated opportunistic pathogen in recipients of allogeneic hematopoietic stem cell transplantation. *Blood*. 2013;122(6):1079-1086. doi:10.1182/blood-2013-03-492363
38. Koenecke C, Pischke S, Heim A, et al. Chronic hepatitis E in hematopoietic stem cell transplant patients in a low-endemic country? *Transpl Infect Dis*. 2012;14(1):103-106. doi:10.1111/j.1399-3062.2011.00702.x
39. Reekie I, Irish D, Ijaz S, et al. Hepatitis E infection in stem cell and solid organ transplant patients: A cross-sectional study: The importance of HEV RNA screening in peri-transplant period. *J Clin Virol*. 2018;107:1-5. doi:10.1016/j.jcv.2018.07.011
40. Bendall R, Ellis V, Ijaz S, Ali R, Dalton H. A comparison of two commercially available anti-HEV IgG kits and a re-evaluation of anti-HEV IgG seroprevalence data in developed countries. *J Med Virol*. 2010;82(5):799-805. doi:10.1002/jmv.21656
41. Pas SD, Streefkerk RHRA, Pronk M, et al. Diagnostic performance of selected commercial HEV IgM and IgG ELISAs for immunocompromised and immunocompetent patients. *J Clin Virol*. 2013;58(4):629-634. doi:10.1016/j.jcv.2013.10.010
42. Avellon A, Morago L, Garcia-Galera del Carmen M, Munoz M, Echevarría J-M. Comparative sensitivity of commercial tests for hepatitis E genotype 3 virus antibody detection. *J Med Virol*. 2015;87(11):1934-1939. doi:10.1002/jmv.24251
43. Hartl J, Otto B, Madden RG, et al. Hepatitis E Seroprevalence in Europe: A meta-analysis. *Viruses*. 2016;8(8). doi:10.3390/v8080211
44. Lapa D, Capobianchi MR, Garbuglia AR. Epidemiology of hepatitis E virus in european countries. *Int J Mol Sci*. 2015;16(10):25711-25743. doi:10.3390/ijms161025711
45. Scotto G, Martinelli D, Centra M, et al. Epidemiological and clinical features of HEV infection: A survey in the district of Foggia (Apulia, Southern Italy). *Epidemiol Infect*. 2014;142(2):287-294.

doi:10.1017/S0950268813001167

46. Mansuy J-M, Bendall R, Legrand-Abravanel F, et al. Hepatitis E virus antibodies in blood donors, France. *Emerging Infect Dis.* 2011;17(12):2309-2312. doi:10.3201/eid1712.110371
47. Kantala T, Maunula L, von Bonsdorff C-H, Peltomaa J, Lappalainen M. Hepatitis E virus in patients with unexplained hepatitis in Finland. *J Clin Virol.* 2009;45(2):109-113. doi:10.1016/j.jcv.2009.03.006
48. Olsøy IB, Henriksen S, Weissbach FH, et al. Seroprevalence of hepatitis E virus (HEV) in a general adult population in Northern Norway: The Tromsø study. *Med Microbiol Immunol.* March 2019. doi:10.1007/s00430-019-00599-5
49. Faber MS, Wenzel JJ, Jilg W, Thamm M, Höhle M, Stark K. Hepatitis E virus seroprevalence among adults, Germany. *Emerging Infect Dis.* 2012;18(10):1654-1657. doi:10.3201/eid1810.111756
50. Nascimento MSJ, Pereira SS, Teixeira J, et al. A nationwide serosurvey of hepatitis E virus antibodies in the general population of Portugal. *Eur J Public Health.* 2018;28(4):720-724. doi:10.1093/eurpub/ckx213
51. Slot E, Hogema BM, Riezebos-Brilman A, Kok TM, Molier M, Zaaier HL. Silent hepatitis E virus infection in Dutch blood donors, 2011 to 2012. *Euro Surveill.* 2013;18(31).
52. Sauleda S, Ong E, Bes M, et al. Seroprevalence of hepatitis E virus (HEV) and detection of HEV RNA with a transcription-mediated amplification assay in blood donors from Catalonia (Spain). *Transfusion.* 2015;55(5):972-979. doi:10.1111/trf.12929
53. Mansuy JM, Gallian P, Dimeglio C, et al. A nationwide survey of hepatitis E viral infection in French blood donors. *Hepatology.* 2016;63(4):1145-1154. doi:10.1002/hep.28436
54. Dalton HR, Stableforth W, Thuraiajah P, et al. Autochthonous hepatitis E in Southwest England: Natural history, complications and seasonal variation, and hepatitis E virus IgG seroprevalence in blood donors, the elderly and patients with chronic liver disease. *Eur J Gastroenterol*

LIST OF FIGURES

Figure 1 - PRISMA flow diagram.

Abbreviations: HEV, Hepatitis E Virus; HSCT, Hematopoietic stem cell transplantation




Figure 1 - PRISMA flow diagram applicable to the study.

Figure 2 - Forest plot of the overall prevalence based on the detection of HEV RNA.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.




Figure 2 - Forest plot of the overall prevalence based on the detection of HEV RNA.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.

Figure 3 - Forest plot of the overall prevalence based on the detection of IgM anti-HEV.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.




Figure 3 - Forest plot of the overall prevalence based on the detection of IgM anti-HEV.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.

Figure 4 - Forest plot of the overall prevalence based on the detection of IgG anti-HEV.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.




Figure 4 - Forest plot of the overall prevalence based on the detection of IgG anti-HEV.

Abbreviations: p, *p*-value; Prev (95%CI), Prevalence 95% Confidence Interval.

LIST OF TABLES

Table 1 - Details of the studies reporting HEV infection in HSCT recipients used in data analysis.

Abbreviations: EIA, Enzyme Immunoassay; HEV, Hepatitis E Virus; HSCT, Hematopoietic stem cell transplantation; IgG, Immunoglobulin G; IgM, Immunoglobulin M; n, sample size; Nested RT-PCR, Nested Reverse Transcription-Polymerase Chain Reaction; RT-qPCR, Reverse Transcription-quantitative Polymerase Chain Reaction.

Table 1 - Details of the studies reporting HEV infection in HSCT recipients used in data analysis.

First author (country, year)	Patients, n	Type of HSCT	HEV Diagnostic Methods	Assay	Positive, n	Prevalence
Tang FF et al. (China, 2019)	177	Allo-HSCT	Anti-HEV IgM Anti-HEV IgG	EIA (MP Diagnostics ®)	5	2.82%
			HEV RNA	Commercial RT-qPCR	2	1.13%
Reekie I, et al. (England, 2018)	259	Allo/Auto-HSCT	HEV RNA	In-house RT-qPCR	1	0.39%
Ankorn MJ, et al. (United Kingdom, 2018)	144	Allo-HSCT	HEV RNA	In-house RT-qPCR	3	2.08%
Willemse SB et al. (Netherlands, 2017)	130	Allo-HSCT	HEV RNA	Commercial RT-qPCR	5	3.85%
Verluis J, et al. (Netherlands, 2013)	328	Allo-HSCT	Anti-HEV IgM Anti-HEV IgG	EIA (Wantai ®)	2 41	0.61% 12.5%
			HEV RNA	In-house RT-qPCR	8	2.44%
Abravanel F, et al. (France, 2012)	88	Allo/Auto-HSCT	Anti-HEV IgM Anti-HEV IgG	EIA (Adaltis ®)	3 11	3.41% 12.5%
			Anti-HEV IgG	EIA (Wantai ®)	32	36.4%
			HEV RNA	In-house RT-qPCR	0	0.00%
Koenecke C, et al. (Germany, 2012)	52	Allo-HSCT	Anti-HEV IgG	EIA (Abbott ®)	3	5.77%
			HEV RNA	In-house Nested RT-PCR	0	0.00%

Abbreviations: EIA, Enzyme Immunoassay; HEV, Hepatitis E Virus; HSCT, Hematopoietic stem cell transplantation; IgG, Immunoglobulin G; IgM, Immunoglobulin M; n, sample size; Nested RT-PCR, Nested Reverse Transcription-Polymerase Chain Reaction; RT-qPCR, Reverse Transcription-quantitative Polymerase Chain Reaction.