

Seroprevalence of Hepatitis E Virus Infection in Pigs from Southern Bulgaria

Iliia Tsachev,¹ Magdalena Baymakova,² Massimo Ciccozzi,³ Roman Pepovich,⁴ Todor Kundurzhiev,⁵ Plamen Marutsov,¹ Kiril K. Dimitrov,⁶ Krasimira Gospodinova,¹ Maria Pishmisheva,⁷ and Liliya Pekova⁸

Abstract

Hepatitis E virus (HEV) has been isolated from humans and several animals' species. During the last years, the knowledge of HEV infection dramatically changed and enriched. The aim of this study was to estimate the seroprevalence of HEV in industrial pigs in different districts of Southern Bulgaria. Three hundred sixty swine serum samples were tested for anti-HEV IgG antibodies. The samples were collected from four industrial farms from three districts of Southern Bulgaria. HEV-specific antibodies in porcine serum were detected by enzyme-linked immunosorbent assay (PrioCHECK HEV Ab porcine). The overall HEV seroprevalence was 60.3%. The seropositivity varied widely depending on age groups and investigated farms. The overall prevalence in weaners was 25%, in fattening pigs 75.8%, and in group of sows was found the highest HEV positivity of 80%. The occurrence of HEV positivity in sows and fattening pigs presented odds ratio (OR) = 17.200 (95% confidence interval [CI]: 8.8–33.7) and OR = 11.342 (95% CI: 6.1–21.0), respectively, compared to weaners. The study indicated that HEV is widespread in industrial farms in Bulgaria and presented high seroprevalence in pigs. The results found that HEV seropositivity showed age dependency. The National Health Authorities should raise awareness of HEV and its zoonotic potential.

Keywords: hepatitis E virus, pigs, industrial farms

Introduction

HEPATITIS E VIRUS (HEV) HAS been isolated from humans and several animals' species. According to the World Health Organization (WHO) database, 20 million human HEV infections are estimated worldwide annually, leading to 3.3 million human cases with clinical presentation, resulting in ~44,000 deaths in 2015 (WHO 2018). HEV is classified into family Hepeviridae, divided in two genera: *Orthohepevirus* and *Piscihepevirus* (ICTV 2017). *Orthohepevirus* included four species: *Orthohepevirus A* (Genotypes: HEV-1–HEV-8), *Orthohepevirus B* (Avian HEV Genotype 2), *Orthohepevirus C* (Germany Rat HEV, Vietnam Rat HEV and Ferret HEV), and *Orthohepevirus D* (Germany Bat HEV) (ICTV 2017, Capai et al. 2018, Pepovich et al. 2019). Genus *Piscihepevirus* has only one species—*Piscihepevirus A*, and

one genotype—Cutthroat Trout HEV (ICTV 2017, Capai et al. 2018). HEV-3, HEV-4, and HEV-7 are zoonotic and infect both humans and animals. HEV-3 has been found in humans, swine, wild boar (ICTV 2017), wild deer (Wild deer HEV Subtype 3b, GenBank accession no. AB189071) (Tei et al. 2003), rabbit (Rabbit HEV Subtype 3ra, GenBank accession no. FJ906895) (Zhao et al. 2009), and mongoose (Mongoose HEV Subtype 3a, GenBank accession no. AB591734) (Nidaira et al. 2012). HEV-4 has been isolated from humans and swine (ICTV 2017). One report demonstrated a human chronic HEV infection caused by HEV-7 (Human HEV 7, GenBank accession no. KT818608), and also HEV stains were isolated from camels and were categorized into genotype 7 (DeHEV 7, GenBank accession no. KJ496143 and KJ496144) (Woo et al. 2014, Lee et al. 2016).

¹Department of Microbiology, Infectious and Parasitic Diseases, Faculty of Veterinary Medicine, Trakia University, Stara Zagora, Bulgaria.

²Department of Infectious Diseases, Military Medical Academy, Sofia, Bulgaria.

³Unit of Medical Statistics and Molecular Epidemiology, Università Campus Bio-Medico di Roma, Rome, Italy.

⁴Department of Infectious Pathology, Hygiene, Technology and Control of Foods from Animal Origin, Faculty of Veterinary Medicine, University of Forestry, Sofia, Bulgaria.

⁵Department of Occupational Medicine, Faculty of Public Health, Medical University, Sofia, Bulgaria.

⁶Department of General and Clinical Pathology, Faculty of Veterinary Medicine, Trakia University, Stara Zagora, Bulgaria.

⁷Department of Infectious Diseases, General Hospital, Pazardzhik, Bulgaria.

⁸Department of Infectious Diseases, Stara Zagora University Hospital, Stara Zagora, Bulgaria.

HEV infection in swine is asymptomatic; although histological liver changes were observed by autopsy (Halbur et al. 2001), the infection could be developed in all age groups: weaners, fattening pigs, and sows, and the main mechanism of HEV transmission among pigs is fecal-oral. Bouwknegt et al. (2008) reported that one HEV-infected pig could potentially infect other 8.8 (95% confidence interval [CI]: 4–19) pigs during the acute phase of infection.

In Bulgaria, the first evidence of human HEV infection was reported in 4 out of 53 (7.5%) cases with acute HEV infection in 1995 (Teoharov et al. 1995). Investigations of human HEV infections in Bulgaria have been started since 2014 (Teoharov et al. 2014, Baymakova et al. 2016, Bruni et al. 2018, Cella et al. 2019), whereas first preliminary data for swine HEV infection in Bulgaria were published in 2018 (Pishmisheva et al. 2018). That report presented an overall seroprevalence of anti-HEV antibodies in swine of 40% (34 positive samples out of all 85 sera) in both investigated districts Pazardzhik and Sliven (Pishmisheva et al. 2018). Up to date, no data are available for HEV prevalence among pigs in other areas of the country.

The aim of this study was to estimate the seroprevalence of HEV in industrial pigs in different districts of Southern Bulgaria.

Materials and Methods

Areas and pigs collection

Three hundred sixty pigs ($n = 360$) were enrolled from four industrial farms (Kalchevo, Yambol, Pishigovo, and Kos-

tinbrod) in three districts of Southern Bulgaria—Yambol, Pazardzhik, and Sofia (Fig. 1). Animals were divided in three age groups: weaners, fattening pigs, and sows. An equal number of samples ($n = 30$) were taken from each age group on each farm. Consequently, a total of 90 samples were tested from each farm. Pigs showed no clinical signs at sampling time point; unfortunately, sex had not been recorded. The pigs' selection in farms, respectively, in age groups was done randomly. The investigated farms were located among plains ($\sim 42^{\circ}48'N$ and $42^{\circ}69'N$ Latitude and $23^{\circ}32'E$ and $26^{\circ}51'E$ Longitude), and the climate is subtropical/continental (average annual temperature $13\text{--}15^{\circ}C$ and precipitations about 650 mm/m^2).

Sampling

Swine blood samples (up to 5 mL per individual) were taken by puncture of the *sinus ophthalmicus*. Blood collection tubes without anticoagulant were kept at room temperature ($20^{\circ}C$) until visible clot retraction, centrifuged at $1500 \times g$ for 10 minutes, and the serum was separated and kept at $-20^{\circ}C$ until processing.

HEV serological assay

The serum samples were tested for HEV antibodies in Laboratory of Infectious Diseases, Faculty of Veterinary Medicine, Trakia University, Stara Zagora, Bulgaria. A commercial enzyme-linked immunosorbent assay (PrioCHECK HEV Ab porcine; Mikrogen GmbH, Neuried, Germany) was used. The



FIG. 1. Geographic distribution of HEV infection in pigs from Southern Bulgaria. HEV, hepatitis E virus.

PrioCHECK HEV Ab porcine is a diagnostic test for detection of HEV-specific antibodies in porcine serum and meat juice samples. Microtiter plate is coated with recombinant HEV antigen of the open reading frame 2 (ORF2) and ORF3 of genotypes HEV-1 and HEV-3. The test has 91.0% sensitivity and 94.1% specificity. The cutoff value was calculated according to the manufacturer's instructions. The cutoff is calculated as mean optical density 450 (OD₄₅₀) of the cutoff control multiplied with 1.2. Results obtained above or equal to the cutoff are considered positive. Results between the OD₄₅₀ of the cutoff control and the cutoff are doubtful. Results obtained below the cut-off are negative.

Ethics statement

The study was approved by the Ethics Committee in Animal Experimentation and Animal Welfare at Trakia University, Stara Zagora (Bulgaria), and was conducted according to the ethical principles of animal experimentation, adopted by the Bulgarian Ministry of Agriculture, Food and Forestry.

Statistical analysis

HEV-positive results among different swine age groups and farms were compared by the chi-squared test. Binary logistic regression was used to evaluate the risk of positive results according to age group. Statistical analysis has been performed by Excel 2007 (Microsoft, Redmond, WA) and SPSS Statistics 19.0 (IBM Corp., Armonk, NY). A *p*-value <0.05 was considered statistically significant.

Results

Positive results for anti-HEV IgG were detected in 217 (60.3%) of all 360 tested sera (Table 1). The overall seropositivity in weaners, fattening pigs, and sows was 25% (mean \pm SD: 25.0 \pm 30.9; 95% CI: 5.2–55.2), 75.8% (mean \pm SD: 75.8 \pm 36.2; 95% CI: 40.3–111.4), and 80% (mean \pm SD: 80.0 \pm 17.6; 95% CI: 62.7–97.3), respectively. The highest HEV seropositivity in age group weaners was found in Kostinbrod (21/30, 70%). The lowest HEV seropositivity in

age group fattening pigs was estimated in Yambol (7/30, 23.3%). In age group of sows, the highest HEV seropositivity was presented in Kostinbrod (29/30, 96.6%). The overall prevalence of anti-HEV antibodies in each farm was as follows: in Kalchevo—58.8% (53/90), in Yambol—43.3% (39/90), in Pishtigovo—50% (45/90), and in Kostinbrod—88.8% (80/90). Based on age groups and different farms, the chi-squared test showed significant differences in HEV seropositivity between the age groups and farms (Table 1). Five percent of all samples were assessed as doubtful.

To estimate the risk for HEV seropositivity, the odds ratio (OR) in different age groups was performed by binary logistic regression. The OR of anti-HEV antibody occurrence in fattening pigs and sows was determined by comparing to group weaners (Table 2). We found that the odds of HEV infection was nearly 11 times higher in fattening pigs and 17 times higher in sows than in the group of weaners.

Discussion

Moderate and high HEV seropositivity among swine were observed in different countries from Southeastern Europe. The estimated anti-HEV positivity in different districts in Romania was 42.7% (65/145) and 49.3% (34/69) (Savuta et al. 2007, 2008), Serbia—34.6% (109/315) (Lupulovic et al. 2010), Greece—80% (76/96) (Siochu et al. 2009), and Croatia—32.9% (469/1424) (Jemersic et al. 2017). In West-European countries of the Mediterranean Basin, the prevalence of swine HEV infection was lower compared to the Balkan Peninsula. In Spain was observed a positivity of 25% (15/60) and 20.4% (233/1141) in different settings and years (Pina et al. 2000, Jimenez de Oya et al. 2011), France—31% (2035/6565) (Rose et al. 2011), and Italy—50.2% (714/1422) (Martinelli et al. 2011). Countries from Western Europe reported relatively high swine HEV seroprevalence. In Switzerland was found 58.1% HEV seropositivity (1161/1999; 95% CI: 55.9–60.2) (Burri et al. 2014), Germany—49.8% and 68.6% (Baechlein et al. 2010, Wacheck et al. 2012), Netherlands—72% in conventional farms (191/265) (Rutjes et al. 2014), Belgium—73% (95% CI: 68.8–77.5) (Thiry et al.

TABLE 1. SEROPREVALENCE OF HEPATITIS E VIRUS INFECTION BY AGE GROUPS IN PIGS FROM SOUTHERN BULGARIA

Age groups	Investigated pigs, n	HEV positive, n (%)	Chi-squared	df	p
Kalchevo					
Weaners	30	0 (0.0)	72.688	2	<0.001
Fattening pigs	30	30 (100.0)			
Sows	30	23 (76.7)			
Yambol					
Weaners	30	5 (16.7)	38.823	2	<0.001
Fattening pigs	30	7 (23.3)			
Sows	30	27 (90.0)			
Pishtigovo					
Weaners	30	4 (13.3)	30.276	2	<0.001
Fattening pigs	30	24 (80.0)			
Sows	30	17 (56.7)			
Kostinbrod					
Weaners	30	21 (70.0)	16.425	2	<0.001
Fattening pigs	30	30 (100.0)			
Sows	30	29 (96.7)			

df, degrees of freedom; HEV, hepatitis E virus.

TABLE 2. LOGISTIC REGRESSION SHOWING THE RELATIONSHIP BETWEEN HEPATITIS E VIRUS-POSITIVE PIGS AND AGE GROUP

Age groups	Investigated pigs, n	HEV positive, n (%)	PE	SE	p	OR	95% CI
Weaners	120	30 (25.0)	NA	NA	NA	1.000	NA
Fattening pigs	120	91 (75.8)	2.429	0.315	<0.001	11.342	6.1–21.0
Sows	120	96 (80.0)	2.845	0.343	<0.001	17.200	8.8–33.7

CI, confidence interval; NA, not applicable; OR, odds ratio; PE, parameter estimate; SE, standard error.

2014), England—85.5% (219/256) and 92.8% (584/629) (Banks et al. 2004, Grierson et al. 2015), and Scotland—61.4% (108/176) (Crossan et al. 2015). Scandinavian countries also had a high prevalence of swine HEV. In Sweden the seropositivity was assessed as 58% (Banks et al. 2004), Denmark—73.2% (Breum et al. 2010), Finland—86.3% (Kantala 2017), and Norway—90% (137/153) (Lange et al. 2017).

In North America, the swine HEV seroprevalence varied widely. In Canada, the reported seropositivity was 18.1% (129/712) in one study and another found 59.4% (594/998) (Meng et al. 1999, Yoo et al. 2001). In USA, the situation was similar and depended on studies—34.5% (29/84) and 68.9% (202/293) (Meng et al. 1997, Withers et al. 2002). The estimated anti-HEV positivity in Mexico was 59.4% and 80% (Cooper et al. 2005, Garcia-Hernandez et al. 2017).

In Asia, the prevalence of HEV antibodies in pigs varied widely. In China, the reported seroprevalence was 78.8% (330/419) and 78.9% (490/621) (Wang et al. 2002, Li et al. 2011). In contrast, Thailand reported 30.7% (23/75) serum samples positive for swine HEV (Meng et al. 1999). Others Asian countries published data in moderate to high range as South Korea—40.7% (57/140) (Meng et al. 1999), Indonesia—72% (71/99) (Wibawa et al. 2004), Taiwan—37.1% (102/275) (Hsieh et al. 1999), and India—42.9% (122/284) (Arankalle et al. 2002).

As shown, the prevalence of HEV antibodies was quite variable between countries, also in different areas in the same countries. This variability could be influenced by study design, diagnostics methods, and tested population. Nevertheless, it seems that pigs could be infected with HEV and be a zoonotic risk for transmission.

The seroprevalence in our study (60.3%) is similar to data found in Switzerland (58.1%) (Burri et al. 2014), Scotland (61.4%) (Crossan et al. 2015), Sweden (58%) (Banks et al. 2004), Canada (59.4%) (Yoo et al. 2001), and Mexico (59.4%) (Garcia-Hernandez et al. 2017). Sows had the highest seroprevalence (80%) in our study. Martinelli et al. (2011) found 70.6% HEV prevalence in sows, also the highest seroprevalence in their study, whereas Danish investigation presented 73.2% positive sows for HEV-IgG (Breum et al. 2010). The results of our study showed an age-dependent seroprevalence (OR = 11.342, $p < 0.001$; OR = 17.200, $p < 0.001$). A similar relationship was observed in another report (Breum et al. 2010). Consequently, HEV occurs in all age groups and pigs could be infected in nursery and fattening periods most likely, corresponding to natural HEV infection (Pavio et al. 2010). Most pigs become infected in 6–8 weeks, virus in feces has peak in 12–14 weeks and decline until 20–22 weeks (Pavio et al. 2010). HEV-IgG appears in 8–9 weeks, increase, and approximately all infected pigs in 22–24 weeks could have HEV-IgG (Pavio et al.

2010). Moreover, Bouwknegt et al. (2009) presented evidence of HEV RNA in urine. They found positive sample in urine of a contact-infected pig at 65 days after the beginning of virus fecal shedding (Bouwknegt et al. 2009). The study highlighted the urine as a potential source for HEV infection among pigs and supposed that contact pathway could be better for transmission of HEV than intravenous inoculation (Bouwknegt et al. 2009). Probably the pig farm density, the husbandry conditions, model of pig farming, animal contact with the environment, sewage system, and water facility could be some reasons for high HEV seroprevalence in pigs. Jemersic et al. (2017) found the highest HEV prevalence in regions with the highest density of pigs and wild boars. Garcia-Hernandez et al. associated the high seroprevalence with short production cycle of fattening and high pig population in farms (Garcia-Hernandez et al. 2017). In our study, the included farms were typical industrial farms with high density of animals, shorter farming for fattening, many herds, and close contact among pigs. These factors probably contributed to the high HEV seroprevalence in pigs. Further projects should be planned to evaluate swine HEV seroprevalence in whole country, molecular analysis for sequences, and human HEV seroprevalence in blood donors.

Conclusions

In conclusion, the preliminary results presented in this study showed a high prevalence of HEV antibodies in pigs in Southern Bulgaria. The evidence for HEV infection in pigs highlights the concerns for zoonotic transmission. National Health Organizations should enhance the control in farms and slaughterhouses, obligatorily record human HEV infection, promote risk factors for transmission in risk population groups, and improve the knowledge about infection among physicians, veterinarians, and general practitioners. Further researches could be realized to a thorough understanding of situation of HEV infection in Bulgaria.

Acknowledgment

This study was funded by the Trakia University, 6000 Stara Zagora, Bulgaria (grant no. 01/2017).

Authors' Contributions

I.T. and R.P.—study design, data collection, data interpretation, article preparation, literature search, and funds collection; M.B. and M.C.—study design, statistical analysis, data interpretation, article preparation, and literature search; T.K.—statistical analysis; P.M. and K.K.D.—data collection and funds collection; K.G.—laboratory analysis; and M.P. and L.P.—funds collection. All authors read and approved the final version of the article.

Author Disclosure Statement

No conflicting financial interests exist.

References

- Arankalle VA, Chobe LP, Joshi MV, Chadha MS, et al. Human and swine hepatitis E viruses from Western India belong to different genotypes. *J Hepatol* 2002; 36:417–425.
- Baechlein C, Schielke A, John R, Ulrich RG, et al. Prevalence of Hepatitis E virus-specific antibodies in sera of German domestic pigs estimated by using different assays. *Vet Microbiol* 2010; 144:187–191.
- Banks M, Heath GS, Grierson SS, King DP, et al. Evidence for the presence of hepatitis E virus in pigs in the United Kingdom. *Vet Rec* 2004; 154:223–227.
- Baymakova M, Sakem B, Plochev K, Popov GT, et al. Epidemiological characteristics and clinical manifestations of hepatitis E virus infection in Bulgaria: A report on 20 patients. *Srp Arh Celok Lek* 2016; 144:63–68.
- Bouwknegt M, Frankena K, Rutjes SA, Wellenberg GJ, et al. Estimation of hepatitis E virus transmission among pigs due to contact-exposure. *Vet Res* 2008; 39:40.
- Bouwknegt M, Rutjes SA, Reusken CB, Stockhofe-Zurwieden N, et al. The course of hepatitis E virus infection in pigs after contact-infection and intravenous inoculation. *BMC Vet Res* 2009; 5:7.
- Breum SO, Hjulsgaard CK, de Deus N, Segales J, et al. Hepatitis E virus is highly prevalent in the Danish pig population. *Vet Microbiol* 2010; 146:144–149.
- Bruni R, Villano U, Equestre M, Chionne P, et al. Hepatitis E virus genotypes and subgenotypes causing acute hepatitis, Bulgaria, 2013–2015. *PLoS One* 2018; 13:e0198045.
- Burri C, Vial F, Ryser-Degiorgis MP, Schwermer H, et al. Seroprevalence of hepatitis E virus in domestic pigs and wild boars in Switzerland. *Zoonoses Public Health* 2014; 61:537–544.
- Capai L, Charrel R, Falchi A. Hepatitis E in high-income countries: What do we know? And what are the knowledge gaps? *Viruses* 2018; 10:E285.
- Cella E, Golkocheva-Markova E, Sagnelli C, Scolamacchia V, et al. Human hepatitis E virus circulation in Bulgaria: Deep Bayesian phylogenetic analysis for viral spread control in the country. *J Med Virol* 2019; 91:132–138.
- Cooper K, Huang FF, Batista L, Rayo CD, et al. Identification of genotype 3 hepatitis E virus (HEV) in serum and fecal samples from pigs in Thailand and Mexico, where genotype 1 and 2 HEV strains are prevalent in the respective human populations. *J Clin Microbiol* 2005; 43:1684–1688.
- Crossan C, Grierson S, Thomson J, Ward A, et al. Prevalence of hepatitis E virus in slaughter-age pigs in Scotland. *Epidemiol Infect* 2015; 143:2237–2240.
- Garcia-Hernandez ME, Cruz-Rivera M, Sanchez-Betancourt JJ, Rico-Chavez O, et al. Seroprevalence of anti-hepatitis E virus antibodies in domestic pigs in Mexico. *BMC Vet Res* 2017; 13:289.
- Grierson S, Heaney J, Cheney T, Morgan D, et al. Prevalence of Hepatitis E virus infection in pigs at the time of slaughter, United Kingdom, 2013. *Emerg Infect Dis* 2015; 21:1396–1401.
- Halbur PG, Kasornrondkub C, Gilbert C, Guenette D, et al. Comparative pathogenesis of infection of pigs with hepatitis E viruses recovered from a pig and a human. *J Clin Microbiol* 2001; 39:918–923.
- Hsieh SY, Meng XJ, Wu YH, Liu ST, et al. Identity of a novel swine hepatitis E virus in Taiwan forming a monophyletic group with Taiwan isolates of human hepatitis E virus. *J Clin Microbiol* 1999; 37:3828–3834.
- International Committee on Taxonomy of Viruses (ICTV). *Hepeviridae* (November 2017). Available at <https://talk.ictvonline.org>
- Jemersic L, Keros T, Maltar L, Barbic L, et al. Differences in hepatitis E virus (HEV) presence in naturally infected seropositive domestic pigs and wild boars—an indication of wild boars having an important role in HEV epidemiology. *Vet Arhiv* 2017; 87:651–663.
- Jimenez de Oya N, de Blas I, Blazquez AB, Martin-Acebes MA, et al. Widespread distribution of hepatitis E virus in Spanish pig herds. *BMC Res Notes* 2011; 4:412.
- Kantala T. Presence of hepatitis E virus (HEV) and markers for HEV infection in production swine, human patients with unexplained hepatitis, and veterinarians in Finland. PhD Thesis, University of Helsinki (Faculty of Veterinary Medicine), Helsinki, Finland, April 28, 2017.
- Lange H, Overbo J, Borgen K, Dudman S, et al. Hepatitis E in Norway: Seroprevalence in humans and swine. *Epidemiol Infect* 2017; 145:181–186.
- Lee GH, Tan BH, Teo EC, Lim SG, et al. Chronic infection with camelid hepatitis E virus in a liver transplant recipient who regularly consumes camel meat and milk. *Gastroenterology* 2016; 150:355–357.e3.
- Li W, Shu X, Pu Y, Bi J, et al. Seroprevalence and molecular detection of hepatitis E virus in Yunnan Province, China. *Arch Virol* 2011; 156:1989–1995.
- Lupulovic D, Lazic S, Prodanov-Radulovic J, Jimenez de Oya N, et al. First serological study of hepatitis E virus infection in backyard pigs from Serbia. *Food Environ Virol* 2010; 2:110–113.
- Martinelli N, Luppi A, Cordioli P, Lombardi G, et al. Prevalence of hepatitis E virus antibodies in pigs in Northern Italy. *Infect Ecol Epidemiol* 2011; 1: DOI: 10.3402/iee.v1i0.7331.
- Meng XJ, Dea S, Engle RE, Friendship R, et al. Prevalence of antibodies to the hepatitis E virus in pigs from countries where hepatitis E is common or is rare in the human population. *J Med Virol* 1999; 59:297–302.
- Meng XJ, Purcell RH, Halbur PG, Lehman JR, et al. A novel virus in swine is closely related to the human hepatitis E virus. *Proc Natl Acad Sci USA* 1997; 94:9860–9865.
- Nidaira M, Takahashi K, Ogura G, Taira K, et al. Detection and phylogenetic analysis of hepatitis E viruses from mongooses in Okinawa, Japan. *J Vet Med Sci* 2012; 74:1665–1668.
- Pavio N, Meng XJ, Renou C. Zoonotic hepatitis E: Animal reservoirs and emerging risks. *Vet Res* 2010; 41:46.
- Pepovich R, Baymakova M, Pishmisheva M, Marutsov P, et al. Current knowledge on *Hepatitis E virus* infection. *Vojnosanit Pregl* 2019; 76: [Epub ahead of Print]. DOI: 10.2298/VSP170815159P.
- Pina S, Buti M, Cotrina M, Piella J, et al. HEV identified in serum from humans with acute hepatitis and in sewage of animal origin in Spain. *J Hepatol* 2000; 33:826–833.
- Pishmisheva M, Baymakova M, Golkocheva-Markova E, Kundurzhiev T, et al. First serological study of hepatitis E virus infection in pigs in Bulgaria. *C R Acad Bulg Sci* 2018; 71:1001–1008.
- Rose N, Lunazzi A, Dorenlor V, Merbah T, et al. High prevalence of Hepatitis E virus in French domestic pigs. *Comp Immunol Microbiol Infect Dis* 2011; 34:419–427.
- Rutjes SA, Bouwknegt M, van der Giessen JW, de Roda Husman AM, et al. Seroprevalence of hepatitis E virus in pigs from different farming systems in The Netherlands. *J Food Prot* 2014; 77:640–642.

- Savuta G, Anita A, Anita D, Ludu L, et al. Preliminary epidemiological investigations regarding hepatitis E virus infection in swine from the North-East of Romania. *Bull USAMV-CN* 2007; 64:356–358.
- Savuta GH, Anita A, Anita D, Ludu L, et al. Seroepidemiological researches regarding swine and human hepatitis E in Romania. *Lucrari Stiintifice Med Vet (Timisoara)* 2008; 41:309–313.
- Siochu A, Tzika E, Alexopoulos C, Kyriakis SC, et al. First report of serological evidence of hepatitis E virus infection in swine in Northern Greece. *Acta Vet (Beograd)* 2009; 59:205–211.
- Tei S, Kitajima N, Takahashi K, Mishiho S. Zoonotic transmission of hepatitis E virus from deer to human beings. *Lancet* 2003; 362:371–373.
- Teoharov P, Kevorkyan A, Raycheva R, Golkocheva-Markova E, et al. Data on the prevalence of hepatitis E virus in Bulgaria. *C R Acad Bulg Sci* 2014; 67:1427–1432.
- Teoharov P, Tiholova M, Draganov P, Lilyanova V, et al. First cases of hepatitis E virus infection in Bulgaria [in Bulgarian]. *Infectology (Sofia)* 1995; 32:17–18.
- Thiry D, Mauroy A, Saegerman C, Thomas I, et al. Estimation of hepatitis E virus (HEV) pig seroprevalence using ELISA and Western blot and comparison between human and pig HEV sequences in Belgium. *Vet Microbiol* 2014; 172:407–414.
- Wacheck S, Werres C, Mohn U, Dorn S, et al. Detection of IgM and IgG against hepatitis E virus in serum and meat juice samples from pigs at slaughter in Bavaria, Germany. *Food-borne Pathog Dis* 2012; 9:655–660.
- Wang YC, Zhang HY, Xia NS, Peng G, et al. Prevalence, isolation, and partial sequence analysis of hepatitis E virus from domestic animals in China. *J Med Virol* 2002; 67:516–521.
- Wibawa ID, Muljono DH, Mulyanto, Suryadarma IG, et al. Prevalence of antibodies to hepatitis E virus among apparently healthy humans and pigs in Bali, Indonesia: Identification of a pig infected with a genotype 4 hepatitis E virus. *J Med Virol* 2004; 73:38–44.
- Withers MR, Correa MT, Morrow M, Stebbins ME, et al. Antibody levels to hepatitis E virus in North Carolina swine workers, non-swine workers, swine, and murids. *Am J Trop Med Hyg* 2002; 66:384–388.
- Woo PC, Lau SK, Teng JL, Tsang AK, et al. New hepatitis E virus genotype in camels, the Middle East. *Emerg Infect Dis* 2014; 20:1044–1048.
- World Health Organization (WHO). Hepatitis E: Key facts (19 September 2018). Available at www.who.int
- Yoo D, Willson P, Pei Y, Hayes MA, et al. Prevalence of hepatitis E virus antibodies in Canadian swine herds and identification of a novel variant of swine hepatitis E virus. *Clin Diagn Lab Immunol* 2001; 8:1213–1219.
- Zhao C, Ma Z, Harrison TJ, Feng R, et al. A novel genotype of hepatitis E virus prevalent among farmed rabbits in China. *J Med Virol* 2009; 81:1371–1379.

Address correspondence to:

*Magdalena Baymakova
Department of Infectious Diseases
Military Medical Academy
Sofia 1606
Bulgaria*

E-mail: dr.baymakova@gmail.com