
barchart(g10 ~ x | g2, subset = g10 != "1")
0.4
0.6
0.8
1
2
10
9
8
7
6
5
4
3
2
0.4
0.6
0.8
x


barchart(unclass(Titanic))
0
200
400
600
800
Adult
Male
Adult
Female
Crew
3rd
2nd
1st
Child
Male
Child
Female
Crew
3rd
2nd
1st
0
200
400
600
800
Freq


barchart(g2)
2
1
0
1
2
3
4
5
Freq


barchart(VADeaths)
70−74
65−69
60−64
55−59
50−54
0
50
100
150
200
250
Freq


barchart(x)
0.4
0.6
0.8
x


barchart(UCBAdmissions)
0
500
1000
1500
Male
Female
Rejected
Admitted
0
500
1000
1500
Freq


bwplot(g2 ~ x + y, outer = TRUE)
0.0
0.2
0.4
0.6
0.8
x
y
2
1
0.0
0.2
0.4
0.6
0.8
x + y


bwplot(y, notch = TRUE)
0.0
0.2
0.4
0.6
0.8
y


densityplot(~x, groups = g2)
Density
0.0
0.5
1.0
1.5
x
0.0
0.5
1.0
1.5


densityplot(y, plot.points = "jitter")
Density
1.0
0.5
0.0
0.0
0.5
1.0
y


dotplot(g10 ~ x | g2)
0.4
0.6
0.8
1
2
10
9
8
7
6
5
4
3
2
1
0.4
0.6
0.8
x


dotplot(unclass(Titanic))
0
200
400
600
Adult
Male
Adult
Female
Child
Male
Child
Female
Crew
3rd
2nd
1st
0
200
400
600
Freq


dotplot(g2)
2
1
4.6
4.8
5.0
5.2
5.4
Freq


dotplot(VADeaths)
70−74
65−69
60−64
55−59
50−54
20
40
60
Freq


dotplot(x)
0.4
0.6
0.8
x


dotplot(UCBAdmissions)
Male
Female
Rejected
Admitted
0
100
200
300
400
500
Freq


histogram(~c(x, y))
Percent of Total
0
10
20
30
0.0
0.2
0.4
0.6
0.8
1.0
c(x, y)


histogram(g2)
Percent of Total
50
40
30
20
10
0
1
2
g2


histogram(c(x, y))
Percent of Total
0
10
20
30
0.0
0.2
0.4
0.6
0.8
1.0
c(x, y)


qqmath(~x + y)
x + y
0.0
0.2
0.4
0.6
0.8
−1
0
1
qnorm


qqmath(x)
x
0.8
0.6
0.4
−1
0
1
qnorm


stripplot(g2 ~ x + y, outer = TRUE)
0.0
0.2
0.4
0.6
0.8
x
y
2
1
0.0
0.2
0.4
0.6
0.8
x + y


stripplot(y, jitter = TRUE)
0.0
0.2
0.4
0.6
0.8
y


qq(g2 ~ x)
0.8
0.6
0.4
2
0.4
0.6
0.8
1


xyplot(y ~ x)
0.8
0.6
0.4
0.2
0.0
y
0.4
0.6
0.8
x


xyplot(ts(x))
0.8
0.6
0.4
2
4
6
8
10
Time


levelplot(y ~ g2 + g10)
g10
10
9
8
7
6
5
4
3
2
1
g2
1
2
0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0


levelplot(UCBAdmissions)
Admitted
Rejected
D
E
F
A
B
C
Female
Male
Gender
Female
Male
Admitted
Rejected
Admitted
Rejected
Admit
500
400
300
200
100
0


levelplot(as.table(x), ...)
1st
2nd
3rd
Crew
Yes
Child
Yes
Adult
No
Child
No
Adult
Female
Male
Sex
Female
Male
1st
2nd
3rd
Crew
Class
700
600
500
400
300
200
100
0


levelplot(VADeaths)
Urban Female
Urban Male
Rural Female
Rural Male
column
50−54
55−59
60−64
65−69
70−74
row
70
60
50
40
30
20
10


contourplot(y ~ g2 + g10)
10
9
8
7
6
5
4
3
2
1
g10
1
2
g2


contourplot(UCBAdmissions)
Admitted
Rejected
D
E
F
A
B
C
Female
Male
Gender
Admitted
Rejected
Admit


contourplot(unclass(Titanic))
1st
2nd
3rd
Crew
Yes
Child
Yes
Adult
No
Child
No
Adult
Female
Male
Sex
Female
Male
1st
2nd
3rd
Crew
Class


contourplot(VADeaths)
Urban Female
Urban Male
Rural Female
Rural Male
column
20
30
40
50
50−54
55−59
60−64
65−69
70−74
row


cloud(g10 ~ x + y)
g10
y
x


cloud(VADeaths)
VADeaths
column
row

cloud(UCBAdmissions)



wireframe(y ~ g2 + g10)




wireframe(VADeaths)
VADeaths
column
row


splom(~cbind(x = x, y = y, g = as.numeric(g2)))
g
y
x
Scatter Plot Matrix


splom(cbind(x = x, y = y, g = as.numeric(g2)))
x
y
g
Scatter Plot Matrix


splom(data.frame(x, y, g2))
g2
y
x
Scatter Plot Matrix


parallelplot(~iris)
Species
Petal.Width
Petal.Length
Sepal.Width
Sepal.Length
Min
Max


parallelplot(data.matrix(iris[1:4]))
Petal.Width
Petal.Length
Sepal.Width
Sepal.Length
Min
Max


parallelplot(iris)
Species
Petal.Width
Petal.Length
Sepal.Width
Sepal.Length
Min
Max


rfs(oneway(y ~ g2))
Fitted Values minus Mean
Residuals
0.2
0.4
0.6
0.8
0.2
0.0
−0.2
−0.4
0.2
0.4
0.6
0.8
f−value


tmd(sort(y) ~ sort(x))
0.00
−0.05
−0.10
−0.15
−0.20
−0.25
difference
0.2
0.4
0.6
0.8
mean


tmd(xyplot(sort(y) ~ sort(x)))
0.00
−0.05
−0.10
−0.15
−0.20
−0.25
difference
0.2
0.4
0.6
0.8
mean


xyplot(y ~ x, pch = 16, cex = 1.5)
0.8
0.6
0.4
0.2
0.0
y
0.4
0.6
0.8
x


xyplot(y ~ x | g2, data = g, cex = c(1, 2))
1
2
0.4
0.6
0.8
0.0
0.2
0.4
0.6
0.8
y
x


densityplot(g$x, plot.points = FALSE)
Density
1.0
0.5
0.0
0.0
0.5
1.0
g$x


plot(equal.count(rnorm(1000)))
Panel
6
5
4
3
2
1
−2
0
2
4
Range