

## Overcoming energetic barriers in acetogenic C1 conversion

Alexander Katsyv<sup>1</sup>, Volker Müller<sup>1\*</sup>

<sup>1</sup>Molecular Microbiology & Bioenergetics, Goethe University Frankfurt, Germany

*Submitted to Journal:*

Frontiers in Bioengineering and Biotechnology

*Specialty Section:*

Synthetic Biology

*ISSN:*

2296-4185

*Article type:*

Review Article

*Received on:*

25 Oct 2020

*Accepted on:*

19 Nov 2020

*Provisional PDF published on:*

19 Nov 2020

*Frontiers website link:*

[www.frontiersin.org](http://www.frontiersin.org)

*Citation:*

Katsyv A and Müller V(2020) Overcoming energetic barriers in acetogenic C1 conversion. *Front. Bioeng. Biotechnol.* 8:1420. doi:10.3389/fbioe.2020.621166

*Copyright statement:*

© 2020 Katsyv and Müller. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution and reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

This Provisional PDF corresponds to the article as it appeared upon acceptance, after peer-review. Fully formatted PDF and full text (HTML) versions will be made available soon.

# Overcoming energetic barriers in acetogenic C1 conversion

Alexander Katsyv and Volker Müller

Department of Molecular Microbiology & Bioenergetics, Institute of Molecular Biosciences, Johann Wolfgang Goethe University, Frankfurt am Main, Germany

<sup>#</sup>correspondence address: Prof. Volker Müller, Department of Molecular Microbiology & Bioenergetics, Institute of Molecular Biosciences, Johann Wolfgang Goethe University, Frankfurt am Main, Germany; Phone: 49-6979829507; Fax: 49-69-79829306;  
E-mail: [vmueller@bio.uni-frankfurt.de](mailto:vmueller@bio.uni-frankfurt.de)

## Abstract

Currently one of the biggest challenges for society is to combat global warming. A solution to this global threat is the implementation of a CO<sub>2</sub>-based bioeconomy and a H<sub>2</sub>-based bioenergy economy. Anaerobic lithotrophic bacteria such as the acetogenic bacteria are key players in the global carbon and H<sub>2</sub> cycle and thus prime candidates as driving forces in a H<sub>2</sub>- and CO<sub>2</sub>-bioeconomy. Naturally, they convert two molecules of CO<sub>2</sub> *via* the Wood-Ljungdahl pathway (WLP) to one molecule of acetyl-CoA which can be converted to different C<sub>2</sub>-products (acetate or ethanol) or elongated to C<sub>4</sub> (butyrate) or C<sub>5</sub>-products (caproate). Since there is no net ATP generation from acetate formation, an electron-transport phosphorylation (ETP) module is hooked up to the WLP. ETP provides the cell with additional ATP, but the ATP gain is very low, only a fraction of an ATP per mol of acetate. Since acetogens live at the thermodynamic edge of life, metabolic engineering to obtain high-value products is currently limited by the low energy status of the cells that allows for the production of only a few compounds with rather low specificity. To set the stage for acetogens as production platforms for a wide range of bioproducts from CO<sub>2</sub>, the energetic barriers have to be overcome. This review summarizes the pathway, the energetics of the pathway and describes ways to overcome energetic barriers in acetogenic C1 conversion.

**Keywords:** carbon capture, biohydrogen, H<sub>2</sub> storage, biofuels, electron transport

## Introduction

In times of global warming there is an immediate need to reduce green house gas emissions. CO<sub>2</sub> is by far the most important based on atmospheric concentrations and there are several chemical as well as biological approaches to reduce the atmospheric CO<sub>2</sub> concentration (Claassen, 1999; Dey, 2007; Anwar et al., 2018). One is to reduce CO<sub>2</sub> emissions at the first place followed by efficient capture and storage of CO<sub>2</sub> (Benson, 2008; Leung, 2014). Lithotrophic microbes that make their biomass from CO<sub>2</sub> are prime candidates to solve both problems (Götz, 2016; Liew et al., 2016; Dürre, 2017; Heijstra et al., 2017). Traditionally, biotechnological processes that compete with oil-based processes use sugars to produce high-value end products (Naik et al., 2010). These approaches do not only produce CO<sub>2</sub> but also compete for sugar that is a feedstock for humans (Ajanovic, 2011). In contrast, lithotrophic organisms do not produce but use CO<sub>2</sub> as feedstock (Bengelsdorf et al., 2018). Lithotrophs fix CO<sub>2</sub> for biomass production with energy derived from solar energy (photolithoautotrophs) or from the oxidation of an inorganic electron donor such as H<sub>2</sub> (chemolithoautotrophs). The latter occurs in the dark and the absence of oxygen, and thus operates at low costs (Rittmann and Herwig, 2012). Lithotrophic, H<sub>2</sub>-oxidizing, CO<sub>2</sub> fixing microorganisms are the methanogenic archaea that only produce methane (Wolfe, 1971; Zehnder and Brock, 1979; Enzmann et al., 2018) or the acetogenic bacteria that produce different end products such as acetate and ethanol or butyrate and formate (Müller et al., 2004; Drake et al., 2008). They are also metabolically flexible and grow lithotrophically on H<sub>2</sub> + CO<sub>2</sub> or on CO, but also heterotrophically on sugars, alcohols, carbonic acids, primary and secondary alcohols (Andreesen et al., 1973; Bache and Pfennig, 1981; Eichler and Schink, 1984; Drake et al., 1997; Drake et al., 2008; Ragsdale and Pierce, 2008; Schuchmann and Müller, 2016). Heterotrophic growth in almost every case goes along with reduction of CO<sub>2</sub> to acetate (Müller, 2003). Thus, acetogens can couple oxidation of various organic and inorganic electron donors to the reduction of CO<sub>2</sub> to acetate or the other before mentioned products.

Therefore, acetogens are the most flexible organisms to be used for a biological approach to capture and store CO<sub>2</sub> in the dark and absence of O<sub>2</sub> (Daniell et al., 2012; Liew et al., 2016; Köpke and Simpson, 2020). Since CO<sub>2</sub> fixation can be driven by H<sub>2</sub> oxidation, these bacteria also capture and store H<sub>2</sub>, a key process in the biohydrogen economy (Bailera, 2017; Müller, 2019; Schwarz and Müller, 2020).

### **Acetogenic bacteria and the Wood-Ljungdahl pathway**

Acetogenic bacteria are a phylogenetically very diverse group of strictly anaerobic bacteria ubiquitous in nature. They are characterized by a reductive pathway in which two mol of CO<sub>2</sub> are reduced to one mol of acetyl-CoA and then further to acetate, the Wood-Ljungdahl pathway (WLP) (Figure 1) (Ljungdahl, 1986; Wood et al., 1986). Reducing power can be derived from the oxidation of organic but also inorganic carbon sources. Some acetogens naturally produce ethanol and butyrate in addition and, therefore, they have come into focus of an alternative, CO<sub>2</sub>-based bioeconomy (Schiel-Bengelsdorf and Dürre, 2012; Dürre, 2017). Ethanol is also produced from CO<sub>2</sub>, H<sub>2</sub> and CO (syngas) on a large industrial scale (Daniell et al., 2012; Bengelsdorf et al., 2018). Apart from these gaseous substrates used, other C1 substrates such as formic acid or methanol are promising feedstocks for an alternative bioeconomy using acetogens as biocatalysts (Cotton et al., 2019; Müller, 2019). In addition to the mere use as production platforms, acetogens are also promising candidates in the H<sub>2</sub>-economy as potential catalysts for H<sub>2</sub> storage or production (Bailera, 2017; Schwarz et al., 2018; Schwarz and Müller, 2020).

Among the naturally occurring carbon fixation pathways the WLP is the only pathway that does not need additional ATP to operate (Ragsdale and Pierce, 2008; Fuchs, 2011). One mol of CO<sub>2</sub> is reduced to formate and then bound to the C1-carrier tetrahydrofolate (THF), driven by the hydrolysis of ATP (Himes and Harmony, 1973; Lovell et al., 1988). From the formyl-THF, water is split off and the produced methenyl-THF is further reduced *via* methylene- to

72 methyl-THF (Clark and Ljungdahl, 1982). The latter condenses with CO and coenzyme A on  
73 the enzyme acetyl-CoA synthase/CO dehydrogenase to acetyl-CoA; the CO derives from the  
74 reduction of another CO<sub>2</sub> by the CO dehydrogenase (Drake et al., 1980; Pezacka and Wood,  
75 1984; Ragsdale and Wood, 1985; Raybuck et al., 1988; Seravalli et al., 1997). Acetyl-CoA is  
76 then converted *via* acetyl phosphate to acetate and ATP. Thus the net synthesis of ATP by  
77 substrate level phosphorylation is zero. Since the bacteria grow by production of acetate from  
78 H<sub>2</sub> + CO<sub>2</sub> there must be additional mechanisms to generate ATP (Schaupp and Ljungdahl,  
79 1974; Drake et al., 1981; Müller, 2003).

### 81 Chemiosmotic energy conservation in acetogens

82 Currently, acetogens can be divided into two bioenergetic groups, the Rnf- and the Ech-  
83 acetogens (Figure 2) (Schuchmann and Müller, 2014). The Rnf- and the Ech-complex are  
84 membrane-bound respiratory enzymes that both use reduced ferredoxin as reductant (Schmehl  
85 et al., 1993; Hedderich and Forzi, 2005; Buckel and Thauer, 2018b). The Rnf complex has six  
86 subunits and catalyzes ferredoxin:NAD<sup>+</sup> oxidoreductase activity; the free energy change of  
87 the electron transport is coupled to the extrusion of ions from the cytoplasm to the periplasm  
88 thus generating an electrochemical ion gradient across the membrane that drives ATP  
89 synthesis *via* a F<sub>1</sub>F<sub>0</sub> ATP synthase (Müller et al., 2008; Biegel and Müller, 2010; Westphal et  
90 al., 2018). The coupling ion maybe Na<sup>+</sup> as in the case of *A. woodii* or H<sup>+</sup> as suggested for  
91 *Clostridium ljungdahlii* or *Clostridium autoethanogenum* (Biegel and Müller, 2010; Tremblay  
92 et al., 2012; Kuhns et al., 2020). Two points are important to make: first, the nature of the  
93 coupling ion used has dramatic consequences for the energetic status, especially under stress  
94 conditions and second, NADH is the end product of this respiration and can be regarded as a  
95 waste product that needs to be recycled (reoxidized). Thus, by metabolic engineering, any  
96 reductive, NADH consuming pathway can be hooked up to the Rnf complex. In the second  
97 group, the Ech-acetogens, with *Thermoanaerobacter kivui* and *Moorella thermoacetica* as

model strains, reduced ferredoxin is oxidized by an 8- or 9-subunit, membrane-bound enzyme that transfers the electron to protons, thus producing molecular  $H_2$  (Pierce et al., 2008; Schölmerich and Müller, 2019). This electron transport is coupled to the export of ions across the cytoplasmic membrane and the electrochemical ion gradient established drives the synthesis of ATP *via* a  $F_1F_0$  ATP synthase (Welte et al., 2010; Schoelmerich and Müller, 2019). Three points are important to make here: first, the nature of the ferredoxin involved and its redox potential is not known. *A. woodii*, for example, has seven genes potentially encoding a ferredoxin. Since the  $CO_2/CO$  couple ( $E_0' [CO_2/CO] = -520$  mV) requires a low potential reductant we calculate with a redox potential of -450 to -500 mV. Second, the potential difference between reduced ferredoxin ( $E_0' [Fd^{2-}/Fd] = \sim -450$  to -500 mV) and protons ( $E_0' [H_2/H^+] = -414$  mV) is only about half ( $\Delta E_0' = +86$  mV to +36 mV) of the difference of the redox couple reduced ferredoxin: $NAD^+$  ( $\Delta E_0' = +180$  mV) and thus, only about half the amount of ions can be translocated per electron transported. These numbers have not been determined for any Rnf- or Ech-complex but based on thermodynamics we consider two ions per two electrons in Rnf and one ion per two electrons in Ech as maximum. Third, the product of respiration is the volatile gas  $H_2$  and thus there is no need for a reductive pathway to be hooked up to the respiration since the electron escapes into the environment as gas. Although this is attractive for the cells it can be deleterious to the cell if the  $H_2$  partial pressure in the ecosystem is so high that it exhibits a thermodynamic backup pressure on the enzyme so that respiration comes to an end. The  $\Delta G^{0'}$  of the reaction ferredoxin: $H^+$  is only -7 kJ/mol to -16.6 kJ/mol. This allows  $H_2$  production only up to a partial pressure of 0.05 MPa. The function of Rnf as respiratory enzyme has been proven by genetic (Westphal et al., 2018) and biochemical experiments (Biegel and Müller, 2010) and recently, the final proof has been obtained using the enzyme from a thermophilic fermenting bacterium, *Thermotoga maritima* (not an acetogen) (Kuhns et al., 2020) as well as the acetogen *A. woodii* (Wiechmann and Müller, unpublished data). The enzyme was purified and reconstituted into liposomes and

catalyzed electron transfer from reduced ferredoxin to  $\text{NAD}^+$ ; this electron transport was coupled to a primary and electrogenic  $\text{Na}^+$  transport into the lumen of the liposomes (Wiechmann and Müller, unpublished data). For Ech, there is evidence that inverted membrane vesicles of the acetogen *T. kivui* couple  $\text{H}^+$  as well as  $\text{Na}^+$  transport into the lumen of the vesicles to  $\text{H}_2$  production from reduced ferredoxin (Schölmerich and Müller, 2019). However, this organism has two Ech-encoding gene clusters and the final proof of ion translocation and the ion used has to await genetic studies and its purification and reconstitution into liposomes. Ion transport coupled to Ech-catalyzed reaction has also been observed in vesicles of the methanogenic archaeon *Methanosarcina mazei* (Welte et al., 2010) and there is no reason to believe that Ech is not as respiratory enzyme in acetogens. This is different in cytochrome-containing acetogens. *A. woodii* and *T. kivui* do not contain cytochromes, but some acetogens like *M. thermoacetica* do. Historically, the discovery of cytochromes in the acetogens *Clostridium formicoaceticum* and *Moorella thermoacetica* was an exciting discovery since it immediately argued for a respiratory chain involved in acetogenesis (Gottwald et al., 1975). Unfortunately, 45 years after their discovery there is no evidence for that. Indeed, it may well be that cytochromes are not involved in acetogenesis ( $\text{CO}_2$  reduction) but reduction of alternative electron acceptors such as nitrate. Since the involvement of cytochromes in acetogenesis is still a molecular hallucination, we will not discuss this further in this review.

The ion gradient established by the Rnf and Ech complex is then used by a  $\text{F}_1\text{F}_0$ -ATP-synthase to drive the synthesis of ATP.  $\text{F}_1\text{F}_0$ -ATP-synthases have been purified from only a few acetogens such as *M. thermoacetica* (Ivey and Ljungdahl, 1986), *Moorella thermoautotrophica* (Das et al., 1997), *A. woodii* (Reidlinger and Müller, 1994) and *Eubacterium limosum* (Litty and Müller, 2020).  $\text{F}_1\text{F}_0$ -ATP-synthases are macromolecular machines that convert electrochemical energy via mechanical energy into chemical energy (ATP) (Müller and Grüber, 2003). They are composed of two motors that are connected by a

central stalk. Ion flow through the membrane-embedded motor made of the rotor (a ring of  $c$  subunits) and the stator ( $a$  subunit) drives rotation of the rotor against the stator. Furthermore, rotation of the  $c$ -ring drives rotation of the central stalk that interacts with the three ATP synthesizing centers in the hydrophilic  $F_1$  domain, leads to the synthesis of ATP from ADP +  $P_i$ . The  $c$  subunit harbours the ion binding-site which are protons in most ATP synthases (Mayer et al., 1986; Meier et al., 2006; Hess et al., 2014) and  $Na^+$  in a few (Meier et al., 2009; Brandt and Muller, 2015; Mayer et al., 2015). Protons are bound to the so-called active carboxylate in helix two of the  $c$  subunit (a aspartate or glutamate residue) (Fillingame et al., 2003) whereas the conserved  $Na^+$ -binding site has in addition two residues, a glutamine in helix one and a serine or threonine residue downstream of the active carboxylate (Rahlf and Müller, 1999; Müller and Grüber, 2003). There are more residues involved in complexing  $Na^+$  but these three are the conserved ones that make the conserved  $Na^+$ -binding motif. *C. ljungdahlii* and *C. autoethanogenum* only have the active carboxylate but not the  $Na^+$ -binding motif (Mayer et al., 1986), whereas *A. woodii* (Reidlinger et al., 1994; Matthies et al., 2014) as well as *E. limosum* (Litty and Müller, 2020) have the conserved  $Na^+$ -binding motif. The activity of the purified enzymes is strictly  $Na^+$  dependent and the enzymes translocate  $Na^+$  after reconstitution into liposomes (Fritz and Müller, 2007; Litty and Müller, 2020).

A critical number for bioenergetic calculations is the number of  $c$  subunits per ring because this gives the number of ions translocated for the synthesis of three moles of ATP. This number has been experimentally determined only for *A. woodii*; in this case the  $c$  ring has 10  $Na^+$ -binding sites (Matthies et al., 2014). Divided by three ATP synthesizing centers that gives 3.3  $Na^+$ /ATP. The non-acetogenic *Clostridium paradoxum* has 11  $H^+$ -binding sites, giving a stoichiometry of 3.6  $H^+$ /ATP (Ferguson et al., 2006; Meier et al., 2006). This number is used in this review to calculate the overall ATP yields in *C. autoethanogenum*.



**Electron carriers in the WLP**

After having discussed that acetogens have either one of the two respiratory enzymes Rnf or Ech the obvious question is: what are the cofactors of the other dehydrogenases/reductases in the WLP and how is the pool of reduced ferredoxin maintained. The first enzyme in the carbonyl branch is the acetyl-CoA synthase/CO dehydrogenase which is also the key enzyme in the entire pathway (Drake et al., 1980; Pezacka and Wood, 1984; Ragsdale and Wood, 1985; Raybuck et al., 1988; Seravalli et al., 1997). Due to the low redox potential of the CO<sub>2</sub>/CO couple ( $E_0' [\text{CO}_2/\text{CO}] = -520 \text{ mV}$ ) only reduced ferredoxin can act as reductant here. How ferredoxin ( $E_0' [\text{Fd}^{2-}/\text{Fd}] = \sim -450 \text{ to } -500 \text{ mV}$ ) is reduced with H<sub>2</sub> ( $E_0' [\text{H}_2/\text{H}^+] = -414 \text{ mV}$ ) has been an enigma for a long time. With the discovery of electron bifurcation to couple endergonic with exergonic redox reactions (Li et al., 2008; Buckel and Thauer, 2018a), the solution was at hand: an electron-bifurcating hydrogenase (Schut and Adams, 2009). Such an enzyme is indeed also present in *A. woodii*. It couples the exergonic electron flow from H<sub>2</sub> to NAD<sup>+</sup> ( $E_0' [\text{NAD}^+/\text{NADH}] = -320 \text{ mV}$ ) to the endergonic electron flow from H<sub>2</sub> to ferredoxin (Schuchmann and Müller, 2012). In this context it is important to note that electron bifurcation is often mistakenly considered as mechanism for energy conservation. It is not, but it saves cellular energy. If electron transport from H<sub>2</sub> to ferredoxin would be driven by ATP, at least one ATP would have to be invested. With electron bifurcation the equivalent of only a fraction of an ATP can be invested. This kind of energy saving is a prerequisite for life at the thermodynamic limit as in acetogens. Therefore, acetogens employ a multitude of different electron-bifurcating enzymes in their metabolism and, actually most of the various electron-bifurcating enzymes known to date are from acetogens (Müller et al., 2018).

The first step in the methyl branch is the reduction of CO<sub>2</sub> to formate. Since the redox potential of the CO<sub>2</sub>/formate couple is -420 mV, neither NAD<sup>+</sup> nor NADP<sup>+</sup> can be used as reductant. The solution is different in different acetogens: some use a ferredoxin-dependent formate dehydrogenase, others a combination of an electron-bifurcating hydrogenase and an

electron-bifurcating formate dehydrogenase and still others use a  $H_2$ -dependent  $CO_2$  reductase (HDCR) (Yamamoto et al., 1983; Nagarajan et al., 2013; Schuchmann and Müller, 2013; Wang et al., 2013a; Wang et al., 2013b). In the latter,  $H_2$  is oxidized by a hydrogenase subunit of the HDCR and the electrons are transferred, most likely by two small iron-sulfur containing electron transfer subunits, to the formate dehydrogenase (Schuchmann and Müller, 2013). This is energetically feasible given the redox potential  $E^{0'}$  of -420 mV of the redox couple  $CO_2$ /formate and -414 mV of the  $H_2/2H^+$  couple. In the electron-bifurcating formate dehydrogenase of *C. autoethanogenum*, the same is achieved but by two consecutive reactions: first, a  $NADP^+$ -specific electron-bifurcating hydrogenase reduces  $NADP^+$  and ferredoxin and second, an electron-bifurcating formate dehydrogenase uses  $NADP^+$  and reduced ferredoxin for  $CO_2$  reduction (Wang et al., 2013a). The next reduction step is the reduction of methenyl-THF to methylene-THF, catalyzed by the methylene-THF dehydrogenase. This enzyme maybe  $NAD^+$  specific as in *A. woodii* or  $NADP^+$  specific as in *C. autoethanogenum* or *Sporomusa ovata* (O'Brien et al., 1973; Ragsdale and Ljungdahl, 1984; Wang et al., 2013a; Kremp et al., 2020). The last reduction step is the reduction of methylene-THF to methyl-THF, catalyzed by the methylene-THF reductase (MTHFR). This enzyme is of special interest in the bioenergetics of acetogens and it was proposed 43 years ago by Thauer *et al.* to be an energetic coupling site (Thauer et al., 1977). This is based on the high redox potential of the pair methylene-/methyl-THF of -200 mV, arguing that its reduction with NADH would deliver sufficient energy to be used in energy conservation (Wohlfarth and Diekert, 1991). Accordingly, it was speculated for a long time that the methylene-THF reductase is the terminal acceptor of a respiratory chain that energizes the cytoplasmic membrane for ATP synthesis (Müller, 2003). However, with the genomic sequences for acetogens and the biochemical data available this can be ruled out. Nevertheless, the enzyme could use electron bifurcation (Buckel and Thauer, 2018a; Müller et al., 2018) to couple the exergonic reduction of methylene- to methyl-THF with the endergonic

reduction of ferredoxin which then drives energy conservation in the respiratory chain. Indeed, the enzymes from *M. thermoacetica* or *C. ljungdahlii* do not couple NADH oxidation to methylene-THF reduction, indicating that a second electron acceptor is missing (Moore et al., 1974; Clark and Ljungdahl, 1984; Mock et al., 2014). Since it is a characteristic of electron-bifurcating enzymes that they are only active in presence of all three reaction partners, the lack of activity was taken to suggest that the enzyme is electron bifurcating to an unknown second acceptor (Mock et al., 2014). However, this still needs to be verified. In contrast, *A. woodii* has a  $\text{NAD}^+$ -dependent, non-electron-bifurcating methylene-THF reductase (Bertsch et al., 2015). In sum, the electron carriers used by the enzymes of the WLP are very different.

### **Electron carriers involved in substrate oxidation**

The same holds true for the electron carriers involved in substrate oxidation. Sugars such as glucose, fructose, mannitol are oxidized by the Embden-Meyerhof-Parnas pathway with  $\text{NAD}^+$  in the glycerol-3-phosphate dehydrogenase reaction and ferredoxin in the pyruvate:ferredoxin oxidoreductase reaction as electron carriers. Other substrates such as ethanol only yield NADH and still others such as formate only yield molecular  $\text{H}_2$  (Schuchmann and Müller, 2013; Bertsch et al., 2016). During lithotrophic growth,  $\text{H}_2$  is oxidized by an electron-bifurcating hydrogenase yielding both, NADH and reduced ferredoxin (Schuchmann and Müller, 2012). In Ech-containing acetogens,  $\text{H}_2$  oxidation could theoretically yield reduced ferredoxin by Ni-Fe hydrogenases (Ech-complex), as shown in the methanogenic archaeon *Methanosarcina barkeri* (Meuer et al., 2002). CO is oxidized by the CO dehydrogenase coupled to reduction of ferredoxin (Seravalli et al., 1997). In sum, oxidation of different substrates yields different reduced electron carries such as NADH, NADPH, reduced ferredoxin,  $\text{H}_2$  or combinations thereof, whereas the WLP only accepts,

species specific, distinct electron donors in the right stoichiometry. Thus, there is an essential need to balance the electrons between the oxidation module and the WLP.


### **Redox balancing: many variations of a theme**

Conversion of redox carriers into each other requires energy and since acetogens live at the thermodynamic edge of life, they have evolved energy-efficient ways. They employ specialized membrane-bound and soluble enzyme systems. The membrane-bound enzymes are the before discussed respiratory enzymes Rnf and Ech. They are both coupled reversibly to the membrane potential. In the above described function they serve to reduce  $\text{NAD}^+$  ( $E_0' [\text{NAD}^+/\text{NADH}] = -320 \text{ mV}$ ) with reduced ferredoxin  $E_0' [\text{Fd}^{2-}/\text{Fd}] = \sim -450 \text{ to } -500 \text{ mV}$ ) as reductant or to reduce protons to  $\text{H}_2$  ( $E_0' [\text{H}_2/\text{H}^+] = -414 \text{ mV}$ ). During growth on low energy substrates that only yield NADH, *A. woodii* employs the Rnf to drive reverse electron transport (Hess et al., 2013b; Westphal et al., 2018). This is actually the function in most organisms and this function was decisive for the name giving: *Rhodobacter nitrogen fixation* (Schmehl et al., 1993). There, the Rnf complex drives the endergonic reduction of ferredoxin, required for nitrogen fixation, with NADH as reductant (Schmehl et al., 1993; Kumagai et al., 1997; Saeki and Kumagai, 1998). Analogously, the Ech complex can drive the endergonic reduction of ferredoxin with  $\text{H}_2$  as reductant (Meuer et al., 2002). The same function is achieved by the soluble electron-bifurcating hydrogenase that couples  $\text{H}_2$  oxidation to the reduction of both,  $\text{NAD}^+$  (exergonic) and ferredoxin (endergonic) (Schuchmann and Müller, 2012). Conversion of  $\text{NADP}^+$  and  $\text{NAD}^+$  is catalyzed with reduced ferredoxin as driver in the Nfn and Stn-type electron-bifurcating transhydrogenases found in acetogens and the electron-bifurcating hydrogenase/formate dehydrogenase complex connects cellular  $\text{H}_2$ ,  $\text{NADP}^+$ , ferredoxin and  $\text{CO}_2$  pools (Wang et al., 2010; Nguyen et al., 2017; Kremp et al., 2020). Together with the difference in electron carrier specificity of the substrate oxidation reaction and the WLP and the different respiratory enzymes this gives thousands of possible

combinations for energy conservation (Figure 3). Since acetogens growing on  $H_2 + CO_2$  make only a fraction of an ATP per mol of acetate formed, a change in the electron carrier specificity of a given metabolic scheme makes a huge difference. Therefore, reliable calculation for the overall ATP gain can only be done with clear conscience for those organisms in which the electron carrier specificity of the redox reaction and the type of respiratory enzyme has been determined experimentally. Metabolic schemes with too many unknown variables are rather harmful than useful. Therefore, we will concentrate on the well-studied acetogen *A. woodii* that grows very robust on  $H_2 + CO_2$  and is a prime candidate also for a biohydrogen economy, but that does not grow on CO or syngas, and on *C. autoethanogenum* (or its close relative, *C. ljungdahlii*), a working horse in the industrial production of biofuels from syngas but a less well understood metabolism.

### Bioenergetics of *A. woodii* and *C. autoethanogenum*

Acetogenesis from  $H_2 + CO_2$  or CO according to equations 1 and 2 goes along with a free energy change of -95 and -175 kJ/mol.



Considering the  $H_2$  and CO concentration in the environment this is sufficient for only 0.3 and 1.5 mol ATP/mol acetate, respectively (Kim and Hegeman, 1983; Novelli et al., 1999; Schuchmann and Müller, 2014). The enzymology and bioenergetics of *A. woodii* and *C. autoethanogenum* for acetogenesis from  $H_2 + CO_2$  or CO is summarized in Figure 4 and 5. In *A. woodii*, not only the electron carriers are known but also the ion/ATP stoichiometry of the ATP synthase is known, the only case for acetogens (Matthies et al., 2014). This allows to calculate the ATP/acetate stoichiometry very accurately with 0.3 mol ATP/mol of acetate with  $H_2 + CO_2$  as substrate (Figure 4A). *A. woodii* does not grow on CO (Bertsch and Müller,

2015) but resting cells are able to oxidize CO according to equation 2 (Diekert et al., 1986). This yields 1.5 ATP/mol of acetate (Figure 4B). For *C. autoethanogenum*, there are two unknowns: the ion/ATP stoichiometry and whether or not the methylene-THF reductase (MTHFR) is electron bifurcating. For the calculations we have assumed an ion/ATP stoichiometry of 3.6 and for the methylene-THF reductase we assume either electron bifurcation with ferredoxin as second electron acceptor or no electron bifurcation. With these two variables, the ATP gain in *C. autoethanogenum* varies from 0.4 - 1 mol/mol of acetate with  $H_2 + CO_2$  as substrate (Figure 5A and C). It should be mentioned that the ATP yield of acetogenesis from  $H_2 + CO_2$  in *C. autoethanogenum* is higher compared to *A. woodii* if an electron-bifurcating methylene-THF reductase is assumed (Figure 5A and B). With CO as electron donor, the energetic of acetogenesis is much better. According to equation 2, the  $\Delta G^{0'}$  is -175 kJ/mol and acetate formation from CO in *C. autoethanogenum* goes along with the synthesis of 1 - 1.5 mol ATP/mol acetate (Figure 5B and D). This is more than from  $H_2 + CO_2$  but both values are still very low compared to fermenting bacteria or even aerobes (Müller, 2008). However, the situation gets worse when products other than acetate are formed from acetyl-CoA (Figure 6).

### **Bioenergetics of the formation of products other than acetate in *A. woodii* and *C. autoethanogenum***

Acetyl-CoA acts as a precursor, not only for the production of acetate, but also for the formation of products like ethanol (Köpke et al., 2011a; Basen et al., 2014; Müller, 2014; Lo et al., 2015), butanol (Dürre et al., 1992), lactate (Gladden, 2004), 2,3 butanediol (Syu, 2001; Köpke et al., 2011b; Hess et al., 2015), acetone (Dürre et al., 1992; Hoffmeister et al., 2016), isobutene (van Leeuwen et al., 2012) or isoprene (Diner et al., 2018) (Figure 6). As deduced above, the reduction of  $CO_2$  to acetyl-CoA with  $H_2$  in *A. woodii* requires 0.7 ATP (Figure 4A), whereas with CO as donor, 0.5 ATP is produced (Figure 4B). *C. autoethanogenum* has

an energy demand of 0.6 ATP when reducing  $\text{CO}_2$  to acetyl-CoA with  $\text{H}_2$  (Figure 5C), whereas with CO as donor, the ATP balance is zero (Figure 5D). Therefore, if the further conversion of acetyl-CoA to the desired product gains ATP, the production from CO will be possible. Whereas, from  $\text{H}_2 + \text{CO}_2$  it depends on the amount of ATP produced. In contrast, if the pathway to the desired product from acetyl-CoA requires an input of ATP, the production from  $\text{H}_2 + \text{CO}_2$  will not be possible. With CO as electron and carbon source it depends on the amount of energy required. As mentioned above, the overall ATP yield can be higher in *C. autoethanogenum*, when an electron-bifurcating methylene-THF reductase is assumed. In this case the ATP balance is zero, when  $\text{CO}_2$  is reduced to acetyl-CoA with  $\text{H}_2$  (Figure 5A), whereas in contrast 0.5 ATP per acetyl-CoA is produced with CO as substrate (Figure 5B).

Ethanol can be made from acetyl-CoA by two reduction steps *via* acetaldehyde (Figure 6). The reduction of acetyl-CoA to ethanol with NADH as electron donor is close to equilibrium ( $\Delta G^0 = -6.2 \text{ kJ/mol}$ ) and catalyzed by NADH-dependent enzymes like the bifunctional AdhE (Thauer et al., 1977; Goodlove et al., 1989; Peng et al., 2008; Yakushi and Matsushita, 2010; Extance et al., 2013; Bertsch et al., 2016). Thus, 2 NADH are required for the reduction of acetyl-CoA to ethanol. When a product like ethanol is formed from  $\text{H}_2 + \text{CO}_2$  *via* acetyl-CoA, the ATP gained in the acetate kinase reaction is missing and the overall ATP gain is negative with a demand of 0.7 ATP/per mol of ethanol in *A. woodii*. In addition, two more reducing equivalents are required; the reduction of  $2 \text{ NAD}^+$  with  $\text{H}_2$  as electron donor yields 0.6 ATP by action of hydrogenase, Rnf complex, and ATP synthase. In sum, ethanol formation has an energy demand of 0.1 ATP (Table 1). With CO as electron donor, ethanol formation would be possible with an ATP yield of 1.7 ATP/ethanol (Table 1). The same holds true for *C. autoethanogenum*. It was shown, that *C. autoethanogenum* can produce ethanol when growing on CO (Abrini et al., 1994; Abubackar et al., 2015) or  $\text{H}_2 + \text{CO}_2$  as electron donor (Mock et al., 2015). Ethanol formation from  $\text{H}_2 + \text{CO}_2$  has an energy demand of 0.2 ATP and with CO an ATP gain of 1.4 ATP/ethanol (Table 2). However, in *C. autoethanogenum*

ethanol production from  $H_2 + CO_2$  could be possible, in comparison to *A. woodii*, if a bifurcating methylene-THF reductase is assumed. In this scenario, *C. autoethanogenum* would gain 0.4 ATP per produced ethanol (Table 2). With CO the energetic for ethanol production in this case are even higher with 1.9 ATP/ethanol (Table 2).

**Table 1. ATP yield for the synthesis of products from acetyl-CoA with  $H_2 + CO_2$  or CO as electron donor in *A. woodii*.**

Product	Key enzymes/intermediates	Conversion (acetyl-CoA as precursor)	ATP yield	
			$H_2+CO_2$	CO
Acetate	Acetate kinase	acetyl-CoA→acetate	0.3	1.5
Ethanol	AIDH/ADH	acetyl-CoA→ethanol	-0.1	1.7
	AOR + ADH		0.3	2.1
Butanol	BDH	2 acetyl-CoA→butanol	-0.2	3.4
	BDH, bifurcating Bcd		0.4	4.0
	AOR		0.2	3.8
	AOR, bifurcating Bcd		0.8	4.4
Isoprene	Mevalonate	3 acetyl-CoA→isoprene + $CO_2$	-4.5	-0.3
Lactate	NADH-dependent LDH	acetyl-CoA + $CO_2$ →lactate	-0.7	1.1
	Bifurcating LDH		-0.1	1.7
2,3-Butanediol	Acetolactate synthase	2 acetyl-CoA→2,3-butanediol	-1.7	1.6
Acetone	Acetoacetate	2 acetyl-CoA→acetone + $CO_2$	-0.4	2.0
Isobutene	Acetone, 3-OH-isovalerate	3 acetyl-CoA→isobutene + 2 $CO_2$	-2.1	1.5

**Table 2. ATP yield for the synthesis of products from acetyl-CoA with  $H_2 + CO_2$  or CO as electron donor in *C. autoethanogenum*.**

Product	Key enzymes/intermediates	Conversion (acetyl-CoA as precursor)	Methylene-THF reductase		ATP yield	
			bifurcating	not bifurcating	$H_2+CO_2$	CO
Acetate	Acetate kinase	acetyl-CoA→acetate	-	+	0.4	1.0
Ethanol	AIDH /ADH	acetyl-CoA→ethanol	-	+	-0.2	1.4
	AOR + ADH		-	+	0.4	1.9



Butanol	BDH	2 acetyl-CoA→butanol	-	+	-0.4	2.8
	BDH, bifurcating Bcd		-	+	0.1	3.3
	AOR		-	+	0.1	3.3
	AOR, bifurcating Bcd		-	+	0.6	3.8
Isoprene	Mevalonate	3 acetyl-CoA→isoprene + CO <sub>2</sub>	-	+	-4.7	-0.9
Lactate	NADH-dependent LDH	acetyl-CoA + CO <sub>2</sub> →lactate	-	+	-0.7	0.9
	Bifurcating LDH		-	+	-0.2	1.4
2,3-Butanediol	Acetolactate synthase	2 acetyl-CoA→2,3-butanediol	-	+	-1.8	1.2
Acetone	Acetoacetate	2 acetyl-CoA→acetone + CO <sub>2</sub>	-	+	-0.5	1.7
Isobutene	Acetone, 3-OH-isovalerate	3 acetyl-CoA→isobutene + 2 CO <sub>2</sub>	-	+	-2.2	1.0
Acetate	Acetate kinase	acetyl-CoA→acetate	+	-	1.0	1.5
Ethanol	AIDH /ADH	acetyl-CoA→ethanol	+	-	0.4	1.9
	AOR + ADH		+	-	1.2	2.4
Butanol	BDH	2 acetyl-CoA→butanol	+	-	0.2	3.3
	BDH, bifurcating Bcd		+	-	0.8	3.9
	AOR		+	-	0.6	3.8
	AOR, bifurcating Bcd		+	-	1.1	4.3
Isoprene	Mevalonate	3 acetyl-CoA→isoprene + CO <sub>2</sub>	+	-	-2.3	-0.2
Lactate	NADH-dependent LDH	acetyl-CoA + CO <sub>2</sub> →lactate	+	-	0.2	1.4
	Bifurcating LDH		+	-	0.8	1.9
2,3-Butanediol	Acetolactate synthase	2 acetyl-CoA→2,3-butanediol	+	-	-1.2	1.9
Acetone	Acetoacetate	2 acetyl-CoA→acetone + CO <sub>2</sub>	+	-	0.1	2.0
Isobutene	Acetone, 3-OH-isovalerate	3 acetyl-CoA→isobutene + 2 CO <sub>2</sub>	+	-	-1.6	1.6

364

365 However, there is a second way of producing ethanol from acetyl-CoA in acetogens.

366 Aldehyde:ferredoxin oxidoreductases (AOR) are capable of catalyzing the reversible

367 reduction of an acid to the corresponding aldehyde, in this case, the reduction of acetate to

368 acetaldehyde (White et al., 1989; Nissen and Basen, 2019). The importance of AOR enzymes

are discussed in more detail later in this review. The redox potential of acetate/acetaldehyde ( $E_0' = -580$  mV) is so negative that a low potential electron donor such as ferredoxin is required (Thauer et al., 1977). The further reduction of acetaldehyde to ethanol could be catalyzed by a monofunctional alcohol dehydrogenase (ADH) (Goodlove et al., 1989), or by the same AdhE as described above. Indeed, *C. autoethanogenum* has an AOR enzyme that together with ADH converts acetate to ethanol (Mock et al., 2015). Under these conditions, ethanol production from  $H_2 + CO_2$  yields 0.4/1.2 ATP/ethanol, while ethanol production from CO gains 1.9/2.4 ATP/ethanol (depending on the MTHFR reaction) (Table 2). *A. woodii* does not have an AOR (Poehlein et al., 2012), but if one is implemented by metabolic engineering, ethanol formation from  $H_2 + CO_2$  would be possible with an ATP yield of 0.3 ATP/ethanol (Table 1). With CO as electron donor, the energetics for ethanol production with an implemented AOR in *A. woodii* are even better, yielding 2.1 ATP/ethanol (Table 1).

The production of other valuable products in *A. woodii* or *C. autoethanogenum* such as lactate, butanol, acetone, isobutene or 2,3-butanediol has an energy demand with  $H_2 + CO_2$  but a higher net ATP gain with CO as electron donor (Table 1 and 2). To form lactate, acetyl-CoA has to be first carboxylated to pyruvate by a pyruvate:ferredoxin oxidoreductase (Figure 6) (Furdui and Ragsdale, 2000). Using NADH as electron donor, pyruvate is further reduced to lactate ( $\Delta G^0 = -25$  kJ/mol) by a lactate dehydrogenase (LDH) (Gladden, 2004). *A. woodii* has an electron-bifurcating lactate dehydrogenase (bLDH), which increases the amount of ATP produced *via* chemiosmosis (Weghoff et al., 2015). Even if a bifurcating LDH is involved, production of lactate from  $H_2 + CO_2$  still requires energy in *A. woodii* or *C. autoethanogenum* by -0.1 or -0.2/-0.7 ATP/lactate (depending on the MTHFR reaction) and, thus lactate production from  $H_2 + CO_2$  is not possible (Table 1 and 2). However, if an electron-bifurcating MTHFR is assumed, lactate production with or without a bifurcating LDH from  $H_2 + CO_2$  is possible (0.2 or 0.8 ATP/lactate) in *C. autoethanogenum* (Table 2). If CO is used as electron donor, the production of lactate with *A. woodii* or *C. autoethanogenum*

will always gain ATP, independent of any electron bifurcation event (MTHFR, bLDH) (Table 1 and 2). Due to the energetics, *A. woodii* can not produce lactate from  $H_2 + CO_2$  (Schoelmerich et al., 2018). In contrast, *C. autoethanogenum* has been shown to produce lactate from syngas (Köpke et al., 2011b; Liew et al., 2017).

For the production of butanol, 2 molecules of acetyl-CoA are required (Dürre et al., 1992). Two molecules of acetyl-CoA are condensed to acetoacetyl-CoA which is reduced to 3-hydroxypropionyl-CoA with NADH (Figure 6). After water is split off, crotonyl-CoA is reduced to butyrate by a butyryl-CoA dehydrogenase (Bcd). Due to the relative positive redox potential of the crotonyl-CoA/butyryl-CoA couple ( $E_0' = -10$  mV), NADH-dependent Bcds can couple crotonyl-CoA reduction to ferredoxin reduction (bBcD) (Li et al., 2008). If the Bcd is not electron bifurcating, the pathway requires 4 NADH and yields 1.2 ATP/butanol for *A. woodii* or 1.6/2.2 ATP/butanol for *C. autoethanogenum* (depending on the MTHFR reaction). If the Bcd is electron bifurcating, the additional conservation of energy via the Rnf complex leads to a total ATP yield of 1.8 ATP/butanol for *A. woodii* or 2.6/3.2 ATP/butanol for *C. autoethanogenum* (depending on the MTHFR reaction). Since 1.4 ATP in *A. woodii* or 1.2 ATP in *C. autoethanogenum* have to be invested to supply 2 acetyl-CoA, the butanol production from  $H_2 + CO_2$  via butyraldehyde dehydrogenase is only possible, if a bifurcating Bcd is involved (Table 1 and 2). In case of *C. autoethanogenum* the energetics for butanol production from  $H_2 + CO_2$  without a bifurcating Bcd (0.2 ATP/butanol) only gains ATP, if an electron-bifurcating MTHFR is assumed (Table 2).

Butanol production can also be achieved by reducing butyrate to butyraldehyde (Nissen and Basen, 2019). AORs have been shown to reduce a broad range of acids to the corresponding aldehydes, also the reduction of butyrate to butyraldehyde (Ni et al., 2012; Perez et al., 2013). The requirement of  $Fd^{2-}$  reduces the energy yield via chemiosmosis, but the formation of butyrate from butyryl-CoA yields 1 ATP via substrate-level phosphorylation. Butanol production via AORs is energetically possible for *A. woodii* and *C. autoethanogenum*

independent on the pathway and electron bifurcation (bBcD, MTHFR) (Table 1 and 2). With CO as electron donor, production of butanol will be, in any case, strongly energy positive, yielding 3.4 - 4.4 ATP/butanol for *A. woodii* (depending on the pathway) or 2.8 - 3.8/3.3 - 4.3 ATP/butanol for *C. authoethanogenum* (depending on the pathway/ MTHFR reaction) (Table 1 and 2).

For the production of 2,3-butanediol, 2 pyruvate are condensed and decarboxylated, yielding acetolactate (Figure 6) (Hess et al., 2015). This is further decarboxylated, giving rise to acetoin. The reduction of acetoin with NADH yields 2,3-butanediol. In sum, 2 acetyl-CoA are reduced with 2  $\text{Fd}^{2-}$  and 2 NADH to 2,3-butanediol. The use of  $\text{H}_2$  or CO as external electron donor makes big differences in the energy balance: with  $\text{H}_2$ , in *A. woodii* 1.7 ATP have to be invested for the synthesis of one 2,3-butanediol, with CO 1.6 ATP are produced for every 2,3-butanediol synthesized (Table 1). Using  $\text{H}_2$  as electron donor in *C. authoethanogenum* 2,3-butanediol production is not possible, even if the MTHFR is electron bifurcating, because the ATP/2,3-butanediol ratio is strongly ATP-dependent with -1.8/-1.2 ATP/2,3-butanediol (depending on the MTHFR reaction) (Table 2). Again, the energetic favourable reduction power of CO enhances the ATP yield also during 2,3-butanediol fermentation (1.2/1.9 ATP/2,3-butanediol; depending on the MTHFR reaction), which makes the production of 2,3-butanediol in *C. authoethanogenum* possible (Köpke et al., 2011b).

Recently, *A. woodii* was shown to produce acetone from  $\text{H}_2 + \text{CO}_2$  after implementation of the corresponding pathway from *Clostridium acetobutylicum*, but only as a side product in small amounts (Hoffmeister et al., 2016). During the production of acetone 2 acetyl-CoA are condensed to acetoacetyl-CoA, which is further converted to acetoacetate and then to acetone (Figure 6) (Dürre et al., 1992). The production of acetone from  $\text{H}_2 + \text{CO}_2$  by *A. woodii* would require an input of ATP (-0.4 ATP/acetone), while production from CO would be coupled to ATP production (2.0 ATP/acetone) (Table 1). In comparison, acetone production from  $\text{H}_2 + \text{CO}_2$  by *C. authoethanogenum* is possible, if an electron-bifurcating MTHFR is assumed

(Table 2). The overall ATP/acetone yield is in this case positive, with 0.1 ATP per produced acetone (Table 2). With CO, *C. authoethanogenum* has an even better ATP yield with 1.7/2.0 ATP/acetone (depending on the MTHFR reaction) (Table 2).

Acetone can be a precursor for further products. For example, for the production of isobutene (Figure 6) (van Leeuwen et al., 2012). With the patented enzyme system for the acetylation of acetone to 3-hydroxy-isovalerate, followed by the ATP-dependent decarboxylation, isobutene could be produced in microorganisms (van Leeuwen et al., 2012). Because an even higher ATP investment is necessary, compared to the production of acetone, the energetics for *A. woodii* or *C. authoethanogenum* with  $H_2 + CO_2$  as substrate are much worse (Table 1 and 2). The production of isobutene from  $H_2 + CO_2$  by *A. woodii* would require an input of ATP (-2.1 ATP/isobutene), while production from CO would be coupled to ATP production (1.5 ATP/isobutene) (Table 1). Using *C. authoethanogenum* for the production of isobutene, an input of ATP of -2.2/-1.6 per isobutene (depending on the MTHFR reaction) with  $H_2 + CO_2$  is needed (Table 2). With CO as electron donor, *C. authoethanogenum* has an ATP gain with 1.0/1.6 ATP/isobutene (depending on the MTHFR reaction) (Table 2).

The production of isoprene is even more energy consuming. To produce isoprene a non-mevalonate pathway has been engineered in acetogens (Diner et al., 2018). The mevalonate pathway starts with the condensation of 2 acetyl-CoA to acetoacetyl-CoA (Figure 6). The addition of a third acetyl-CoA yields 3-hydroxy-3-methyl-glutaryl-CoA (HMG-CoA), which is reduced by a  $NAD^+$ -dependent HMG-CoA reductase to mevalonate. After two consequent ATP-dependent phosphorylations, diphosphomevalonate is decarboxylated at the expense of ATP, giving rise to isopentenyl diphosphate (IPP) (Kuzuyama, 2002). After an isomerization, the diphosphate bond is hydrolyzed, yielding isoprene. Thus, in total, the conversion of 3 acetyl-CoA to isoprene requires 2 NADPH and 3 ATP. Because of the high energy input needed, both *A. woodii* or *C. authoethanogenum* have an ATP-demand per produced isoprene when growing on  $H_2 + CO_2$  or CO as substrate under all possible scenarios (Table 1 and 2).

## Overcoming energetic barriers

The reduction of CO<sub>2</sub> with H<sub>2</sub> or with CO as reductant is energetically limited and currently only two products, ethanol and acetate can be produced on an industrial scale (Daniell et al., 2012; Bengelsdorf et al., 2018). As outlined above, ethanol formation from H<sub>2</sub> + CO<sub>2</sub> via acetyl-CoA to acetaldehyde and ethanol requires the net input of ATP and should not be possible with high specificity. However, it is possible in some species, and these species have an enzyme that activates acetate ( $\Delta G^{0'} = -15.4$  kJ/mol) with reduced ferredoxin as electron donor, the aldehyde:ferredoxin oxidoreductase (AOR) (White et al., 1989; Chan et al., 1995; Heider et al., 1995; Nissen and Basen, 2019). Therefore, ethanol formation still includes the acetate kinase reaction. Although additional reduced ferredoxin is required as driving force and thus missing as driving force for ATP synthesis, the overall process yields 1.2 ATP/mol ethanol. Acetogens like *A. woodii* that do not have an AOR do not produce ethanol from H<sub>2</sub> + CO<sub>2</sub>, but with the advent of genetic methods in acetogens, implementing an AOR into the acetogenic metabolism would be the first choice to increase the energetics and overcome the energetic barrier to acetyl-CoA formation. In contrast, *C. autoethanogenum* produce ethanol from H<sub>2</sub> + CO<sub>2</sub> (Mock et al., 2015; Liew et al., 2017). The importance of AOR enzymes for higher ethanol production rates were shown, for example, in *Pyrococcus furiosus* or *C. autoethanogenum* (Basen et al., 2014; Liew et al., 2017). An engineered *P. furiosus* strain converted glucose to ethanol via acetate and acetaldehyde, catalyzed by the host-encoded aldehyde ferredoxin oxidoreductase (AOR) and heterologously expressed AdhA, in an energy-conserving, redox-balanced pathway (Basen et al., 2014; Müller, 2014). Implementing an indirect ethanol pathway by deletion of *AdhE* in the genome of *C. autoethanogenum* shifted the conversion of acetate to acetaldehyde via AOR (Liew et al., 2017). Using this strategy the generated strains produced up to 180% more ethanol and also accumulated up to 38% less of acetate. The first heterologous AOR production in an acetogenic bacterium was recently described for *Clostridium carboxidivorans* (Cheng et al., 2019). A plasmid-based

implementation of the AOR genes lead to higher ethanol production rates during growth on glucose. Furthermore strains overexpressing AOR showed CO<sub>2</sub> re-assimilation during heterotrophic growth on glucose (Cheng et al., 2019). However, the selectivity of AORs to reduce short chain fatty acids, like acetate, can also easily be achieved with simple changes in growth conditions (Abubackar et al., 2015; Mock et al., 2015; Martin et al., 2016; Richter et al., 2016; Richter, 2016). For syngas-fermenting bacteria, that are using the AOR-ADH pathway, solventogenesis (alcohol production) occurs preferably when growth is limited due to nutrient limitations or at low pH (Daniell et al., 2012; Abubackar et al., 2015; Mock et al., 2015; Al-Shorgani et al., 2018). For example, results from bioreactor studies with continuous CO supply revealed that the shift from high to low pH values improves ethanol production by *C. autoethanogenum* without any accumulation of acetic acid (Abubackar et al., 2015). Low pH is therefore not only beneficial for ethanol production *via* syngas fermentation, but also an effective method for decreasing acetate concentration in the fermentation broth (Martin et al., 2016; Richter et al., 2016; Abubackar et al., 2015).

The same principle would also apply to other alcohols. Butanol is an interesting biofuel and can be produced by acetogens from the condensation of two mol acetyl-CoA to acetoacetyl-CoA followed by a reduction to butyryl-CoA, *via* a series of reactions known from fatty acid oxidation (see above) (Dürre, 2007; Köpke et al., 2011c). Butyryl-CoA can be either oxidized to butyrate giving one ATP or reduced to butanol without the production of an ATP. In this pathway, the acetate kinase-produced ATP's (2 molecules) are missing and although one ATP is gained by the butyrate kinase, the overall ATP gain is negative and butyrate cannot be produced from H<sub>2</sub> + CO<sub>2</sub>. With CO as substrate the energetics are a bit better. The oxidation of CO is coupled to more ferredoxin reduction, which can be used as driving force to produce additional ATP. But butanol is not produced for energetic reasons, since a redirection of butyryl-CoA to butanol results in a loss of another ATP. Here, implementation of an AOR by genetic engineering would bring the energetics over the hurdle to produce butanol (Ni et al.,

2012; Perez et al., 2013). AORs so far characterized are rather unspecific and can oxidize chain aldehydes (e.g. formaldehyde, acetaldehyde, crotonaldehyde), branched chain aldehydes (e.g. isovalerylaldehyde) as well as aromatic aldehydes (phenylacetaldehyde, benzaldehyde) or reduce acetate and butyrate (Nissen and Basen, 2019). The specificity and efficiency of the AOR could be improved by directed evolution (Turner, 2003; Dalby, 2011).

As outlined above, energy conservation during acetogenesis from  $H_2 + CO_2$  involves substrate level phosphorylation as well as electron transport phosphorylation. The former yields one ATP per acetate, the latter much less, in the order of 0.3 and less per mol of acetate. So far, every acetogen examined has only one of the two respiratory enzymes, Ech (Schölmerich and Müller, 2019) or Rnf (Biegel et al., 2011). Interestingly, Ech complexes have thus far only been found in thermophilic species (Schölmerich and Müller, 2019). One idea to improve the energetics is to implement by genetic engineering an Ech complex into *A. woodii*, *C. autoethanogenum* or others, but this requires an Ech complex from a mesophilic bacterium. Fortunately, Ech-containing mesophilic anaerobes do exist, for example in the *Butyrivrio* clade (Figure 7) (Schölmerich et al., 2020) (Hackmann and Firkins, 2015). Currently, it is tested whether these can be functionally produced in *A. woodii*. Another obstacle is that *A. woodii* uses  $Na^+$  as coupling ion for the ATP synthase and the Rnf complex (Hess et al., 2013b; Matthies et al., 2014); ideally, the Ech complex implemented should also translocate  $Na^+$  but so far, the ion specificity of these Ech complexes as well as other is not known, simply due to the fact that they have never been purified and reconstituted into liposomes. However, even if an Ech complex is a proton and not a sodium ion pump, the proton gradient established would be converted to a  $Na^+$  gradient by a  $Na^+/H^+$  antiporter present in *A. woodii* (Biegel and Müller, 2011). The same principle could be used *vice versa*, but this would require a thermophilic Rnf complex, for example from *Thermotoga maritima* (Kuhns et al., 2020).



Membrane-bound cytochromes are well-known electron carriers in membrane-bound electron transport chains; their presence would immediately suggest an electron transport chain and electron transport phosphorylation in the organism they have been discovered in (Gottwald et al., 1975; Das et al., 1989; Das and Ljungdahl, 2003). Methanogenic archaea also use the Wood-Ljungdahl pathway for CO<sub>2</sub> fixation and methane formation but how this is coupled to energy conservation has been hotly debated a while ago (Blaut and Gottschalk, 1984; Lancaster, 1989). Some favoured substrate level phosphorylation by an unknown reaction in the WLP, other some sort of to be discovered electron transport phosphorylation. The discovery of b<sub>559</sub>-type cytochromes in *Methanosarcina barkeri* in 1979 by Kühn and Gottschalk was a ground-breaking study that excited the entire field and paved the road to the discovery of electron transport phosphorylation in this model methanogen (Kühn et al., 1979; Blaut and Gottschalk, 1984). Later, methanophenazine (Abken et al., 1998) was found as additional electron carrier in these cells and the heterodisulfide reductase as electron acceptor (Bäumer et al., 1998). Now, different electron transport chains with different electron input modules in different methanogens are well established (Deppenmeier, 2002; Schlegel and Müller, 2013; Welte and Deppenmeier, 2014). In this historic context was the discovery of *b*-type cytochromes in *Moorella thermoacetica* in 1975 by Gottwald et al.. That was immediately taken as indication for cytochrome-dependent electron transport chain as part of a chemiosmotic mechanism of ATP synthesis. Later, reduction of cytochromes as well as the concomitant generation of a membrane potential in vesicles of *C. thermoautotrophicum* was detected (Hugenholtz and Ljungdahl, 1989). The most conclusive evidence that the cytochromes are involved in coupling the WLP to energy conservation was presented by Kamlage and Blaut in 1993. By showing that a cytochrome-deficient mutant was no longer able to oxidize methyl groups to CO<sub>2</sub> or reduce CO<sub>2</sub> to the level of a methyl group, they laid the foundation that cytochromes are involved as electron carrier in the methylene-THF reductase, a reaction that was already postulated in 1977 by Thauer et al. to be the most likely

energy conserving site since reduction of methylene-THF to Methyl-THF ( $\Delta G^0 = -23.1$  kJ/mol) with  $\text{NAD}^+$  as reductant is the most exergonic of the pathway. An involvement of cytochromes in  $\text{CO}_2$  reduction is also supported by the finding that nitrate, an alternative electron acceptor in *M. thermoacetica*, represses the synthesis of the *b*-type cytochrome (Fröstl et al., 1996; Arendsen et al., 1999). An involvement of cytochromes in the methylene-THF reductase reaction was also very recently speculated by Keller et al. (2019) for *Thermacetogenium phaeum*. However, it is far from being settled whether or not there is indeed a third, cytochrome-dependent respiratory chain in addition to Ech- and Rnf-containing respiratory chains in acetogens. This must be addressed in future studies using the genetic tools that have been developed. However, it should be noted that Rnf- and Ech-acetogens such as *A. woodii* and *T. kivui* do not have cytochromes (Müller, 2003). Others seem to have Rnf plus cytochromes such as *Sporomusa* (Kamlage et al., 1993; Poehlein et al., 2013) or *Clostridium aceticum* (Poehlein et al., 2015) or Ech plus cytochromes such as *M. thermoacetica* (Gottwald et al., 1975; Pierce et al., 2008). Important for the context here is, that the current industrially relevant cells do not have cytochromes. However, implementing cytochromes by genetic engineering is not an easy task since a rather extensive biogenesis machinery as well as genes encoding biosynthesis of quinones are required (Thöny-Meyer, 1997).

In addition to  $\text{CO}_2$ , some acetogens can reduce a number of alternative substrates such as pyruvate (Misoph and Drake, 1996), fumarate (Dorn et al., 1978; Matthies et al., 1993), aromatic acrylates (Bache and Pfennig, 1981; Tschech and Pfennig, 1984; Misoph et al., 1996), inorganic sulfur compounds (Beatty and Ljungdahl, 1990; 1991; Hattori et al., 2000), and nitrate or nitrite (Seifritz et al., 1993; Fröstl et al., 1996; Arendsen et al., 1999; Seifritz et al., 2003). For industrial applications, the electron acceptor would have to be added as well as the product removed from the fermenter, thus raising the costs of operation and the price of the product. Of these, the use of aromatic acrylates, nitrate and nitrite have been studied to

some extent (Bache and Pfennig, 1981; Tschech and Pfennig, 1984; Seifritz et al., 1993; Fröstl et al., 1996; Arendsen et al., 1999; Seifritz et al., 2003). Aromatic acrylates such as caffeate are reduced by *A. woodii* simultaneously with CO<sub>2</sub> when H<sub>2</sub> or fructose is the electron donor, or sequential with CO<sub>2</sub> first, when methanol is the electron donor (Dilling et al., 2007). Caffeate respiration is linked to energy conservation *via* Rnf and ATP synthase and co-utilization of caffeate and CO<sub>2</sub> would enhance the ATP level (Imkamp and Müller, 2002; Hess et al., 2013a). However, caffeate is rather toxic and cells do not tolerate more than ~ 5 mM (Parke and Ornston, 2004), which, in addition to the costs, makes the approach economically less favourable. Nitrate is rather inexpensive but the effect of nitrate is discussed very controversially. Under heterotrophic conditions, nitrate is stimulatory but lithotrophic growth of *M. thermoactica* is inhibited (Seifritz et al., 2002). The basis for this inhibition remains to be elucidated, one study argues that the activity of enzymes of the WLP is not altered but only cytochromes disappear (Seifritz et al., 2002) whereas the other argues that also the activities of key enzymes are down regulated by nitrate (Fröstl et al., 1996). Mechanistically, it is not known whether nitrate reduction is coupled to energy conservation *via* a nitrate-dependent ion-motive respiratory chain or whether the presence of nitrate redirects electrons to the Rnf- and Ech-complexes thus providing more fuel for chemiosmotic ATP synthesis. In contrast, *C. ljungdahlii* co-utilized CO<sub>2</sub> and nitrate and this enhanced biomass formation from H<sub>2</sub> + CO<sub>2</sub> (Emerson et al., 2019). Unfortunately, ethanol production from H<sub>2</sub> + CO<sub>2</sub> was strongly reduced by nitrate, indicating that the electron from acetyl-CoA reduction to ethanol goes to nitrate. However, in pH-controlled bioreactors nitrate improved growth and ethanol formation but resulted in stochastic inhibition events (Klask et al., 2020). Clearly, more basic research has to be done on nitrate reduction and its possible coupling to energy conservation and in addition, one has to remember that nitrate reduction is only observed in a few species and, for example, not in *A. woodii*.

So far, we have discussed alterations in the pathway of carbon and electrons in acetogenesis from  $\text{H}_2 + \text{CO}_2$  or CO to increase chemiosmotic ATP synthesis. An alternative is to bring in a reaction (sequence) that is coupled to substrate level phosphorylation. To phosphorylate ADP the reaction must have a phosphoryl group transfer potential that is more negative than  $-31.8$  kJ/mol (Thauer et al., 1977). This restricts the reactions to only a few:


Creatine kinase:



Arginine kinase:



Carbamate kinase:



Succinyl-CoA synthetase:



Acetate kinase:



Butyrate kinase:




Phosphoglycerate kinase:



$$(\Delta G^{0'} = -51.9 \text{ kJ/mol})$$

Pyruvate kinase:




Arginine kinase and creatine kinase (equation 3 and 4) play an essential role in ATP buffering systems in invertebrates (Wallimann et al., 1992) and are not useful here since they require previous activation of the substrate by ATP. Reaction 4 is part of the arginine deaminase pathway, a pathway found in eukaryotes (Novak et al., 2016) as well as prokaryotes (Deibel,

1964) that provide ATP from a fermentative pathway. In that pathway arginine is converted to ornithine, ammonium, and  $\text{CO}_2$ , while generating ATP from ADP and phosphate. The enzymes involved in the three steps of the pathway are arginine deiminase (ADI) (Petrack et al., 1957), ornithine transcarbamylase (OTC) (Kalman et al., 1965; Hernandez and Johnson, 1967), and carbamate kinase (CK) (Grisolia et al., 1962). The first reaction, catalyzed by ADI, is the deamination of arginine to yield citrulline and  $\text{NH}_4^+$  (Petrack et al., 1957). OTC then catalyzes the conversion of citrulline and inorganic phosphate into carbamoyl-phosphate and ornithine (Kalman et al., 1965; Hernandez and Johnson, 1967). Finally, CK catalyzes the hydrolysis of carbamoyl phosphate to form  $\text{CO}_2$  and  $\text{NH}_4^+$ , while the phosphate group is used to regenerate ATP from ADP (Grisolia et al., 1962). Indeed, feeding arginine (20 mM) promoted growth advantages in *A. woodii* (Beck et al., 2020). A 69% higher maximal  $\text{OD}_{600}$  and about 60% lower acetate yield per biomass was obtained in the presence of arginine (Beck et al., 2020). However, the costs of arginine and the removal of ornithine and ammonium have to be considered. Succinyl-CoA synthetase is a reaction of the tricarboxylic acid cycle (TCC) (Hager, 1962); this could be useful by feeding citrate which is converted by the enzymes of the TCC to succinate. Uptake of citrate (Dimroth and Thomer, 1986) and export of succinate (Kimmich et al., 1991) must be ensured by the appropriate transporter; another disadvantage is that the reaction sequence requires four electrons (2 NADH). Acetate kinase and butyrate kinase have been discussed above as part of the butyrate/butanol pathway. Intermediates of this pathway are CoA esters, which are expensive, instable and probably not taken up (Banis et al., 1976). Feeding the precursors acetate or acetoacetate would require an ATP-dependent activation, so that the net ATP gain by SLP would be zero.

An additional strategy is to provide organic electron donors that are acetogenic substrates. It is known for some acetogens that they can utilize different substrates simultaneously, for example  $\text{H}_2 + \text{CO}_2$  together with glucose (Braun and Gottschalk, 1981). The WLP is ideally suited for capturing  $\text{CO}_2$  to increase the product yields. Homoacetogenesis according to

equation 11 and 12 can be regarded as mixotrophic growth: sugar is oxidized to acetate giving rise to ATP synthesis by the reactions of pyruvate kinase, phosphoglycerate kinase and acetate kinase.



The electrons gained during glycolysis are captured by CO<sub>2</sub> reduction (Schuchmann and Müller, 2016). Alternatively, interspecies H<sub>2</sub> transfer can be used instead of the WLP to get rid of surplus reducing equivalents as shown for *A. woodii* (Winter and Wolfe, 1980; Wiechmann et al., 2020) or *T. kivui* (Moon et al., 2020). In principle any oxidative pathway can be hooked up to the WLP; this is what makes the WLP and acetogens so interesting for biotechnology. On the downside, mixotrophy is limited by the redox balance (Schuchmann and Müller, 2016). Reduction of 2 CO<sub>2</sub> to acetate requires 8 electrons. Therefore, the first requirement for a pathway coupled to the WLP is the release of 8 electrons per 2 CO<sub>2</sub> produced to allow complete CO<sub>2</sub> fixation. This is the case for oxidation of glucose to 2 acetate by typical fermentations (Schuchmann and Müller, 2016). However, this approach is limited to end products with the same oxidation state as acetate, since glucose oxidation does not provide enough electrons. Electron limitation can be overcome by supplying additional electron donors such as H<sub>2</sub> or electrons provided by a cathode for direct electron uptake (Bajracharya et al., 2015; Philips, 2020). In this case, H<sub>2</sub> provides an unlimited source of reducing power (Liu and Suflita, 1995). Under these conditions, ethanol production from glucose and H<sub>2</sub> according to equation 13 is possible.




Other examples of acetogenic substrates are primary (Eichler and Schink, 1984; 1985) and secondary alcohols (Zellner and Winter, 1987). Oxidation of ethanol under anaerobic conditions is only possible by specialized bacteria or archaea that either transfer the electrons

to H<sub>2</sub> which is used by a syntrophic partner or to an electron acceptor such as CO<sub>2</sub> or sulfate (Eichler and Schink, 1985; Seitz et al., 1990; Schink and Friedrich, 1994; Schink, 1997). Butandiol oxidation requires the presence of CO<sub>2</sub> according to equation 14.



The energy released in this reaction (calculated without phosphorylation of ADP) is -89 kJ/mol 2,3-butanediol. The same is true for oxidation of methanol (equation 15) or ethanol (equation 16).



The strategies described so far all apply to pure cultures of acetogens. A fundamentally different strategy is to use consortia of different organisms (Stams, 1994; Stams and Plugge, 2009; Morris et al., 2013; Kouzuma et al., 2015; Angenent et al., 2016). The main products of acetogens such as acetate and ethanol are the substrates for chain elongation *via* reverse  $\beta$ -oxidation (Grootscholten et al., 2013; Angenent et al., 2016). Through a combination of syngas fermentation and chain elongation, C<sub>4</sub>, C<sub>6</sub> and C<sub>8</sub> products have been obtained by mixed cultures in open culture fermentation (Angenent et al., 2016). Alternatively, the products of H<sub>2</sub> + CO<sub>2</sub> or syngas fermentation, ethanol and/or acetate, can be fed to pure cultures of anaerobes such *Clostridium kluyveri* (Barker and Taha, 1942) to produce n-caproic acid (Barker and Taha, 1942; Gildemyn et al., 2017) or to aerobes such as yeasts (Kerbs et al., 1952; Okada, 1981) or *Acinetobacter baylyi* (Salcedo-Vite et al., 2019). Using this approach, the entire tool box is available for the production of higher value compounds.

## Concluding remarks

In times of global climate change alternatives must be explored for a future-proof (bio)technology and energy industry. Renewable energy sources must be used in processes

that do not emit CO<sub>2</sub> or, even better, capture CO<sub>2</sub> and convert it into valuable compounds that are otherwise made petrochemically. In addition, a future-proof energy industry must take into account molecular H<sub>2</sub> as promising energy carrier. Different biological solutions are available that solve the most urgent future problems of humanity and are currently tested. Among those are the acetogenic bacteria with their outstanding options to solve these problems. They are the most efficient biocatalysts for H<sub>2</sub> storage around and their activity is orders of magnitude higher than any chemical catalyst. They are already used on an industrial scale to convert syngas into ethanol and many more products are currently under investigation. They can not only use C<sub>1</sub> gases but also methanol, other primary and secondary alcohols, carbonic acids as well as sugars as driving force for CO<sub>2</sub> reduction and there is no other group of anaerobes that has nearly the same metabolic flexibility for CO<sub>2</sub> reduction that acetogens. The construction of new acetogenic bioplatfroms is still in its infancy but steadily increasing; the genetic tool box required is also steadily increasing. Acetogens have a CO<sub>2</sub> fixation pathway that is energy neutral and does not need additional ATP when acetate is the product. Other products that originate from acetyl-CoA require net input of ATP and, therefore, the main obstacle to overcome in acetogenic fermentations is the energy barrier. This review summarized current knowledge on the strategies to increase the energy charge of the bacteria, some of which have been tested; others could be tested in the future. The road is paved for acetogens to become the key players to achieve a better and more sustainable future for all.

## Acknowledgements

This work was funded by financial support of the Federal Ministry for Education and Research (BMBF) for the OBAC project (ERA-IB-16-018).



**Figure legends**

**Figure 1. The Wood-Ljungdahl pathway of CO<sub>2</sub> reduction.** Substrates fed directly into the pathway are shown to the left and right. Acetyl-CoA is the precursor of biomass and biocommodities (dotted arrows). [H], reducing equivalents; THF, tetrahydrofolic acid; [CO], enzyme-bound CO; large dashed arrows, oxidative direction of the pathway

**Figure 2. Respiratory enzyme complexes Rnf and Ech in acetogens.** Acetogens are classified in Rnf- (A) or Ech-containing (B) organisms. Exergonic electron transfer leads to the translocation of H<sup>+</sup>/Na<sup>+</sup> across the cytoplasmic membrane and the electrochemical H<sup>+</sup>/Na<sup>+</sup> potential is then the driving force for ATP synthesis. Reducing equivalents are oxidized via different pathways, results in different product formations. Rnf, model of *A. woodii*; Ech, model of *T. kivui*; [H], reducing equivalents; FMN, flavin mononucleotide

**Figure 3. Modularity of acetogenesis.** Reducing equivalents are generated during substrate utilization in acetogens. To maintain the redox balancing, different enzymes, like Rnf, Ech, electron-bifurcating hydrogenase, Nfn/Stn complex or the electron-bifurcating hydrogenase/formate dehydrogenase leads to a variety of different electron carriers, which can be reoxidized in the WLP. THF, tetrahydrofolic acid; Hyd, electron-bifurcating hydrogenase; FDH/Hyl, electron-bifurcating formate dehydrogenase; Nfn, electron-bifurcating, ferredoxin-dependent transhydrogenase; Stn, *Sporomusa* type Nfn

**Figure 4. Bioenergetics of acetate formation from H<sub>2</sub> + CO<sub>2</sub> and CO in *A. woodii*.** The reducing equivalents for the reductive steps in the WLP during CO<sub>2</sub> reduction (A) are provided by an H<sub>2</sub>-oxidizing, electron-bifurcating hydrogenase which reduces ferredoxin and NAD<sup>+</sup>. The reducing equivalents for the reductive steps during CO oxidation (B) are provided

by the CO-oxidizing CODH/ACS which reduces ferredoxin. Excess  $\text{Fd}^{2-}$  is oxidized by the Rnf complex which reduces  $\text{NAD}^+$  and builds up a  $\text{Na}^+$  gradient. This gradient drives ATP synthesis *via* the  $\text{Na}^+$ -dependent ATP synthase. In total, 0.3 ATP from  $\text{H}_2 + \text{CO}_2$  and 1.5 ATP from CO can be synthesized per acetate produced. CODH/ACS, CO dehydrogenase/acetyl coenzyme A synthase; THF, tetrahydrofolic acid

**Figure 5. Bioenergetics of acetate formation from  $\text{H}_2 + \text{CO}_2$  and CO in *C. autoethanogenum*.** The reducing equivalents for the reductive steps in the WLP during  $\text{CO}_2$  reduction (A, C) are provided by an  $\text{H}_2$ -oxidizing, electron-bifurcating formatehydrogenase complex (HytA-E/FDH) which reduces Fd,  $\text{NADP}^+$  and  $\text{CO}_2$ . The reducing equivalents for the reductive steps during CO oxidation (B, D) are provided by the CO-oxidizing CODH/ACS which reduces Fd. The Nfn complex is converting reducing equivalents Fd, NADH and NADPH. The methylene-THF reductase is assumed to be electron bifurcating in A and B. In C and D the methylene-THF reductase is not electron bifurcating. Excess  $\text{Fd}^{2-}$  is oxidized by the Rnf complex which reduces  $\text{NAD}^+$  and builds up a  $\text{H}^+$  gradient. This gradient drives ATP synthesis *via* the  $\text{H}^+$ -dependent ATP synthase. In total, 0.4/1 ATP from  $\text{H}_2 + \text{CO}_2$  and 1/1.5 ATP from CO (depending on the MTHFR reaction) can be synthesized per acetate produced. CODH/ACS, CO dehydrogenase/acetyl coenzyme A synthase; THF, tetrahydrofolic acid; Nfn, electron-bifurcating and ferredoxin-dependent transhydrogenase

**Figure 6. Metabolic pathways of acetyl-CoA reduction.** Acetyl-CoA is synthesized *via* the Wood–Ljungdahl pathway and can be reduced to lactate, 2,3 butanediol, ethanol, acetat, butanol, isoprene, acetone or isobutene. Cofactors involved in different metabolic pathways are indicated.

**Figure 7. Ech gene-cluster of representative mesophilic anaerobes.** Ech genes are highly distributed in the mesophilic *Butyrivibrio* clade. Typical anaerobes are for example *Pseudobutyrvibrio ruminis*, *Butyrivibrio proteoclasticus*, *Pseudobutyrvibrio xylanivorans* or *Butyrivibrio fibrisolvens*. Here shown are the Ech gene-cluster-comparison against representative archaea *M. mazei* and *M. barkeri*. Ech gene-cluster of mesophilic anaerobes show similar genetic organisation.

## References

- Abken, H.J., Tietze, M., Brodersen, J., Baumer, S., Beifuss, U., and Deppenmeier, U. (1998). Isolation and characterization of methanophenazine and function of phenazines in membrane-bound electron transport of *Methanosarcina mazei* Gö1. *J. Bacteriol.* 180, 2027-2032. doi: 10.1128/JB.180.8.2027-2032.1998
- Abrini, J., Naveau, H., and Nyns, E.J. (1994). *Clostridium autoethanogenum*, sp. nov., an anaerobic bacterium that produces ethanol from carbon monoxide. *Arch. Microbiol.* 161, 345-351. doi: 10.1007/BF00303591
- Abubackar, H.N., Veiga, M.C., and Kennes, C. (2015). Carbon monoxide fermentation to ethanol by *Clostridium autoethanogenum* in a bioreactor with no accumulation of acetic acid. *Bioresour. Technol.* 186, 122-127. doi: 10.1016/j.biortech.2015.02.113
- Ajanovic, A. (2011). Biofuels versus food production: Does biofuels production increase food prices? *Energy* 36, 2070-2076. doi: 10.1016/j.energy.2010.05.019
- Al-Shorgani, N.K.N., Kalil, M.S., Yusoff, W.M.W., and Hamid, A.A. (2018). Impact of pH and butyric acid on butanol production during batch fermentation using a new local isolate of *Clostridium acetobutylicum* YM1. *Saudi. J. Biol. Sci.* 25, 339-348. doi: 10.1016/j.sjbs.2017.03.020
- Andreesen, J.R., Schaupp, A., Neurauter, C., Brown, A., and Ljungdahl, L.G. (1973). Fermentation of glucose, fructose and xylose by *Clostridium thermoaceticum*: effect of metals on growth yield, enzymes and the synthesis of acetate from CO<sub>2</sub>. *J. Bacteriol.* 114, 743-751. doi: 10.1128/JB.114.2.743-751.1973
- Angenent, L.T., Richter, H., Buckel, W., Spirito, C.M., Steinbusch, K.J., Plugge, C.M., et al. (2016). Chain elongation with reactor microbiomes: Open-culture biotechnology to produce biochemicals. *Environ. Sci. Technol.* 50, 2796-2810. doi: 10.1021/acs.est.5b04847
- Anwar, M.N., Fayyaz, A., Sohail, N.F., Khokhar, M.F., Baqar, M., Khan, W.D., et al. (2018). CO<sub>2</sub> capture and storage: A way forward for sustainable environment. *J. Environ. Manage.* 226, 131-144. doi: 10.1016/j.jenvman.2018.08.009
- Arendsen, A.F., Soliman, M.Q., and Ragsdale, S.W. (1999). Nitrate-dependent regulation of acetate biosynthesis and nitrate respiration by *Clostridium thermoaceticum*. *J. Bacteriol.* 181, 1489-1495. doi: 10.1128/JB.181.5.1489-1495.1999
- Bache, R., and Pfennig, N. (1981). Selective isolation of *Acetobacterium woodii* on methoxylated aromatic acids and determination of growth yields. *Arch. Microbiol.* 130, 255-261. doi: 10.1007/BF00459530
- Bailera, M.L., Lisbona, P., Romeo, L.M., and Espatolero, S. (2017). Power to gas projects review: Lab, pilot and demo plants for storing renewable energy and CO<sub>2</sub>. *Renew. Sustain. Energy. Rev.* 69, 292-312. doi: 10.1016/j.rser.2016.11.130
- Bajracharya, S., ter Heijne, A., Benetton, X.D., Vanbroekhoven, K., Buisman, C.J., Strik, D.P.B.T.B., et al. (2015). Carbon dioxide reduction by mixed and pure cultures in

- 869 microbial electrosynthesis using an assembly of graphite felt and stainless steel as a  
870 cathode. *Bioresour. Technol.* 195, 14-24. doi: 10.1016/j.biortech.2015.05.081
- 871 Banis, R.J., Roberts, C.S., Stokes, G.B., and Tove, S.B. (1976). Procedure for enzymatic  
872 synthesis and isolation of radioactive long chain acyl-CoA esters. *Anal. Biochem.* 73,  
873 1-8. doi: 10.1016/0003-2697(76)90135-4
- 874 Barker, H.A., and Taha, S.M. (1942). *Clostridium kluyverii*, an organism concerned in the  
875 formation of caproic acid from ethyl alcohol. *J. Bacteriol.* 43, 347-363. doi:  
876 10.1128/JB.43.3.347-363.1942
- 877 Basen, M., Schut, G.J., Nguyen, D.M., Lipscomb, G.L., Benn, R.A., Prybol, C.J., et al.  
878 (2014). Single gene insertion drives bioalcohol production by a thermophilic archaeon.  
879 *Proc. Natl. Acad. Sci. U.S.A.* 111, 17618-17623. doi: 10.1073/pnas.1413789111
- 880 Bäumer, S., Murakami, E., Brodersen, J., Gottschalk, G., Ragsdale, S.W., and Deppenmeier,  
881 U. (1998). The F<sub>420</sub>H<sub>2</sub>:heterodisulfide oxidoreductase system from *Methanosarcina*  
882 species. 2-Hydroxyphenazine mediates electron transfer from F<sub>420</sub>H<sub>2</sub> dehydrogenase to  
883 heterodisulfide reductase. *FEBS Lett.* 428, 295-298. doi: 10.1016/s0014-  
884 5793(98)00555-9
- 885 Beaty, P.S., and Ljungdahl, L.G. (1990). Thiosulfate reduction by *Clostridium*  
886 *thermoaceticum* and *Clostridium thermoautotrophicum* during growth on methanol.  
887 *Ann. Meet. Am. Soc. Microbiol.* 199, 1-7.
- 888 Beaty, P.S., and Ljungdahl, L.G. (1991). Growth of *Clostridium thermoaceticum* on  
889 methanol, ethanol or dimethylsulfoxide. *Ann. Meet. Am. Soc. Microbiol.* K-131, 236.
- 890 Beck, M.H., Flaiz, M., Bengelsdorf, F.R., and Dürre, P. (2020). Induced heterologous  
891 expression of the arginine deiminase pathway promotes growth advantages in the  
892 strict anaerobe *Acetobacterium woodii*. *Appl. Microbiol. Biotechnol.* 104, 687-699.  
893 doi: 10.1007/s00253-019-10248-9
- 894 Bengelsdorf, F.R., Beck, M.H., Erz, C., Hoffmeister, S., Karl, M.M., Riegler, P., et al. (2018).  
895 Bacterial anaerobic synthesis gas (syngas) and CO<sub>2</sub>+H<sub>2</sub> Fermentation. *Adv. Appl.*  
896 *Microbiol.* 103, 143-221. doi: 10.1016/bs.aambs.2018.01.002
- 897 Benson, S.M., and Orr, F.M. (2008). Carbon dioxide capture and storage. *MRS Bulletin* 33,  
898 303-305. doi: 10.1557/mrs2008.63
- 899 Bertsch, J., and Müller, V. (2015). CO metabolism in the acetogen *Acetobacterium woodii*.  
900 *Appl. Environ. Microbiol.* 81, 5949-5956. doi: 10.1128/AEM.01772-15
- 901 Bertsch, J., Öppinger, C., Hess, V., Langer, J.D., and Müller, V. (2015). A heterotrimeric  
902 NADH-oxidizing methylenetetrahydrofolate reductase from the acetogenic bacterium  
903 *Acetobacterium woodii*. *J. Bacteriol.* 197, 1681-1689. doi: 10.1128/JB.00048-15
- 904 Bertsch, J., Siemund, A.L., Kremp, F., and Müller, V. (2016). A novel route for ethanol  
905 oxidation in the acetogenic bacterium *Acetobacterium woodii*: The  
906 acetaldehyde/ethanol dehydrogenase pathway. *Environ. Microbiol.* 18, 2913-2922.  
907 doi: 10.1111/1462-2920.13082

- 908 Biegel, E., and Müller, V. (2010). Bacterial Na<sup>+</sup>-translocating ferredoxin:NAD<sup>+</sup>  
909 oxidoreductase. *Proc. Natl. Acad. Sci. U.S.A.* 107, 18138-18142. doi:  
910 10.1073/pnas.1010318107
- 911 Biegel, E., and Müller, V. (2011). A Na<sup>+</sup>-translocating pyrophosphatase in the acetogenic  
912 bacterium *Acetobacterium woodii*. *J. Biol. Chem.* 286, 6080-6084. doi:  
913 10.1074/jbc.M110.192823
- 914 Biegel, E., Schmidt, S., González, J.M., and Müller, V. (2011). Biochemistry, evolution and  
915 physiological function of the Rnf complex, a novel ion-motive electron transport  
916 complex in prokaryotes. *Cell. Mol. Life Sci.* 68, 613-634. doi: 10.1007/s00018-010-  
917 0555-8
- 918 Blaut, M., and Gottschalk, G. (1984). Protonmotive force-driven synthesis of ATP during  
919 methane formation from molecular hydrogen and formaldehyde or carbon dioxide in  
920 *Methanosarcina barkeri*. *FEMS Microbiol. Lett.* 24, 103-107. doi: 10.1111/j.1574-  
921 6968.1984.tb01253.x
- 922 Brandt, K., and Müller, V. (2015). Hybrid rotors in F<sub>1</sub>F<sub>o</sub> ATP synthases: subunit  
923 composition, distribution, and physiological significance. *Biol. Chem.* 396, 1031-  
924 1042. doi: 10.1515/hsz-2015-0137
- 925 Braun, K., and Gottschalk, G. (1981). Effect of molecular hydrogen and carbon dioxide on  
926 chemo-organotrophic growth of *Acetobacterium woodii* and *Clostridium aceticum*.  
927 *Arch. Microbiol.* 128, 294-298. doi: 10.1007/BF00422533
- 928 Buckel, W., and Thauer, R.K. (2018a). Flavin-based electron bifurcation, a new mechanism  
929 of biological energy coupling. *Chem. Rev.* 118, 3862-3886. doi:  
930 10.1021/acs.chemrev.7b00707
- 931 Buckel, W., and Thauer, R.K. (2018b). Flavin-based electron bifurcation, ferredoxin,  
932 flavodoxin, and anaerobic respiration with protons (Ech) or NAD<sup>+</sup> (Rnf) as electron  
933 acceptors: a historical review. *Front. Microbiol.* 9, 401. doi:  
934 10.3389/fmicb.2018.00401
- 935 Chan, M.K., Mukund, S., Kletzin, A., Adams, M.W.W., and Rees, D.C. (1995). Structure of a  
936 hyperthermophilic tungstopterin enzyme, aldehyde ferredoxin oxidoreductase. *Science*  
937 267, 1463-1469. doi: 10.1126/science.7878465
- 938 Cheng, C., Li, W., Lin, M., and Yang, S.T. (2019). Metabolic engineering of *Clostridium*  
939 *carboxidivorans* for enhanced ethanol and butanol production from syngas and  
940 glucose. *Bioresour. Technol.* 284, 415-423. doi: 10.1016/j.biortech.2019.03.145
- 941 Claassen, P.A.M., van Lier, J.B., Lopez Contreras, A.M., van Niel, E.W.J., Sijtsma, L.,  
942 Stams, A.J.M., et al. (1999). Utilisation of biomass for the supply of energy carriers.  
943 *Appl. Microbiol. Biotechnol.* 52, 741-755. doi: 10.1007/s002530051586
- 944 Clark, J.E., and Ljungdahl, L.G. (1982). Purification and properties of 5,10-  
945 methenyltetrahydrofolate cyclohydrolase from *Clostridium formicoaceticum*. *J. Biol.*  
946 *Chem.* 257, 3833-3836.

- 947 Clark, J.E., and Ljungdahl, L.G. (1984). Purification and properties of 5,10-  
948 methylenetetrahydrofolate reductase, an iron-sulfur flavoprotein from *Clostridium*  
949 *formicoaceticum*. *J. Biol. Chem.* 259, 10845-10849.
- 950 Cotton, C.A., Claassens, N.J., Benito-Vaquerizo, S., and Bar-Even, A. (2019). Renewable  
951 methanol and formate as microbial feedstocks. *Curr. Opin. Biotechnol.* 62, 168-180.  
952 doi: 10.1016/j.copbio.2019.10.002
- 953 Dalby, P.A. (2011). Strategy and success for the directed evolution of enzymes. *Curr. Opin.*  
954 *Struct. Biol.* 21, 473-480. doi: 10.1016/j.sbi.2011.05.003
- 955 Daniell, J., Köpke, M., and Simpson, S.D. (2012). Commercial biomass syngas fermentation.  
956 *Energies* 5, 5372-5417. doi: 10.3390/en5125372
- 957 Das, A., Hugenholtz, J., Van Halbeek, H., and Ljungdahl, L.G. (1989). Structure and function  
958 of a menaquinone involved in electron transport in membranes of *Clostridium*  
959 *thermoautotrophicum* and *Clostridium thermoaceticum*. *J. Bacteriol.* 171, 5823-5829.  
960 doi: 10.1128/jb.171.11.5823-5829.1989
- 961 Das, A., Ivey, D.M., and Ljungdahl, L.G. (1997). Purification and reconstitution into  
962 proteoliposomes of the F<sub>1</sub>F<sub>0</sub> ATP synthase from the obligately anaerobic gram-  
963 positive bacterium *Clostridium thermoautotrophicum*. *J. Bacteriol.* 179, 1714-1720.  
964 doi: 10.1128/jb.179.5.1714-1720.1997
- 965 Das, A., and Ljungdahl, L.G. (2003). "Electron-transport systems in acetogens," in  
966 *Biochemistry and physiology of anaerobic bacteria*, ed. L.G. Ljungdahl, M.W.  
967 Adams, L.L. Barton, J.G. Ferry and M.K. Johnson. (New York: Springer), 191-204.
- 968 Deibel, R.H. (1964). Utilization of arginine as an energy source for the growth of  
969 *Streptococcus faecalis*. *J. Bacteriol.* 87, 988-992. doi: 10.1128/JB.87.5.988-992.1964
- 970 Deppenmeier, U. (2002). The unique biochemistry of methanogenesis. *Prog. Nucleic. Acid.*  
971 *Res. Mol. Biol.* 71, 223-283. doi: 10.1016/s0079-6603(02)71045-3
- 972 Dey, G.R. (2007). Chemical reduction of CO<sub>2</sub> to different products during photo catalytic  
973 reaction on TiO<sub>2</sub> under diverse conditions: an overview. *Journal of natural gas*  
974 *chemistry* 16, 217-226. doi: 10.1016/S1003-9953(07)60052-8
- 975 Diekert, G., Schrader, E., and Harder, W. (1986). Energetics of CO formation and CO  
976 oxidation in cell suspensions of *Acetobacterium woodii*. *Arch. Microbiol.* 144, 386-  
977 392. doi: 10.1007/BF00409889
- 978 Dilling, S., Imkamp, F., Schmidt, S., and Müller, V. (2007). Regulation of caffeate respiration  
979 in the acetogenic bacterium *Acetobacterium woodii*. *Appl. Environ. Microbiol.* 73,  
980 3630-3636. doi: 10.1128/AEM.02060-06
- 981 Dimroth, P., and Thomer, A. (1986). Citrate transport in *Klebsiella pneumoniae*. *Biol. Chem.*  
982 *Hoppe Seyler* 367, 813-823. doi: 10.1515/bchm3.1986.367.2.813
- 983 Diner, B.A., Fan, J., Scotcher, M.C., Wells, D.H., and Whited, G.M. (2018). Synthesis of  
984 heterologous mevalonic acid pathway enzymes in *Clostridium ljungdahlii* for the  
985 conversion of fructose and of syngas to mevalonate and isoprene. *Appl. Environ.*  
986 *Microbiol.* 84, e01723-01717. doi: 10.1128/AEM.01723-17

- 987 Dorn, M., Andreesen, J.R., and Gottschalk, G. (1978). Fermentation of fumarate and L-malate  
988 by *Clostridium formicoaceticum*. *J. Bacteriol.* 133, 26-32.
- 989 Drake, H.L., Daniel, S., Küsel, K., Matthies, C., Kuhner, C., and Braus-Strohmeyer, S.  
990 (1997). Acetogenic bacteria: what are the *in situ* consequences of their diverse  
991 metabolic diversities? *Biofactors* 1, 13-24. doi: 10.1002/biof.5520060103
- 992 Drake, H.L., Gößner, A.S., and Daniel, S.L. (2008). Old acetogens, new light. *Ann. N. Y.*  
993 *Acad. Sci.* 1125, 100-128. doi: 10.1196/annals.1419.016
- 994 Drake, H.L., Hu, S.I., and Wood, H.G. (1980). Purification of carbon monoxide  
995 dehydrogenase, a nickel enzyme from *Clostridium thermocaceticum*. *J. Biol. Chem.*  
996 255, 7174-7180.
- 997 Drake, H.L., Hu, S.I., and Wood, H.G. (1981). Purification of five components from  
998 *Clostridium thermoaceticum* which catalyze synthesis of acetate from pyruvate and  
999 methyltetrahydrofolate. *J. Biol. Chem.* 56, 11137-11144
- 1000 Dürre, P. (2017). Gas fermentation - a biotechnological solution for today's challenges.  
1001 *Microb. Biotechnol.* 10, 14-16. doi: 10.1111/1751-7915.12431
- 1002 Dürre, P. (2007). Biobutanol: An attractive biofuel. *Biotechnol. J.* 2, 1525-1534. doi:  
1003 10.1002/biot.200700168
- 1004 Dürre, P., Bahl, H., and Gottschalk, G. (1992). Acetone-butanol fermentation - basis of a  
1005 modern biotechnological process. *Chem. Ing. Tech.* 64, 491-498. doi:  
1006 10.1002/cite.330640603
- 1007 Eichler, B., and Schink, B. (1984). Oxidation of primary aliphatic alcohols by *Acetobacterium*  
1008 *carbinolicum* sp. nov., a homoacetogenic anaerobe. *Arch. Microbiol.* 140, 147-152.  
1009 doi: 10.1007/BF00454917
- 1010 Eichler, B., and Schink, B. (1985). Fermentation of primary alcohols and diols and pure  
1011 culture of syntrophically alcohol-oxidizing anaerobes. *Arch. Microbiol.* 143, 60-66.  
1012 doi: 10.1007/BF00414769
- 1013 Emerson, D.F., Woolston, B.M., Liu, N., Donnelly, M., Currie, D.H., and Stephanopoulos, G.  
1014 (2019). Enhancing hydrogen-dependent growth of and carbon dioxide fixation by  
1015 *Clostridium ljungdahlii* through nitrate supplementation. *Biotechnol. Bioeng.* 116,  
1016 294-306. doi: 10.1002/bit.26847
- 1017 Enzmann, F., Mayer, F., Rother, M., and Holtmann, D. (2018). Methanogens: biochemical  
1018 background and biotechnological applications. *AMB Expr.* 8, 1. doi: 10.1186/s13568-  
1019 017-0531-x
- 1020 Extance, J., Crennell, S.J., Eley, K., Cripps, R., Hough, D.W., and Danson, M.J. (2013).  
1021 Structure of a bifunctional alcohol dehydrogenase involved in bioethanol generation in  
1022 *Geobacillus thermoglucosidasius*. *Acta. Crystallogr. D. Biol. Crystallogr.* 69, 2104-  
1023 2115. doi: 10.1107/S0907444913020349
- 1024 Ferguson, S.A., Keis, S., and Cook, G.M. (2006). Biochemical and molecular characterization  
1025 of a Na<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub>-ATPase from the thermoalkaliphilic bacterium  
1026 *Clostridium paradoxum*. *J. Bacteriol.* 188, 5045-5054. doi: 10.1128/JB.00128-06



- 1027 Fillingame, R.H., Angevine, C.M., and Dmitriev, O.Y. (2003). Mechanics of coupling proton  
1028 movements to *c*-ring rotation in ATP synthase. *FEBS Lett.* 555, 29-34. doi:  
1029 10.1016/S0005-2728(02)00250-5
- 1030 Fritz, M., and Müller, V. (2007). An intermediate step in the evolution of ATPases - the F<sub>1</sub>F<sub>0</sub>-  
1031 ATPase from *Acetobacterium woodii* contains F-type and V-type rotor subunits and is  
1032 capable of ATP synthesis. *FEBS J.* 274, 3421-3428. doi: 10.1111/j.1742-  
1033 4658.2007.05874.x
- 1034 Fröstl, J.M., Seifritz, C., and Drake, H.L. (1996). Effect of nitrate on the autotrophic  
1035 metabolism of the acetogens *Clostridium thermoautotrophicum* and *Clostridium*  
1036 *thermoaceticum*. *J. Bacteriol.* 178, 4597-4603. doi: 10.1128/jb.178.15.4597-  
1037 4603.1996
- 1038 Fuchs, G. (2011). Alternative pathways of carbon dioxide fixation: Insights into the early  
1039 evolution of life? *Annu. Rev. Microbiol.* 65, 631-658. doi: 10.1146/annurev-micro-  
1040 090110-102801
- 1041 Furdui, C., and Ragsdale, S.W. (2000). The role of pyruvate ferredoxin oxidoreductase in  
1042 pyruvate synthesis during autotrophic growth by the Wood-Ljungdahl pathway. *J.*  
1043 *Biol. Chem.* 275, 28494-28499. doi: 10.1074/jbc.M003291200M003291200
- 1044 Gildemyn, S., Molitor, B., Usack, J.G., Nguyen, M., Rabaey, K., and Angenent, L.T. (2017).  
1045 Upgrading syngas fermentation effluent using *Clostridium kluyveri* in a continuous  
1046 fermentation. *Biotechnol. Biofuels* 10, 83. doi: 10.1186/s13068-017-0764-6
- 1047 Gladden, L.B. (2004). Lactate metabolism: a new paradigm for the third millennium. *J.*  
1048 *Physiol.* 558, 5-30. doi: 10.1113/jphysiol.2003.058701jphysiol.2003.058701
- 1049 Goodlove, P.E., Cunningham, P.R., Parker, J., and Clark, D.P. (1989). Cloning and sequence  
1050 analysis of the fermentative alcohol-dehydrogenase-encoding gene of *Escherichia*  
1051 *coli*. *Gene* 85, 209-214. doi: 10.1016/0378-1119(89)90483-6
- 1052 Gottwald, M., Andreesen, J.R., LeGall, J., and Ljungdahl, L.G. (1975). Presence of  
1053 cytochrome and menaquinone in *Clostridium formicoaceticum* and *Clostridium*  
1054 *thermoaceticum*. *J. Bacteriol.* 122, 325-328.
- 1055 Götz, M.L., Mörs, J., Koch, F., Graf, A.M., Bajohr, F., Reimert, S., et al. (2016). Renewable  
1056 power-to-gas: A technological and economic review. *Renew. Energy* 85, 1371-1390.  
1057 doi: 10.1016/j.renene.2015.07.066
- 1058 Grisolia, S., Harmon, P., and Rajman, L. (1962). Studies on the mechanism of action of  
1059 carbamate kinase. *Biochim. Biophys. Acta.* 62, 293-299. doi: 10.1016/0006-  
1060 3002(62)90042-2
- 1061 Grootsholten, T.I., Steinbusch, K.J., Hamelers, H.V., and Buisman, C.J. (2013). Chain  
1062 elongation of acetate and ethanol in an upflow anaerobic filter for high rate MCFA  
1063 production. *Bioresour. Technol.* 135, 440-445. doi: 10.1016/j.biortech.2012.10.165
- 1064 Hackmann, T.J., and Firkins, J.L. (2015). Electron transport phosphorylation in rumen  
1065 butyrovibrios: unprecedented ATP yield for glucose fermentation to butyrate. *Front.*  
1066 *Microbiol.* 6, 622. doi: 10.3389/fmicb.2015.00622

- 1067 Hager, L.B. (1962). "Succinyl CoA synthetase" in *The enzymes*, ed. P.D. Boyer, H. Lardy,  
1068 and K. Myrback (New York: Academic Press), 387-399.
- 1069 Hattori, S., Kamagata, Y., Hanada, S., and Shoun, H. (2000). *Thermacetogenium phaeum* gen.  
1070 nov., sp. nov., a strictly anaerobic, thermophilic, syntrophic acetate-oxidizing  
1071 bacterium. *Int. J. Syst. Evol. Microbiol.* 50, 1601-1609. doi: 10.1099/00207713-50-4-  
1072 1601
- 1073 Hedderich, R., and Forzi, L. (2005). Energy-converting [NiFe] hydrogenases: more than just  
1074 H<sub>2</sub> activation. *J. Mol. Microbiol. Biotechnol.* 10, 92-104. doi: 10.1159/000091557
- 1075 Heider, J., Ma, K., and Adams, M.W. (1995). Purification, characterization, and metabolic  
1076 function of tungsten-containing aldehyde ferredoxin oxidoreductase from the  
1077 hyperthermophilic and proteolytic archaeon *Thermococcus* strain ES-1. *J. Bacteriol.*  
1078 177, 4757-4764. doi: 10.1128/jb.177.16.4757-4764.1995
- 1079 Heijstra, B.D., Leang, C., and Juminaga, A. (2017). Gas fermentation: cellular engineering  
1080 possibilities and scale up. *Microb. Cell Fact.* 16, 60. doi: 10.1186/s12934-017-0676-y
- 1081 Hernandez, E., and Johnson, M.J. (1967). Anaerobic growth yields of *Aerobacter cloacae* and  
1082 *Escherichia coli*. *J. Bacteriol.* 94, 991-995. doi: 10.1128/JB.94.4.991-995.1967
- 1083 Hess, V., Gonzalez, J.M., Parthasarathy, A., Buckel, W., and Müller, V. (2013a). Caffeate  
1084 respiration in the acetogenic bacterium *Acetobacterium woodii*: a coenzyme A loop  
1085 saves energy for caffeate activation. *Appl. Environ. Microbiol.* 79, 1942-1947. doi:  
1086 10.1128/AEM.03604-12
- 1087 Hess, V., Oyrik, O., Trifunovic, D., and Müller, V. (2015). 2,3-butanediol metabolism in the  
1088 acetogen *Acetobacterium woodii*. *Appl. Environ. Microbiol.* 81, 4711-4719. doi:  
1089 10.1128/AEM.00960-15
- 1090 Hess, V., Schuchmann, K., and Müller, V. (2013b). The ferredoxin:NAD<sup>+</sup> oxidoreductase  
1091 (Rnf) from the acetogen *Acetobacterium woodii* requires Na<sup>+</sup> and is reversibly coupled  
1092 to the membrane potential. *J. Biol. Chem.* 288, 31496-31502. doi:  
1093 10.1074/jbc.M113.510255
- 1094 Himes, R.H., and Harmony, J.A. (1973). Formyltetrahydrofolate synthetase. *CRC Crit. Rev.*  
1095 *Biochem.* 1, 501-535. doi: 10.3109/10409237309105441
- 1096 Hoffmeister, S., Gerdorf, M., Bengelsdorf, F.R., Linder, S., Fluchter, S., Oztürk, H., et al.  
1097 (2016). Acetone production with metabolically engineered strains of *Acetobacterium*  
1098 *woodii*. *Metab. Eng.* 36, 37-47. doi: 10.1016/j.ymben.2016.03.001
- 1099 Hugenholtz, J., and Ljungdahl, L.G. (1989). Electron transport and electrochemical proton  
1100 gradient in membrane vesicles of *Clostridium thermoaceticum*. *J. Bacteriol.* 171,  
1101 2873-2875. doi: 10.1128/jb.171.5.2873-2875.1989
- 1102 Imkamp, F., and Müller, V. (2002). Chemiosmotic energy conservation with Na<sup>+</sup> as the  
1103 coupling ion during hydrogen-dependent caffeate reduction by *Acetobacterium*  
1104 *woodii*. *J. Bacteriol.* 184, 1947-1951. doi: 10.1128/JB.184.7.1947-1951.2002

- 1105 Ivey, D.M., and Ljungdahl, L.G. (1986). Purification and characterization of the F<sub>1</sub>-ATPase  
1106 from *Clostridium thermoaceticum*. *J. Bacteriol.* 165, 252-257. doi:  
1107 10.1128/jb.165.1.252-257.1986
- 1108 Kalman, S.M., Duffield, P.H., and Brzozowski, T. (1965). Identity in *Escherichia coli* of  
1109 carbamyl phosphokinase and an activity which catalyzes amino group transfer from  
1110 glutamine to ornithine in citrulline synthesis. *Biochem. Biophys. Res. Commun.* 18,  
1111 530-537. doi: 10.1016/0006-291x(65)90786-2
- 1112 Kamlage, B., and Blaut, M. (1993). Isolation of a cytochrome-deficient mutant strain of  
1113 *Sporomusa sphaeroides* not capable of oxidizing methyl groups. *J. Bacteriol.* 175,  
1114 3043-3050. doi: 10.1128/jb.175.10.3043-3050.1993
- 1115 Kamlage, B., Boelter, A., and Blaut, M. (1993). Spectroscopic and potentiometric  
1116 characterization of cytochromes in 2 *Sporomusa* species and their expression during  
1117 growth on selected substrates. *Arch. Microbiol.* 159, 189-196. doi:  
1118 10.1007/BF00250281
- 1119 Keller, A., Schink, B., and Müller, N. (2019). Alternative pathways of acetogenic ethanol and  
1120 methanol degradation in the thermophilic anaerobe *Thermacetogenium phaeum*.  
1121 *Front. Microbiol.* 10, 423. doi: 10.3389/fmicb.2019.00423
- 1122 Kerbs, H.A., Gurin, S., and Eggleston, L.V. (1952). The pathway of oxidation of acetate in  
1123 baker's yeast. *Biochem. J.* 51, 614-628. doi: 10.1042/bj0510614
- 1124 Kim, Y.M., and Hegeman, G.D. (1983). Oxidation of carbon monoxide by bacteria. *Int. Rev.*  
1125 *Cytol.* 81, 1-32. doi: 10.1016/s0074-7696(08)62333-5
- 1126 Kimmich, G.A., Randles, J., and Bennett, E. (1991). Sodium-dependent succinate transport by  
1127 isolated chick intestinal cells. *Am. J. Physiol.* 260, C1151-C1157. doi:  
1128 10.1152/ajpcell.1991.260.6.C1151
- 1129 Klask, C.M., Kliem-Kuster, N., Molitor, B., and Angenent, L.T. (2020). Nitrate feed improves  
1130 growth and ethanol production of *Clostridium ljungdahlii* with CO<sub>2</sub> and H<sub>2</sub>, but results  
1131 in stochastic inhibition events. *Front. Microbiol.* 11, 724. doi:  
1132 10.3389/fmicb.2020.00724
- 1133 Köpke, M., Mihalcea, C., Bromley, J.C., and Simpson, S.D. (2011a). Fermentative production  
1134 of ethanol from carbon monoxide. *Curr. Opin. Biotechnol.* 22, 320-325. doi:  
1135 10.1016/j.copbio.2011.01.005
- 1136 Köpke, M., Mihalcea, C., Liew, F., Tizard, J.H., Ali, M.S., Conolly, J.J., et al. (2011b). 2,3-  
1137 butanediol production by acetogenic bacteria, an alternative route to chemical  
1138 synthesis, using industrial waste gas. *Appl. Environ. Microbiol.* 77, 5467-5475. doi:  
1139 10.1128/AEM.00355-11
- 1140 Köpke, M., and Simpson, S.D. (2020). Pollution to products: recycling of 'above ground'  
1141 carbon by gas fermentation. *Curr. Opin. Biotechnol.* 65, 180-189. doi:  
1142 10.1016/j.copbio.2020.02.017
- 1143 Köpke, M., Steffi, N., and Dürre, P. (2011c). "The past, present, and future of biofuels—  
1144 biobutanol as promising alternative," in *Biofuel Production - Recent Developments*  
1145 *and Prospects*, ed. D.M.A.D.S. Bernardes. (Rijeka, Croatia: InTech), 451-486.

- 1146 Kouzuma, A., Kato, S., and Watanabe, K. (2015). Microbial interspecies interactions: recent  
1147 findings in syntrophic consortia. *Front. Microbiol.* 6, 477. doi:  
1148 10.3389/fmicb.2015.00477
- 1149 Kremp, F., Roth, J., and Müller, V. (2020). The *Sporomusa* type Nfn is a novel type of  
1150 electron-bifurcating transhydrogenase that links the redox pools in acetogenic bacteria.  
1151 *Sci. Rep.* 10, 14872. doi: 10.1038/s41598-020-71038-2
- 1152 Kühn, W., Fiebig, K., Walther, R., and Gottschalk, G. (1979). Presence of a cytochrome  $b_{559}$   
1153 in *Methanosarcina barkeri*. *FEBS Lett.* 105, 271-274. doi: 10.1016/0014-  
1154 5793(79)80627-4
- 1155 Kuhns, M., Trifunović, D., Huber, H., and Müller, V. (2020). The Rnf complex is a  $\text{Na}^+$   
1156 coupled respiratory enzyme in a fermenting bacterium, *Thermotoga maritima*.  
1157 *Commun. Biol.* 3, 431. doi: 10.1038/s42003-020-01158-y
- 1158 Kumagai, H., Fujiwara, T., Matsubara, H., and Saeki, K. (1997). Membrane localization,  
1159 topology, and mutual stabilization of the *rnfABC* gene products in *Rhodobacter*  
1160 *capsulatus* and implications for a new family of energy-coupling NADH  
1161 oxidoreductases. *Biochemistry* 36, 5509-5521. doi: 10.1021/bi970014q
- 1162 Kuzuyama, T. (2002). Mevalonate and nonmevalonate pathways for the biosynthesis of  
1163 isoprene units. *Biosci. Biotechnol. Biochem.* 66, 1619-1627. doi: 10.1271/bbb.66.1619
- 1164 Lancaster, J.R. (1989). Sodium, protons, and energy coupling in the methanogenic bacteria. *J.*  
1165 *Bioenerg. Biomembr.* 21, 717-740. doi: 10.1007/BF00762689
- 1166 Leung, D.L.C., Caramanna, G., and Maroto-Valer, M.M. (2014). An overview of current  
1167 status of carbon dioxide capture and storage technologies. *Renewable and sustainable*  
1168 *energy reviews* 39, 426-443. doi: 10.1016/j.rser.2014.07.093
- 1169 Li, F., Hinderberger, J., Seedorf, H., Zhang, J., Buckel, W., and Thauer, R.K. (2008). Coupled  
1170 ferredoxin and crotonyl coenzyme A (CoA) reduction with NADH catalyzed by the  
1171 butyryl-CoA dehydrogenase/Etf complex from *Clostridium kluyveri*. *J. Bacteriol.* 190,  
1172 843-850. doi: 10.1128/JB.01417-07
- 1173 Liew, F., Henstra, A.M., Köpke, M., Winzer, K., Simpson, S.D., and Minton, N.P. (2017).  
1174 Metabolic engineering of *Clostridium autoethanogenum* for selective alcohol  
1175 production. *Metab. Eng.* 40, 104-114. doi: 10.1016/j.ymben.2017.01.007
- 1176 Liew, F., Martin, M.E., Tappel, R.C., Heijstra, B.D., Mihalcea, C., and Köpke, M. (2016).  
1177 Gas fermentation - a flexible platform for commercial scale production of low-carbon-  
1178 fuels and chemicals from waste and renewable feedstocks. *Front. Microbiol.* 7, 694.  
1179 doi: 10.3389/fmicb.2016.00694
- 1180 Liu, S., and Suflita, J.M. (1995).  $\text{H}_2$  as an energy source for mixotrophic acetogenesis from  
1181 the reduction of  $\text{CO}_2$  and syringate by *Acetobacterium woodii* and *Eubacterium*  
1182 *limosum*. *Curr. Microbiol.* 31, 245-250. doi: 10.1007/BF00298382
- 1183 Litty, D., and Müller, V. (2020). A  $\text{Na}^+$   $\text{A}_1\text{A}_0$  ATP synthase with a V-type *c* subunit in a  
1184 mesophilic bacterium. *FEBS J.* 287, 3012-3023. doi: 10.1111/febs.15193

- 1185 Ljungdahl, L.G. (1986). The autotrophic pathway of acetate synthesis in acetogenic bacteria.  
1186 *Ann. Rev. Microbiol.* 40, 415-450. doi: 10.1146/annurev.mi.40.100186.002215
- 1187 Lo, J., Zheng, T., Hon, S., Olson, D.G., and Lynd, L.R. (2015). The bifunctional alcohol and  
1188 aldehyde dehydrogenase gene, *adhE*, is necessary for ethanol production in  
1189 *Clostridium thermocellum* and *Thermoanaerobacterium saccharolyticum*. *J. Bacteriol.*  
1190 197, 1386-1393. doi: 10.1128/JB.02450-14
- 1191 Lovell, C.R., Przybyla, A., and Ljungdahl, L.G. (1988). Cloning and expression in  
1192 *Escherichia coli* of the *Clostridium thermoaceticum* gene encoding thermostable  
1193 formyltetrahydrofolate synthetase. *Arch. Microbiol.* 149, 280-285. doi:  
1194 10.1007/BF00411642
- 1195 Martin, M.E., Richter, H., Saha, S., and Angenent, L.T. (2016). Traits of selected *Clostridium*  
1196 strains for syngas fermentation to ethanol. *Biotechnol. Bioeng.* 113, 531-539. doi:  
1197 10.1002/bit.25827
- 1198 Matthies, C., Freiberger, A., and Drake, H.L. (1993). Fumarate dissimilation and differential  
1199 reductant flow by *Clostridium formicoaceticum* and *Clostridium aceticum*. *Arch.*  
1200 *Microbiol.* 160, 273-278. doi: 10.1007/BF00292076
- 1201 Matthies, D., Zhou, W., Klyszejko, A.L., Anselmi, C., Yildiz, O., Brandt, K., et al. (2014).  
1202 High-resolution structure and mechanism of an F/V-hybrid rotor ring in a Na<sup>+</sup>-coupled  
1203 ATP synthase. *Nat. Commun.* 5, 5286. doi: 10.1038/ncomms6286
- 1204 Mayer, F., Ivey, D.M., and Ljungdahl, L.G. (1986). Macromolecular organization of F<sub>1</sub>-  
1205 ATPase isolated from *Clostridium thermoaceticum* as revealed by electron  
1206 microscopy. *J. Bacteriol.* 166, 1128-1130. doi: 10.1128/jb.166.3.1128-1130.1986  
1207
- 1208 Mayer, F., Lim, J.K., Langer, J.D., Kang, S.G., and Müller, V. (2015). Na<sup>+</sup> transport by the  
1209 A<sub>1</sub>A<sub>0</sub>-ATP synthase purified from *Thermococcus onnurineus* and reconstituted into  
1210 liposomes. *J. Biol. Chem.* 290, 6994-7002. doi: 10.1074/jbc.M114.616862  
1211
- 1212 Meier, T., Ferguson, S.A., Cook, G.M., Dimroth, P., and Vonck, J. (2006). Structural  
1213 investigations of the membrane-embedded rotor ring of the F-ATPase from  
1214 *Clostridium paradoxum*. *J. Bacteriol.* 188, 7759-7764. doi: 10.1128/JB.00934-06  
1215
- 1216 Meier, T., Krah, A., Bond, P.J., Pogoryelov, D., Diederichs, K., and Faraldo-Gómez, J.D.  
1217 (2009). Complete ion-coordination structure in the rotor ring of Na<sup>+</sup>-dependent F-ATP  
1218 synthases. *J. Mol. Biol.* 391, 498-507. doi: 10.1016/j.jmb.2009.05.082  
1219
- 1220 Meuer, J., Kuettner, H.C., Zhang, J.K., Hedderich, R., and Metcalf, W.W. (2002). Genetic  
1221 analysis of the archaeon *Methanosarcina barkeri* Fusaro reveals a central role for Ech  
1222 hydrogenase and ferredoxin in methanogenesis and carbon fixation. *Proc. Natl. Acad.*  
1223 *Sci. U.S.A.* 99, 5632-5637. doi: 10.1073/pnas.072615499072615499
- 1224 Misoph, M., Daniel, S.L., and Drake, H.L. (1996). Bidirectional usage of ferulate by the  
1225 acetogen *Peptostreptococcus productus* U-1: CO<sub>2</sub> and aromatic acrylate groups as  
1226 competing electron acceptors. *Microbiology* 142, 1983-1988. doi: 10.1099/13500872-  
1227 142-8-1983

- 1228 Misoph, M., and Drake, H.L. (1996). Effect of CO<sub>2</sub> on the fermentation capacities of the  
1229 acetogen *Peptostreptococcus productus* U-1. *J. Bacteriol.* 178, 3140-3145. doi:  
1230 10.1128/jb.178.11.3140-3145.1996
- 1231 Mock, J., Wang, S., Huang, H., Kahnt, J., and Thauer, R.K. (2014). Evidence for a  
1232 hexaheteromeric methylenetetrahydrofolate reductase in *Moorella thermoacetica*. *J.*  
1233 *Bacteriol.* 196, 3303-3314. doi: 10.1128/JB.01839-14
- 1234 Mock, J., Zheng, Y., Müller, A.P., Ly, S., Tran, L., Segovia, S., et al. (2015). Energy  
1235 conservation associated with ethanol formation from H<sub>2</sub> and CO<sub>2</sub> in *Clostridium*  
1236 *autoethanogenum* involving electron bifurcation. *J. Bacteriol.* 197, 2965-2980. doi:  
1237 10.1128/JB.00399-15
- 1238 Moon, J., Jain, S., Müller, V., and Basen, M. (2020). Homoacetogenic conversion of mannitol  
1239 by the thermophilic acetogenic bacterium *Thermoanaerobacter kivui* requires external  
1240 CO<sub>2</sub>. *Front. Microbiol.* 11, 571736. doi: 10.3389/fmicb.2020.571736.
- 1241 Moore, M.R., O'Brien, W.E., and Ljungdahl, L.G. (1974). Purification and characterization of  
1242 nicotinamide adenine dinucleotide-dependent methylenetetrahydrofolate  
1243 dehydrogenase from *Clostridium formicoaceticum*. *J. Biol. Chem.* 249, 5250-5253.
- 1244 Morris, B.E., Henneberger, R., Huber, H., and Moissl-Eichinger, C. (2013). Microbial  
1245 syntrophy: interaction for the common good. *FEMS Microbiol. Rev.* 37, 384-406. doi:  
1246 10.1111/1574-6976.12019
- 1247 Müller, V. (2003). Energy conservation in acetogenic bacteria. *Appl. Environ. Microbiol.* 69,  
1248 6345-6353. doi: 10.1128/aem.69.11.6345-6353.2003
- 1249 Müller, V. (2008). "Bacterial fermentation" in *Encyclopedia of life sciences*. (Chichester:  
1250 John Wiley & Sons Ltd). doi: 10.1002/9780470015902.a0001415.pub2
- 1251 Müller, V. (2014). Bioalcohol production by a new synthetic route in a hyperthermophilic  
1252 archaeon. *Proc. Natl. Acad. Sci. U.S.A.* 111, 17352-17353. doi:  
1253 10.1073/pnas.1420385111
- 1254 Müller, V. (2019). New horizons in acetogenic conversion of one-carbon substrates and  
1255 biological hydrogen storage. *Trends Biotechnol.* 37, 1344-1354. doi:  
1256 10.1016/j.tibtech.2019.05.008
- 1257 Müller, V., Chowdhury, N.P., and Basen, M. (2018). Electron bifurcation: A long-hidden  
1258 energy-coupling mechanism. *Annu. Rev. Microbiol.* 72, 331-353. doi:  
1259 10.1146/annurev-micro-090816-093440.
- 1260 Müller, V., and Grüber, G. (2003). ATP synthases: structure, function and evolution of unique  
1261 energy converters. *Cell. Mol. Life Sci.* 60, 474-494. doi: 10.1007/s000180300040
- 1262 Müller, V., Imkamp, F., Biegel, E., Schmidt, S., and Dilling, S. (2008). Discovery of a  
1263 ferredoxin:NAD<sup>+</sup>-oxidoreductase (Rnf) in *Acetobacterium woodii*: a novel potential  
1264 coupling site in acetogens. *Ann. N. Y. Acad. Sci.* 1125, 137-146. doi:  
1265 10.1196/annals.1419.011

1266

- 1267 Müller, V., Inkamp, F., Rauwolf, A., Küsel, K., and Drake, H.L. (2004). "Molecular and  
1268 cellular biology of acetogenic bacteria," in *Strict and facultative anaerobes: Medical  
1269 and environmental aspects*, ed. M. Nakano and P. Zuber. (Norfolk: Horizon Scientific  
1270 Press), 251-281.
- 1271 Nagarajan, H., Sahin, M., Nogales, J., Latif, H., Lovley, D.R., Ebrahim, A., et al. (2013).  
1272 Characterizing acetogenic metabolism using a genome-scale metabolic reconstruction  
1273 of *Clostridium ljungdahlii*. *Microb. Cell Fact.* 12, 118. doi: 10.1186/1475-2859-12-  
1274 118
- 1275 Naik, S.N., Goud, V.V., Rout, P.K., and Dalai, A.K. (2010). Production of first and second  
1276 generation biofuels: A comprehensive review. *Renew. Sust. Energ. Rev.* 14, 578-597.  
1277 doi: 10.1016/j.rser.2009.10.003
- 1278 Nguyen, D.M.N., Schut, G.J., Zadvornyy, O.A., Tokmina-Lukaszewska, M., Poudel, S.,  
1279 Lipscomb, G.L., et al. (2017). Two functionally distinct NADP<sup>+</sup>-dependent ferredoxin  
1280 oxidoreductases maintain the primary redox balance of *Pyrococcus furiosus*. *J. Biol.*  
1281 *Chem.* 292, 14603-14616. doi: 10.1074/jbc.M117.794172
- 1282 Ni, Y., Hagedoorn, P.L., Xu, J.H., Arends, I.W., and Hollmann, F. (2012). A biocatalytic  
1283 hydrogenation of carboxylic acids. *Chem. Commun.* 48, 12056-12058. doi:  
1284 10.1039/C2CC36479D
- 1285 Nissen, L.S., and Basen, M. (2019). The emerging role of aldehyde:ferredoxin  
1286 oxidoreductases in microbially-catalyzed alcohol production. *J. Biotechnol.* 306, 105-  
1287 117. doi: 10.1016/j.jbiotec.2019.09.005
- 1288 Novak, L., Zubacova, Z., Karnkowska, A., Kolisko, M., Hroudova, M., Stairs, C.W., et al.  
1289 (2016). Arginine deiminase pathway enzymes: evolutionary history in metamonads  
1290 and other eukaryotes. *BMC Evol. Biol.* 16, 197. doi: 10.1186/s12862-016-0771-4
- 1291 Novelli, P.C., Lang, P.M., Masarie, K.A., Hurst, D.F., Myers, R., and Elkins, J.W. (1999).  
1292 Molecular hydrogen in the troposphere: Global distribution and budget. *J. Geophys.*  
1293 *Res.* 104, 30427-30444. doi: Doi 10.1029/1999jd900788
- 1294 O'Brien, W.E., Brewer, J.M., and Ljungdahl, L.G. (1973). Purification and characterization of  
1295 thermostable 5,10-methylenetetrahydrofolate dehydrogenase from *Clostridium*  
1296 *thermoaceticum*. *J. Biol. Chem.* 248, 403-408.
- 1297 Okada, W., Fukuda, H., and Morikawa, H. (1981). Kinetic expressions of ethanol production  
1298 rate and ethanol consumption rate in baker's yeast cultivation. *J. Ferment. Technol.* 59,  
1299 103-109.
- 1300 Parke, D., and Ornston, L.N. (2004). Toxicity caused by hydroxycinnamoyl-coenzyme A  
1301 thioester accumulation in mutants of *Acinetobacter* sp. strain ADP1. *Appl. Environ.*  
1302 *Microbiol.* 70, 2974-2983. doi: 10.1128/aem.70.5.2974-2983.2004
- 1303 Peng, H., Wu, G., and Shao, W. (2008). The aldehyde/alcohol dehydrogenase (AdhE) in  
1304 relation to the ethanol formation in *Thermoanaerobacter ethanolicus* JW200.  
1305 *Anaerobe* 14, 125-127. doi: 10.1016/j.anaerobe.2007.09.004

- 1306 Perez, J.M., Richter, H., Loftus, S.E., and Angenent, L.T. (2013). Biocatalytic reduction of  
1307 short-chain carboxylic acids into their corresponding alcohols with syngas  
1308 fermentation. *Biotechnol. Bioeng.* 110, 1066-1077. doi: 10.1002/bit.24786
- 1309 Petrack, B., Sullivan, L., and Ratner, S. (1957). Behavior of purified arginine desiminase from  
1310 *S. faecalis*. *Arch. Biochem. Biophys.* 69, 186-197. doi: 10.1016/0003-9861(57)90485-x
- 1311 Pezacka, E., and Wood, H.G. (1984). The synthesis of acetyl-CoA by *Clostridium*  
1312 *thermoaceticum* from carbon dioxide, hydrogen, coenzyme A and  
1313 methyltetrahydrofolate. *Arch. Microbiol.* 137, 63-69. doi: 10.1007/BF00425809
- 1314 Philips, J. (2020). Extracellular electron uptake by acetogenic bacteria: Does H<sub>2</sub> consumption  
1315 favor the H<sub>2</sub> evolution reaction on a cathode or metallic iron? *Front. Microbiol.* 10,  
1316 2997. doi: 10.3389/fmicb.2019.02997
- 1317 Pierce, E., Xie, G., Barabote, R.D., Saunders, E., Han, C.S., Detter, J.C., et al. (2008). The  
1318 complete genome sequence of *Moorella thermoacetica* (f. *Clostridium*  
1319 *thermoaceticum*). *Environ. Microbiol.* 10, 2550-2573. doi: 10.1111/j.1462-  
1320 2920.2008.01679.x
- 1321 Poehlein, A., Cebulla, M., Ilg, M.M., Bengelsdorf, F.R., Schiel-Bengelsdorf, B., Whited, G.,  
1322 et al. (2015). The complete genome sequence of *Clostridium aceticum*: a missing link  
1323 between Rnf- and cytochrome-containing autotrophic acetogens. *mBio* 6, e01168-  
1324 01115. doi: 10.1128/mBio.01168-15
- 1325 Poehlein, A., Gottschalk, G., and Daniel, R. (2013). First insights into the genome of the  
1326 Gram-negative, endospore-forming organism *Sporomusa ovata* strain H1 DSM 2662.  
1327 *Genome Announc.* 1, e00734-00713. doi: 10.1128/genomeA.00734-13
- 1328 Poehlein, A., Schmidt, S., Kaster, A.-K., Goenrich, M., Vollmers, J., Thürmer, A., et al.  
1329 (2012). An ancient pathway combining carbon dioxide fixation with the generation  
1330 and utilization of a sodium ion gradient for ATP synthesis. *PLoS One* 7, e33439. doi:  
1331 10.1371/journal.pone.0033439
- 1332 Ragsdale, S.W., and Ljungdahl, L.G. (1984). Purification and properties of NAD-dependent  
1333 5,10-methylenetetrahydrofolate dehydrogenase from *Acetobacterium woodii*. *J. Biol.*  
1334 *Chem.* 259, 3499-3503.
- 1335 Ragsdale, S.W., and Pierce, E. (2008). Acetogenesis and the Wood-Ljungdahl pathway of  
1336 CO<sub>2</sub> fixation. *Biochim. Biophys. Acta* 1784, 1873-1898. doi:  
1337 10.1016/j.bbapap.2008.08.012
- 1338 Ragsdale, S.W., and Wood, H.G. (1985). Acetate biosynthesis by acetogenic bacteria.  
1339 Evidence that carbon monoxide dehydrogenase is the condensing enzyme that  
1340 catalyzes the final steps in the synthesis. *J. Biol. Chem.* 260, 3970-3977.
- 1341 Rahlfs, S., and Müller, V. (1999). Sequence of subunit *a* of the Na<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub>-  
1342 ATPase of *Acetobacterium woodii*: proposal for residues involved in Na<sup>+</sup> binding.  
1343 *FEBS Lett.* 453, 35-40. doi: 10.1016/s0014-5793(99)00576-1
- 1344



- 1345 Raybuck, S.A., Bastian, N.R., Orme-Johnson, W.H., and Walsh, C.T. (1988). Kinetic  
1346 characterization of the carbon monoxide-acetyl-CoA (carbonyl group) exchange  
1347 activity of the acetyl-CoA synthesizing CO dehydrogenase from *Clostridium*  
1348 *thermoaceticum*. *Biochemistry* 27, 7698-7702. doi: 10.1021/bi00420a019
- 1349 Reidlinger, J., Mayer, F., and Müller, V. (1994). The molecular structure of the Na<sup>+</sup>-  
1350 translocating F<sub>1</sub>F<sub>0</sub>-ATPase of *Acetobacterium woodii*, as revealed by electron  
1351 microscopy, resembles that of H<sup>+</sup>-translocating ATPases. *FEBS Lett.* 356, 17-20. doi:  
1352 10.1016/0014-5793(94)01222-9  
1353
- 1354 Reidlinger, J., and Müller, V. (1994). Purification of ATP synthase from *Acetobacterium*  
1355 *woodii* and identification as a Na<sup>+</sup>-translocating F<sub>1</sub>F<sub>0</sub>-type enzyme. *Eur. J. Biochem.*  
1356 223, 275-283. doi: 10.1111/j.1432-1033.1994.tb18992.x  
1357
- 1358 Richter, H., Molitor, B., Diender, M., Sousa, D.Z., and Angenent, L.T. (2016). A narrow pH  
1359 range supports butanol, hexanol, and octanol production from syngas in a continuous  
1360 co-culture of *Clostridium ljungdahlii* and *Clostridium kluyveri* with in-line product  
1361 extraction. *Front. Microbiol.* 7, 1773. doi: 10.3389/fmicb.2016.01773
- 1362 Richter, H., Molitor, B., Wei, H., Chen, W., Aristilde, L., and Angenent, L.T. (2016). Ethanol  
1363 production in syngas-fermenting *Clostridium ljungdahlii* is controlled by  
1364 thermodynamics rather than by enzyme expression. *Energ. Environ. Sci.* 9, 2392-  
1365 2399. doi: 10.1039/C6EE01108J
- 1366 Rittmann, S., and Herwig, C. (2012). A comprehensive and quantitative review of dark  
1367 fermentative biohydrogen production. *Microb. Cell Fact.* 11, 115. doi: 10.1186/1475-  
1368 2859-11-115
- 1369 Saeki, K., and Kumagai, H. (1998). The *rnf* gene products in *Rhodobacter capsulatus* play an  
1370 essential role in nitrogen fixation during anaerobic DMSO-dependent growth in the  
1371 dark. *Arch. Microbiol.* 169, 464-467. doi: 10.1007/s002030050598
- 1372 Salcedo-Vite, K., Sigala, J.C., Segura, D., Gosset, G., and Martinez, A. (2019). *Acinetobacter*  
1373 *baylyi* ADP1 growth performance and lipid accumulation on different carbon sources.  
1374 *Appl. Microbiol. Biotechnol.* 103, 6217-6229. doi: 10.1007/s00253-019-09910-z
- 1375 Schaupp, A., and Ljungdahl, L.G. (1974). Purification and properties of acetate kinase from  
1376 *Clostridium thermoaceticum*. *Arch. Microbiol.* 100, 121-129. doi:  
1377 10.1007/BF00446312
- 1378 Schiel-Bengelsdorf, B., and Dürre, P. (2012). Pathway engineering and synthetic biology  
1379 using acetogens. *FEBS Lett.* 586, 2191-2198. doi: 10.1016/j.febslet.2012.04.043
- 1380 Schink, B. (1997). Energetics of syntrophic cooperation in methanogenic degradation.  
1381 *Microbiol. Mol. Biol. Rev.* 61, 262-280.
- 1382 Schink, B., and Friedrich, M. (1994). Energetics of syntrophic fatty acid oxidation. *FEMS*  
1383 *Microbiol. Rev.* 15, 85-94.
- 1384 Schlegel, K., and Müller, V. (2013). Evolution of Na<sup>+</sup> and H<sup>+</sup> bioenergetics in methanogenic  
1385 archaea. *Biochem. Soc. Trans.* 41, 421-426. doi: 10.1042/BST20120294

- 1386 Schmehl, M., Jahn, A., Meyer zu Vilsendorf, A., Hennecke, S., Masepohl, B., Schuppler, M.,  
1387 et al. (1993). Identification of a new class of nitrogen fixation genes in *Rhodobacter*  
1388 *capsulatus*: a putative membrane complex involved in electron transport to  
1389 nitrogenase. *Mol. Gen. Genet.* 241, 602-615. doi: 10.1007/BF00279903
- 1390 Schölmerich, M.C., Katsyv, A., Sung, W., Mijic, V., Wiechmann, A., Kottenhahn, P., et al.  
1391 (2018). Regulation of lactate metabolism in the acetogenic bacterium *Acetobacterium*  
1392 *woodii*. *Environ. Microbiol.* 20, 4587-4595. doi: 10.1111/1462-2920.14412
- 1393 Schölmerich, M.C., and Müller, V. (2019). Energy-converting hydrogenases: The link  
1394 between H<sub>2</sub> metabolism and energy conservation. *Cell. Mol. Life Sci.* 77, 1461-1481.  
1395 doi: 10.1007/s00018-019-03329-5
- 1396 Schölmerich, M.C., Katsyv, A., Dönig, J., Hackmann, T.J., and Müller, V. (2020). Energy  
1397 conservation involving 2 respiratory circuits. *Proc. Natl. Acad. Sci. U.S.A.* 117, 1167-  
1398 1173. doi: 10.1073/pnas.1914939117
- 1399 Schölmerich, M.C., and Müller, V. (2019). Energy conservation by a hydrogenase-dependent  
1400 chemiosmotic mechanism in an ancient metabolic pathway. *Proc. Natl. Acad. Sci.*  
1401 *U.S.A.* 116, 6329-6334. doi: 10.1007/s00018-019-03329-5
- 1402 Schuchmann, K., and Müller, V. (2012). A bacterial electron bifurcating hydrogenase. *J. Biol.*  
1403 *Chem.* 287, 31165–31171. doi: 10.1074/jbc.M112.395038
- 1404 Schuchmann, K., and Müller, V. (2013). Direct and reversible hydrogenation of CO<sub>2</sub> to  
1405 formate by a bacterial carbon dioxide reductase. *Science* 342, 1382-1385. doi:  
1406 10.1126/science.1244758
- 1407 Schuchmann, K., and Müller, V. (2014). Autotrophy at the thermodynamic limit of life: A  
1408 model for energy conservation in acetogenic bacteria. *Nat. Rev. Microbiol.* 12, 809-  
1409 821. doi: 10.1038/nrmicro3365
- 1410 Schuchmann, K., and Müller, V. (2016). Energetics and application of heterotrophy in  
1411 acetogenic bacteria. *Appl. Environ. Microbiol.* 82, 4056-4069. doi:  
1412 10.1128/AEM.00882-16
- 1413 Schut, G.J., and Adams, M.W. (2009). The iron-hydrogenase of *Thermotoga maritima* utilizes  
1414 ferredoxin and NADH synergistically: a new perspective on anaerobic hydrogen  
1415 production. *J. Bacteriol.* 191, 4451-4457. doi: 10.1128/JB.01582-08
- 1416 Schwarz, F.M., and Müller, V. (2020). Whole-cell biocatalysis for hydrogen storage and  
1417 syngas conversion to formate using a thermophilic acetogen. *Biotechnol. Biofuels* 13,  
1418 32. doi: 10.1186/s13068-020-1670-x
- 1419 Schwarz, F.M., Schuchmann, K., and Müller, V. (2018). Hydrogenation of CO<sub>2</sub> at ambient  
1420 pressure catalyzed by a highly active thermostable biocatalyst. *Biotechnol. Biofuels*  
1421 11, 237. doi: 10.1186/s13068-018-1236-3
- 1422 Seifritz, C., Daniel, S.L., Gößner, A., and Drake, H.L. (1993). Nitrate as a preferred electron  
1423 sink for the acetogen *Clostridium thermoaceticum*. *J. Bacteriol.* 175, 8008-8013. doi:  
1424 10.1128/jb.175.24.8008-8013.1993

- 1425 Seifritz, C., Drake, H.L., and Daniel, S.L. (2003). Nitrite as an energy-conserving electron  
1426 sink for the acetogenic bacterium *Moorella thermoacetica*. *Curr. Microbiol.* 46, 329-  
1427 333. doi: 10.1007/s00284-002-3830-6
- 1428 Seifritz, C., Fröstl, J.M., Drake, H.L., and Daniel, S.L. (2002). Influence of nitrate on oxalate-  
1429 and glyoxylate-dependent growth and acetogenesis by *Moorella thermoacetica*. *Arch.*  
1430 *Microbiol.* 178, 457-464. doi: 10.1007/s00203-002-0475-6
- 1431 Seitz, H.J., Schink, B., Pfennig, N., and Conrad, R. (1990). Energetics of syntrophic ethanol  
1432 oxidation in defined chemostat cocultures. *Arch. Microbiol.* 155, 89-93. doi:  
1433 10.1007/BF00291279
- 1434 Seravalli, J., Kumar, M., Lu, W.P., and Ragsdale, S.W. (1997). Mechanism of carbon  
1435 monoxide oxidation by the carbon monoxide dehydrogenase/acetyl-CoA synthase  
1436 from *Clostridium thermoaceticum*: Kinetic characterization of the intermediates.  
1437 *Biochemistry* 36, 11241-11251. doi: 10.1021/bi970590m
- 1438 Stams, A.J.M. (1994). Metabolic interactions between anaerobic bacteria in methanogenic  
1439 environments. *Antonie van Leeuwenhoek* 66, 271-294. doi: 10.1007/BF00871644
- 1440 Stams, A.J.M., and Plugge, C.M. (2009). Electron transfer in syntrophic communities of  
1441 anaerobic bacteria and archaea. *Nature Reviews Microbiology* 7, 568-577. doi:  
1442 10.1038/nrmicro2166
- 1443 Syu, M.J. (2001). Biological production of 2,3-butanediol. *Appl. Microbiol. Biotechnol.* 55,  
1444 10-18. doi: 10.1007/s002530000486
- 1445 Thauer, R.K., Jungermann, K., and Decker, K. (1977). Energy conservation in chemotrophic  
1446 anaerobic bacteria. *Bact. Rev.* 41, 100-180.
- 1447 Thöny-Meyer, L. (1997). Biogenesis of respiratory cytochromes in bacteria. *Microbiol. Mol.*  
1448 *Biol. Rev.* 61, 337-376.
- 1449 Tremblay, P.L., Zhang, T., Dar, S.A., Leang, C., and Lovley, D.R. (2012). The Rnf complex  
1450 of *Clostridium ljungdahlii* is a proton-translocating ferredoxin:NAD<sup>+</sup> oxidoreductase  
1451 essential for autotrophic growth. *MBio* 4, e00406-00412. doi: 10.1128/mBio.00406-12
- 1452 Tschech, A., and Pfennig, N. (1984). Growth yield increase linked to caffeate reduction in  
1453 *Acetobacterium woodii*. *Arch. Microbiol.* 137, 163-167. doi: 10.1007/BF00414460
- 1454 Turner, N.J. (2003). Directed evolution of enzymes for applied biocatalysis. *Trends*  
1455 *Biotechnol.* 21, 474-478. doi: 10.1016/j.tibtech.2003.09.001
- 1456 van Leeuwen, B.N., van der Wulp, A.M., Duijnste, I., van Maris, A.J., and Straathof, A.J.  
1457 (2012). Fermentative production of isobutene. *Appl. Microbiol. Biotechnol.* 93, 1377-  
1458 1387. doi: 10.1007/s00253-011-3853-7
- 1459 Wallimann, T., Wyss, M., Brdiczka, D., Nicolay, K., and Eppenberger, H.M. (1992).  
1460 Intracellular compartmentation, structure and function of creatine kinase isoenzymes  
1461 in tissues with high and fluctuating energy demands: the 'phosphocreatine circuit' for  
1462 cellular energy homeostasis. *Biochem. J.* 281, 21-40. doi: 10.1042/bj2810021

- 1463 Wang, S., Huang, H., Kahnt, J., Müller, A.P., Köpke, M., and Thauer, R.K. (2013a). NADP-  
1464 specific electron-bifurcating [FeFe]-hydrogenase in a functional complex with formate  
1465 dehydrogenase in *Clostridium autoethanogenum* grown on CO. *J. Bacteriol.* 195,  
1466 4373-4386. doi: 10.1128/JB.00678-13
- 1467 Wang, S., Huang, H., Kahnt, J., and Thauer, R.K. (2013b). *Clostridium acidurici* electron-  
1468 bifurcating formate dehydrogenase. *App. Environ. Microbiol.* 79, 6176-6179. doi:  
1469 10.1128/AEM.02015-13
- 1470 Wang, S., Huang, H., Moll, J., and Thauer, R.K. (2010). NADP<sup>+</sup> reduction with reduced  
1471 ferredoxin and NADP<sup>+</sup> reduction with NADH are coupled *via* an electron bifurcating  
1472 enzyme complex in *Clostridium kluyveri*. *J. Bacteriol.* 192, 5115-5123. doi:  
1473 10.1128/JB.00612-10
- 1474 Weghoff, M.C., Bertsch, J., and Müller, V. (2015). A novel mode of lactate metabolism in  
1475 strictly anaerobic bacteria. *Environ. Microbiol.* 17, 670-677. doi: 10.1111/1462-2920
- 1476 Welte, C., and Deppenmeier, U. (2014). Bioenergetics and anaerobic respiratory chains of  
1477 acetoclastic methanogens. *Biochim. Biophys. Acta* 1837, 1130-1147. doi:  
1478 10.1016/j.bbabi.2013.12.002
- 1479 Welte, C., Krätzer, C., and Deppenmeier, U. (2010). Involvement of Ech hydrogenase in  
1480 energy conservation of *Methanosarcina mazei*. *FEBS J.* 277, 3396-3403. doi:  
1481 10.1111/j.1742-4658.2010.07744.x
- 1482 Westphal, L., Wiechmann, A., Baker, J., Minton, N.P., and Müller, V. (2018). The Rnf  
1483 complex is an energy coupled transhydrogenase essential to reversibly link cellular  
1484 NADH and ferredoxin pools in the acetogen *Acetobacterium woodii*. *J. Bacteriol.* 200,  
1485 e00357-00318. doi: 10.1128/JB.00357-18
- 1486 White, H., Strobl, G., Feicht, R., and Simon, H. (1989). Carboxylic acid reductase: a new  
1487 tungsten enzyme catalyses the reduction of non-activated carboxylic acids to  
1488 aldehydes. *Eur. J. Biochem.* 184, 89-96. doi: 10.1111/j.1432-1033.1989.tb14993.x
- 1489 Wiechmann, A., Ciurus, S., Oswald, F., Seiler, V.N., and Müller, V. (2020). It does not  
1490 always take two to tango: “Syntrophy” *via* hydrogen cycling in one bacterial cell.  
1491 *ISME J.* 14, 1561-1570 doi: 10.1038/s41396-020-0627-1
- 1492 Winter, J.U., and Wolfe, R.S. (1980). Methane formation from fructose by syntrophic  
1493 associations of *Acetobacterium woodii* and different strains of methanogens. *Arch.*  
1494 *Microbiol.* 124, 73-79. doi: 10.1007/BF00407031
- 1495 Wohlfarth, G., and Diekert, G. (1991). Thermodynamics of methylenetetrahydrofolate  
1496 reduction to methyltetrahydrofolate and its implications for the energy metabolism of  
1497 homoacetogenic bacteria. *Arch. Microbiol.* 155, 378-381. doi: 10.1007/BF00243458
- 1498 Wolfe, R.S. (1971). Microbial formation of methane. *Adv. Microbiol. Physiol.* 6, 107-146.  
1499 doi: 10.1016/S0065-2911(08)60068-5
- 1500 Wood, H.G., Ragsdale, S.W., and Pezacka, E. (1986). The acetyl-CoA pathway of autotrophic  
1501 growth. *FEMS Microbiol. Rev.* 39, 345-362. doi: 10.1016/0378-1097(86)90022-4

- 1502 Yakushi, T., and Matsushita, K. (2010). Alcohol dehydrogenase of acetic acid bacteria:  
1503 structure, mode of action, and applications in biotechnology. *Appl. Microbiol.*  
1504 *Biotechnol.* 86, 1257-1265. doi: 10.1007/s00253-010-2529-z
- 1505 Yamamoto, I., Saiki, T., Liu, S.M., and Ljungdahl, L.G. (1983). Purification and properties of  
1506 NADP-dependent formate dehydrogenase from *Clostridium thermoaceticum*, a  
1507 tungsten-selenium-iron protein. *J. Biol. Chem.* 258, 1826-1832.
- 1508 Zehnder, A.J., and Brock, T.D. (1979). Methane formation and methane oxidation by  
1509 methanogenic bacteria. *J. Bacteriol.* 137, 420-432. doi: 10.1128/JB.137.1.420-  
1510 432.1979
- 1511 Zellner, G., and Winter, J. (1987). Secondary alcohols as hydrogen donors for CO<sub>2</sub>-reduction  
1512 by methanogens. *FEMS Microbiol. Lett.* 44, 323-328. doi: 10.1111/j.1574-  
1513 6968.1987.tb02309.x
- 1514

Provisional

Figure 01.TIF




Figure 02.TIF




Figure 03.TIF






Figure 04.TIF




Figure 05.TIF




Figure 06.TIF




Figure 07.TIF

