

Covers of sets of groups

Peter J. Cameron, University of St Andrews



Algebra and Combinatorics Seminar
18 January 2024

Cauchy's theorem for the prime 6

Cauchy's theorem says that, if a prime p divides the order of a finite group G , then G contains a subgroup isomorphic to C_p (the cyclic group of order p).

Cauchy's theorem for the prime 6

Cauchy's theorem says that, if a prime p divides the order of a finite group G , then G contains a subgroup isomorphic to C_p (the cyclic group of order p).

One of our new results is the following:

Theorem

If 6 divides the order of a finite group G , then G contains a subgroup isomorphic to C_6 , D_6 (the dihedral group of order 6), or A_4 (the alternating group of degree 4).

Cauchy's theorem for the prime 6

Cauchy's theorem says that, if a prime p divides the order of a finite group G , then G contains a subgroup isomorphic to C_p (the cyclic group of order p).

One of our new results is the following:

Theorem

If 6 divides the order of a finite group G , then G contains a subgroup isomorphic to C_6 , D_6 (the dihedral group of order 6), or A_4 (the alternating group of degree 4).

More about this later ...

The authors

There are still elementary questions about finite groups which have not yet been explored.

The authors

There are still elementary questions about finite groups which have not yet been explored.

This research began on 16 October 2023, when Hamid Reza Dorbidi (Jiroft) sent me a preliminary manuscript. We worked on it, and put a paper on the arXiv the following month (2311.16562).

The authors

There are still elementary questions about finite groups which have not yet been explored.

This research began on 16 October 2023, when Hamid Reza Dorbidi (Jiroft) sent me a preliminary manuscript. We worked on it, and put a paper on the arXiv the following month (2311.16562).

David Craven (Birmingham) and Benjamin Sambale (Heidelberg) saw it on the arXiv and contributed a number of improvements, so we invited them to join the team.

The authors

There are still elementary questions about finite groups which have not yet been explored.

This research began on 16 October 2023, when Hamid Reza Dorbidi (Jiroft) sent me a preliminary manuscript. We worked on it, and put a paper on the arXiv the following month (2311.16562).

David Craven (Birmingham) and Benjamin Sambale (Heidelberg) saw it on the arXiv and contributed a number of improvements, so we invited them to join the team. We are about to submit a revised version to the arXiv.

The problem

There has been a great deal of interest among finite group theorists in describing the subgroups of a given group, in particular the maximal subgroups. Colva's book (with John Bray and Derek Holt) is an important example.

The problem

There has been a great deal of interest among finite group theorists in describing the subgroups of a given group, in particular the maximal subgroups. Colva's book (with John Bray and Derek Holt) is an important example.

Dorbidi's question was a kind of "inverse problem". Given a finite set \mathcal{F} of finite groups, call a group G a **cover** of \mathcal{F} if every group in \mathcal{F} can be embedded in G . What can we say about covers?

The problem

There has been a great deal of interest among finite group theorists in describing the subgroups of a given group, in particular the maximal subgroups. Colva's book (with John Bray and Derek Holt) is an important example.

Dorbidi's question was a kind of "inverse problem". Given a finite set \mathcal{F} of finite groups, call a group G a **cover** of \mathcal{F} if every group in \mathcal{F} can be embedded in G . What can we say about covers?

In particular, let us call an \mathcal{F} -cover G cover **minimal** if no proper subgroup of G is an \mathcal{F} -cover, and **minimum** if no group of smaller order is an \mathcal{F} -cover. We are particularly interested in minimal and minimum \mathcal{F} -covers.

The problem

There has been a great deal of interest among finite group theorists in describing the subgroups of a given group, in particular the maximal subgroups. Colva's book (with John Bray and Derek Holt) is an important example.

Dorbidi's question was a kind of "inverse problem". Given a finite set \mathcal{F} of finite groups, call a group G a **cover** of \mathcal{F} if every group in \mathcal{F} can be embedded in G . What can we say about covers?

In particular, let us call an \mathcal{F} -cover G cover **minimal** if no proper subgroup of G is an \mathcal{F} -cover, and **minimum** if no group of smaller order is an \mathcal{F} -cover. We are particularly interested in minimal and minimum \mathcal{F} -covers.

For any set \mathcal{F} , there is a minimal \mathcal{F} -cover: take the direct product of the groups in \mathcal{F} (this is a cover), and take a subgroup minimal with respect to embedding all the groups in \mathcal{F} .

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

- ▶ How large is a minimum \mathcal{F} -cover?

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

- ▶ How large is a minimum \mathcal{F} -cover?
- ▶ If all the groups in \mathcal{F} have some property \mathcal{P} , is there at least one minimum \mathcal{F} -cover which has property \mathcal{P} ?

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

- ▶ How large is a minimum \mathcal{F} -cover?
- ▶ If all the groups in \mathcal{F} have some property \mathcal{P} , is there at least one minimum \mathcal{F} -cover which has property \mathcal{P} ?
- ▶ Clearly \mathcal{F} has only finitely many minimum covers. Can it have infinitely many minimal covers?

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

- ▶ How large is a minimum \mathcal{F} -cover?
- ▶ If all the groups in \mathcal{F} have some property \mathcal{P} , is there at least one minimum \mathcal{F} -cover which has property \mathcal{P} ?
- ▶ Clearly \mathcal{F} has only finitely many minimum covers. Can it have infinitely many minimal covers?
- ▶ Given a positive integer n , what is the size of a minimum $\mathcal{F}(n)$ -cover, where $\mathcal{F}(n)$ consists of all the groups of order n ? (I will say simply an *n -cover*.)

Some guiding problems

Given a finite set \mathcal{F} of finite groups:

- ▶ How large is a minimum \mathcal{F} -cover?
- ▶ If all the groups in \mathcal{F} have some property \mathcal{P} , is there at least one minimum \mathcal{F} -cover which has property \mathcal{P} ?
- ▶ Clearly \mathcal{F} has only finitely many minimum covers. Can it have infinitely many minimal covers?
- ▶ Given a positive integer n , what is the size of a minimum $\mathcal{F}(n)$ -cover, where $\mathcal{F}(n)$ consists of all the groups of order n ? (I will say simply an *n -cover*.)

I will say something about each of these questions.

Preliminary results

Cayley's Theorem tells us that

Theorem

The symmetric group S_n is an n -cover.

Preliminary results

Cayley's Theorem tells us that

Theorem

The symmetric group S_n is an n -cover.

Also Sylow's theorem tells us that

Theorem

Let $N = p_1^{a_1} \cdots p_r^{a_r}$, where p_1, \dots, p_r are distinct primes. Then, for each i , the Sylow p_i -subgroup of S_n is a $p_i^{a_i}$ -cover.

Preliminary results

Cayley's Theorem tells us that

Theorem

The symmetric group S_n is an n -cover.

Also Sylow's theorem tells us that

Theorem

Let $N = p_1^{a_1} \cdots p_r^{a_r}$, where p_1, \dots, p_r are distinct primes. Then, for each i , the Sylow p_i -subgroup of S_n is a $p_i^{a_i}$ -cover.

In particular, if $n = p^m$ with p prime, then the Sylow p -subgroup of S_n is a p^m -cover, of order $p^{(p^m-1)/(p-1)}$.

Prime power order

This upper bound for the order of a p^m -cover is doubly exponential in m . By contrast, we have:

Prime power order

This upper bound for the order of a p^m -cover is doubly exponential in m . By contrast, we have:

Theorem

The order of a minimum p^m -cover (for p prime) is at least $p^{(2/27+o(1))m^2}$.

Prime power order

This upper bound for the order of a p^m -cover is doubly exponential in m . By contrast, we have:

Theorem

The order of a minimum p^m -cover (for p prime) is at least $p^{(2/27+o(1))m^2}$.

The proof uses Sims' result for the number of groups of order p^m ; a p -group embedding them all must be fairly large.

Prime power order

This upper bound for the order of a p^m -cover is doubly exponential in m . By contrast, we have:

Theorem

The order of a minimum p^m -cover (for p prime) is at least $p^{(2/27+o(1))m^2}$.

The proof uses Sims' result for the number of groups of order p^m ; a p -group embedding them all must be fairly large.

Question

Is the order of a minimum p^m -cover of the form $p^{f(m)}$ where f is polynomial?

Prime power order

This upper bound for the order of a p^m -cover is doubly exponential in m . By contrast, we have:

Theorem

The order of a minimum p^m -cover (for p prime) is at least $p^{(2/27+o(1))m^2}$.

The proof uses Sims' result for the number of groups of order p^m ; a p -group embedding them all must be fairly large.

Question

Is the order of a minimum p^m -cover of the form $p^{f(m)}$ where f is polynomial?

We know that the values for $p^m = 2^3, 2^4, p^3$ (p an odd prime) are respectively $2^5, 2^8$, and p^6 respectively.

Two primes

Let q and r be distinct primes. Does $\{C_q, C_r\}$ have finitely or infinitely many minimal covers?

Two primes

Let q and r be distinct primes. Does $\{C_q, C_r\}$ have finitely or infinitely many minimal covers?

For $\{q, r\} = \{2, 3\}$, a lengthy argument shows that the only minimal covers are C_6 , D_6 and A_4 . This is the theorem with which I began.

Two primes

Let q and r be distinct primes. Does $\{C_q, C_r\}$ have finitely or infinitely many minimal covers?

For $\{q, r\} = \{2, 3\}$, a lengthy argument shows that the only minimal covers are C_6 , D_6 and A_4 . This is the theorem with which I began.

If q and r are odd and at least one is greater than 5, then there are infinitely many minimal covers.

Two primes

Let q and r be distinct primes. Does $\{C_q, C_r\}$ have finitely or infinitely many minimal covers?

For $\{q, r\} = \{2, 3\}$, a lengthy argument shows that the only minimal covers are C_6 , D_6 and A_4 . This is the theorem with which I began.

If q and r are odd and at least one is greater than 5, then there are infinitely many minimal covers.

For, by Dirichlet's Theorem, there are infinitely many primes p congruent to 1 (mod q) and to -1 (mod r). Then $G = \text{PSL}(2, p)$ is a $\{C_q, C_r\}$ -cover. If $r > 5$, then the only maximal subgroup of G containing C_r is D_{p+1} , which does not contain C_q . The argument is similar in the other case.

Two primes

Let q and r be distinct primes. Does $\{C_q, C_r\}$ have finitely or infinitely many minimal covers?

For $\{q, r\} = \{2, 3\}$, a lengthy argument shows that the only minimal covers are C_6 , D_6 and A_4 . This is the theorem with which I began.

If q and r are odd and at least one is greater than 5, then there are infinitely many minimal covers.

For, by Dirichlet's Theorem, there are infinitely many primes p congruent to 1 (mod q) and to -1 (mod r). Then $G = \text{PSL}(2, p)$ is a $\{C_q, C_r\}$ -cover. If $r > 5$, then the only maximal subgroup of G containing C_r is D_{p+1} , which does not contain C_q . The argument is similar in the other case.

Remaining pairs of primes have not yet been settled.

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

- ▶ If G is soluble, then it is one of three possibilities: C_{qr} , an elementary abelian q -group with C_r acting irreducibly on it, or an elementary abelian r -group with C_q acting irreducibly on it.

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

- ▶ If G is soluble, then it is one of three possibilities: C_{qr} , an elementary abelian q -group with C_r acting irreducibly on it, or an elementary abelian r -group with C_q acting irreducibly on it.
- ▶ Otherwise, G has a unique maximal normal subgroup N , and G/N is a minimal $\{C_q, C_r\}$ -cover.

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

- ▶ If G is soluble, then it is one of three possibilities: C_{qr} , an elementary abelian q -group with C_r acting irreducibly on it, or an elementary abelian r -group with C_q acting irreducibly on it.
- ▶ Otherwise, G has a unique maximal normal subgroup N , and G/N is a minimal $\{C_q, C_r\}$ -cover.
- ▶ Now take $\{q, r\} = \{2, 3\}$.
In 1977 (pre-CFSG), Podufalov showed that a simple group with no element of order 6 must be $\text{PSL}(2, q)$, $\text{PSL}(3, q)$, $\text{PSU}(3, q)$ or $\text{Sz}(q)$ for some prime power q .

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

- ▶ If G is soluble, then it is one of three possibilities: C_{qr} , an elementary abelian q -group with C_r acting irreducibly on it, or an elementary abelian r -group with C_q acting irreducibly on it.
- ▶ Otherwise, G has a unique maximal normal subgroup N , and G/N is a minimal $\{C_q, C_r\}$ -cover.
- ▶ Now take $\{q, r\} = \{2, 3\}$.
In 1977 (pre-CFSG), Podufalov showed that a simple group with no element of order 6 must be $\text{PSL}(2, q)$, $\text{PSL}(3, q)$, $\text{PSU}(3, q)$ or $\text{Sz}(q)$ for some prime power q .
- ▶ Suzuki groups can't occur since their orders are not divisible by 3. The others all involve $\text{PSL}(2, p)$, where q is a power of p .

Sketch proof

Let G be a minimal $\{C_q, C_r\}$ -cover.

- ▶ If G is soluble, then it is one of three possibilities: C_{qr} , an elementary abelian q -group with C_r acting irreducibly on it, or an elementary abelian r -group with C_q acting irreducibly on it.
- ▶ Otherwise, G has a unique maximal normal subgroup N , and G/N is a minimal $\{C_q, C_r\}$ -cover.
- ▶ Now take $\{q, r\} = \{2, 3\}$.
In 1977 (pre-CFSG), Podufalov showed that a simple group with no element of order 6 must be $\text{PSL}(2, q)$, $\text{PSL}(3, q)$, $\text{PSU}(3, q)$ or $\text{Sz}(q)$ for some prime power q .
- ▶ Suzuki groups can't occur since their orders are not divisible by 3. The others all involve $\text{PSL}(2, p)$, where q is a power of p .
- ▶ $\text{PSL}(2, 3) \cong A_4$, while for other p they have D_6 as a subgroup.

Minimal covers

It is not too hard to show that any minimal 2^2 -cover has order 2^3 (and indeed, there are just two, namely $C_4 \times C_2$ and D_8), so that they are minimum covers.

Minimal covers

It is not too hard to show that any minimal 2^2 -cover has order 2^3 (and indeed, there are just two, namely $C_4 \times C_2$ and D_8), so that they are minimum covers.

However, we have:

Theorem

There are infinitely many minimal 2^3 -covers. Indeed, for any $m \geq 4$, the group $SD_{2^m} \times C_2$ is a minimal 2^3 -cover.

Minimal covers

It is not too hard to show that any minimal 2^2 -cover has order 2^3 (and indeed, there are just two, namely $C_4 \times C_2$ and D_8), so that they are minimum covers.

However, we have:

Theorem

There are infinitely many minimal 2^3 -covers. Indeed, for any $m \geq 4$, the group $SD_{2^m} \times C_2$ is a minimal 2^3 -cover.

Here SD_{2^m} is the **semi-dihedral group**

$$\langle a, b : a^{2^{m-1}} = b^2 = 1, b^{-1}ab = a^{2^{m-2}-1} \rangle.$$

Nilpotent groups

Theorem

Let \mathcal{F} be a set of finite nilpotent groups. Then there is a minimum \mathcal{F} -cover which is nilpotent.

Nilpotent groups

Theorem

Let \mathcal{F} be a set of finite nilpotent groups. Then there is a minimum \mathcal{F} -cover which is nilpotent.

To see this, just take the direct product of \mathcal{F}_p -covers, where \mathcal{F}_p is the set of Sylow p -subgroups of the groups in \mathcal{F} for all relevant primes.

Nilpotent groups

Theorem

Let \mathcal{F} be a set of finite nilpotent groups. Then there is a minimum \mathcal{F} -cover which is nilpotent.

To see this, just take the direct product of \mathcal{F}_p -covers, where \mathcal{F}_p is the set of Sylow p -subgroups of the groups in \mathcal{F} for all relevant primes.

Of course we cannot say that all minimum \mathcal{F} -covers are nilpotent. For example, if $\mathcal{F} = \{(C_2)^2, C_3, C_5\}$, then a minimum \mathcal{F} -cover has order 60, and any group of order 60 having these as its Sylow subgroups is an \mathcal{F} -cover, including the simple group A_5 .

Abelian groups

For abelian groups, the position is very different. We can work out the order of the smallest abelian cover of any set of finite abelian groups; but we cannot prove that at least one minimum cover must be abelian!

Abelian groups

For abelian groups, the position is very different. We can work out the order of the smallest abelian cover of any set of finite abelian groups; but we cannot prove that at least one minimum cover must be abelian!

Arguing as in the last theorem, it suffices to find the smallest cover of a set \mathcal{F} of abelian p -groups, where p is prime.

Abelian groups

For abelian groups, the position is very different. We can work out the order of the smallest abelian cover of any set of finite abelian groups; but we cannot prove that at least one minimum cover must be abelian!

Arguing as in the last theorem, it suffices to find the smallest cover of a set \mathcal{F} of abelian p -groups, where p is prime.

Any such group has a canonical form

$$G = C_{p^{a_1}} \times \cdots \times C_{p^{a_r}}$$

where $a_1 \geq \cdots \geq a_r$. Now given a set of finite abelian p -groups, we can write them all in canonical form and assume that the value of r is the same for each (by adding trivial factors if necessary). Then the smallest abelian cover has canonical form whose i th factor is the largest group occurring as the i th factor of one of the groups in \mathcal{F} .

As a corollary, we can show the following.

As a corollary, we can show the following.
Define a function f on the natural numbers by

$$f(n) = \sum_{k=1}^n \left\lfloor \frac{n}{k} \right\rfloor.$$

As a corollary, we can show the following.

Define a function f on the natural numbers by

$$f(n) = \sum_{k=1}^n \left\lfloor \frac{n}{k} \right\rfloor.$$

This function was considered by Dirichlet, who showed that $f(n) = n(\log n + 2\gamma - 1) + O(\sqrt{n})$, where γ is the **Euler–Mascheroni constant**. It is given as sequence A006218 in the On-Line Encyclopedia of Integer Sequences, where a number of occurrences of it are noted. But the one given here seems to be new.

As a corollary, we can show the following.

Define a function f on the natural numbers by

$$f(n) = \sum_{k=1}^n \left\lfloor \frac{n}{k} \right\rfloor.$$

This function was considered by Dirichlet, who showed that $f(n) = n(\log n + 2\gamma - 1) + O(\sqrt{n})$, where γ is the **Euler–Mascheroni constant**. It is given as sequence A006218 in the On-Line Encyclopedia of Integer Sequences, where a number of occurrences of it are noted. But the one given here seems to be new.

Theorem

The smallest order of an abelian group containing every abelian group of order p^n is $p^{f(n)}$.

As a corollary, we can show the following.

Define a function f on the natural numbers by

$$f(n) = \sum_{k=1}^n \left\lfloor \frac{n}{k} \right\rfloor.$$

This function was considered by Dirichlet, who showed that $f(n) = n(\log n + 2\gamma - 1) + O(\sqrt{n})$, where γ is the **Euler–Mascheroni constant**. It is given as sequence A006218 in the On-Line Encyclopedia of Integer Sequences, where a number of occurrences of it are noted. But the one given here seems to be new.

Theorem

The smallest order of an abelian group containing every abelian group of order p^n is $p^{f(n)}$.

This is proved by applying the preceding theorem, first computing that the largest size of the k th component in the canonical form of an abelian group of order p^n is $p^{\lfloor n/k \rfloor}$.

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).
- ▶ Abelian: Don't know.

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).
- ▶ Abelian: Don't know.
- ▶ Nilpotent: Yes (we saw this earlier).

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).
- ▶ Abelian: Don't know.
- ▶ Nilpotent: Yes (we saw this earlier).
- ▶ Soluble: No (proof coming up).

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).
- ▶ Abelian: Don't know.
- ▶ Nilpotent: Yes (we saw this earlier).
- ▶ Soluble: No (proof coming up).
- ▶ Simple: See later.

Questions and answers

Let \mathcal{P} be a property of finite groups. If \mathcal{F} is a finite set of finite groups, does there exist a minimum \mathcal{F} -cover with property \mathcal{P} ?

- ▶ Cyclic: Yes (the cyclic group whose order is the least common multiple of the groups in \mathcal{F} does the job).
- ▶ Abelian: Don't know.
- ▶ Nilpotent: Yes (we saw this earlier).
- ▶ Soluble: No (proof coming up).
- ▶ Simple: See later.

For soluble groups, take $\mathcal{F} = \{D_{10}, A_4\}$. The least common multiple of their orders is 60, and it is an easy exercise to show that the only group of order 60 containing both is A_5 .

Simple groups

Theorem

Let M and N be simple groups. Then a minimum cover of $\{M, N\}$ is either simple or $M \times N$.

Simple groups

Theorem

Let M and N be simple groups. Then a minimum cover of $\{M, N\}$ is either simple or $M \times N$.

Both cases occur:

Example Let $M = A_5$ and $N = \text{PSL}(2, 8)$. The orders of these groups are 60 and 504. Their least common multiple is 2520 and their product is 30240. The only simple groups with order divisible by 2520 and not greater than 30240 are A_7 , A_8 and $\text{PSL}(3, 4)$; none of these embed $\text{PSL}(2, 8)$. By the theorem, the unique minimum $\{M, N\}$ -cover is $M \times N$.

Simple groups

Theorem

Let M and N be simple groups. Then a minimum cover of $\{M, N\}$ is either simple or $M \times N$.

Both cases occur:

Example Let $M = A_5$ and $N = \text{PSL}(2, 8)$. The orders of these groups are 60 and 504. Their least common multiple is 2520 and their product is 30240. The only simple groups with order divisible by 2520 and not greater than 30240 are A_7 , A_8 and $\text{PSL}(3, 4)$; none of these embed $\text{PSL}(2, 8)$. By the theorem, the unique minimum $\{M, N\}$ -cover is $M \times N$.

Example Let $M = A_6$ and $N = \text{PSL}(2, 7)$. Their orders are 360 and 168, with least common multiple 2520. There is a unique simple group of order 2520, namely A_7 , which embeds both M and N ; so A_7 is the unique minimum $\{M, N\}$ -cover.

Can a pair of simple groups have more than one minimum cover?

Can a pair of simple groups have more than one minimum cover?

There are two possibilities: Either two simple groups of the same order are minimum covers, or $\{M, N\}$ has minimum covers L and $M \times N$.

Can a pair of simple groups have more than one minimum cover?

There are two possibilities: Either two simple groups of the same order are minimum covers, or $\{M, N\}$ has minimum covers L and $M \times N$.

In the former case, we know that a pair of simple groups of the same order must be either $\{A_8, \text{PSL}(3, 4)\}$, or $\{\text{PSp}(2m, q), \text{P}\Omega(2m + 1, q)\}$ with $m \geq 3$ and q odd.

Can a pair of simple groups have more than one minimum cover?

There are two possibilities: Either two simple groups of the same order are minimum covers, or $\{M, N\}$ has minimum covers L and $M \times N$.

In the former case, we know that a pair of simple groups of the same order must be either $\{A_8, \text{PSL}(3, 4)\}$, or $\{\text{PSp}(2m, q), \text{P}\Omega(2m + 1, q)\}$ with $m \geq 3$ and q odd.

In the latter case, there are infinitely many triples of simple groups $\{L, M, N\}$ with $|L| = |M| \cdot |N|$. (The smallest is $\{A_6, \text{PSL}(2, 8), A_9\}$.) Can we determine all such triples? And can both L and $M \times N$ be minimum covers of the same pair of simple groups?

Radicals and residuals

The answer to our earlier question is "Yes" for classes of groups not having normal subgroups of certain types, or quotients of certain types.

Radicals and residuals

The answer to our earlier question is “Yes” for classes of groups not having normal subgroups of certain types, or quotients of certain types.

Theorem

Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, none of which has a non-trivial \mathcal{X} -group as a quotient, and let G be a minimal \mathcal{F} -cover. Then G has no non-trivial \mathcal{X} -group as a quotient.

Radicals and residuals

The answer to our earlier question is “Yes” for classes of groups not having normal subgroups of certain types, or quotients of certain types.

Theorem

Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, none of which has a non-trivial \mathcal{X} -group as a quotient, and let G be a minimal \mathcal{F} -cover. Then G has no non-trivial \mathcal{X} -group as a quotient.

Proof.

Suppose that $G/N \in \mathcal{X}$. Then for any group $H \in \mathcal{F}$, $H/H \cap N \cong HN/N \leq G/N \in \mathcal{X}$. So $H/H \cap N \in \mathcal{X}$ which implies $H \subseteq N$. By minimality of G , we have $N = G$. □

Applications

- ▶ Let \mathcal{X} be the class of finite abelian groups. The condition that G has no non-trivial homomorphism to an \mathcal{X} -group means that G is perfect. So we deduce that, if every group in \mathcal{F} is perfect, then any minimal \mathcal{F} -cover is perfect.

Applications

- ▶ Let \mathcal{X} be the class of finite abelian groups. The condition that G has no non-trivial homomorphism to an \mathcal{X} -group means that G is perfect. So we deduce that, if every group in \mathcal{F} is perfect, then any minimal \mathcal{F} -cover is perfect.
- ▶ Let \mathcal{X} be the class of finite solvable groups. The condition that G has no non-trivial homomorphism to an \mathcal{X} -group means that G is equal to its solvable residual. So, if every group in \mathcal{F} is equal to its solvable residual, then the same is true of any minimal \mathcal{F} -cover.

A dual result

Let us call an \mathcal{F} -cover G **co-minimal** if no quotient of G is an \mathcal{F} -cover. Note that a minimum cover is both minimal and co-minimal.

A dual result

Let us call an \mathcal{F} -cover G **co-minimal** if no quotient of G is an \mathcal{F} -cover. Note that a minimum cover is both minimal and co-minimal.

Theorem

Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, and suppose that no group in \mathcal{F} has a non-trivial normal \mathcal{X} -subgroup. Let G be a co-minimal cover of \mathcal{F} . Then G has no non-trivial normal \mathcal{X} -subgroup.

A dual result

Let us call an \mathcal{F} -cover G **co-minimal** if no quotient of G is an \mathcal{F} -cover. Note that a minimum cover is both minimal and co-minimal.

Theorem

Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, and suppose that no group in \mathcal{F} has a non-trivial normal \mathcal{X} -subgroup. Let G be a co-minimal cover of \mathcal{F} . Then G has no non-trivial normal \mathcal{X} -subgroup.

Applications:

- ▶ If no group in \mathcal{F} has a non-trivial abelian normal subgroup, then a co-minimal cover has no non-trivial abelian normal subgroup.

A dual result

Let us call an \mathcal{F} -cover G **co-minimal** if no quotient of G is an \mathcal{F} -cover. Note that a minimum cover is both minimal and co-minimal.

Theorem

Suppose that \mathcal{X} is a subgroup-closed class of finite groups. Let \mathcal{F} be a finite set of finite groups, and suppose that no group in \mathcal{F} has a non-trivial normal \mathcal{X} -subgroup. Let G be a co-minimal cover of \mathcal{F} . Then G has no non-trivial normal \mathcal{X} -subgroup.

Applications:

- ▶ If no group in \mathcal{F} has a non-trivial abelian normal subgroup, then a co-minimal cover has no non-trivial abelian normal subgroup.
- ▶ Similarly with “soluble” replacing “abelian”.

A dual problem

We can dualise the entire set-up. Given a finite set \mathcal{F} of finite groups, say that G is a **dual cover** for \mathcal{F} if every group in \mathcal{F} is a homomorphic image of G .

A dual problem

We can dualise the entire set-up. Given a finite set \mathcal{F} of finite groups, say that G is a **dual cover** for \mathcal{F} if every group in \mathcal{F} is a homomorphic image of G .

Dualising the earlier concepts, we say that a dual cover G is **minimal** if no proper homomorphic image of G is a dual cover, and **minimum** if no dual cover is smaller than G .

A dual problem

We can dualise the entire set-up. Given a finite set \mathcal{F} of finite groups, say that G is a **dual cover** for \mathcal{F} if every group in \mathcal{F} is a homomorphic image of G .

Dualising the earlier concepts, we say that a dual cover G is **minimal** if no proper homomorphic image of G is a dual cover, and **minimum** if no dual cover is smaller than G .

Because of duality for abelian groups, the results for these groups are the same as those for covers described earlier.

A dual problem

We can dualise the entire set-up. Given a finite set \mathcal{F} of finite groups, say that G is a **dual cover** for \mathcal{F} if every group in \mathcal{F} is a homomorphic image of G .

Dualising the earlier concepts, we say that a dual cover G is **minimal** if no proper homomorphic image of G is a dual cover, and **minimum** if no dual cover is smaller than G .

Because of duality for abelian groups, the results for these groups are the same as those for covers described earlier. But for non-abelian groups, essentially nothing is known.

Some problems

Question

Is it true that a finite set of finite abelian groups has a minimum cover which is abelian?

Some problems

Question

Is it true that a finite set of finite abelian groups has a minimum cover which is abelian?

Question

What is the order of magnitude of the function F such that a minimum p^m -cover has order $p^{F(m)}$? Is it polynomial, or exponential?

Some problems

Question

Is it true that a finite set of finite abelian groups has a minimum cover which is abelian?

Question

What is the order of magnitude of the function F such that a minimum p^m -cover has order $p^{F(m)}$? Is it polynomial, or exponential?

Question

Develop the theory of dual covers.

Some problems

Question

Is it true that a finite set of finite abelian groups has a minimum cover which is abelian?

Question

What is the order of magnitude of the function F such that a minimum p^m -cover has order $p^{F(m)}$? Is it polynomial, or exponential?

Question

Develop the theory of dual covers.

Question

What happens for semigroups?



... for your attention.