


An algebraist looks at discrete mathematics

Peter J. Cameron, University of St Andrews



ADMA Colloquium Lecture, 18 May 2024



Summary

For most of my career, I have worked mostly in discrete mathematics.

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

I want to tell you three stories about places where algebra and discrete mathematics have come together profitably. I had some part in all three of these. The topics are

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

I want to tell you three stories about places where algebra and discrete mathematics have come together profitably. I had some part in all three of these. The topics are

1. Root systems and graph spectra.

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

I want to tell you three stories about places where algebra and discrete mathematics have come together profitably. I had some part in all three of these. The topics are

1. Root systems and graph spectra.
2. The countable random graph and the Urysohn space.

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

I want to tell you three stories about places where algebra and discrete mathematics have come together profitably. I had some part in all three of these. The topics are

1. Root systems and graph spectra.
2. The countable random graph and the Urysohn space.
3. A beautiful graph from the Mathieu group M_{11} .

Summary

For most of my career, I have worked mostly in discrete mathematics.

But I began life as an algebraist.

I want to tell you three stories about places where algebra and discrete mathematics have come together profitably. I had some part in all three of these. The topics are

1. Root systems and graph spectra.
2. The countable random graph and the Urysohn space.
3. A beautiful graph from the Mathieu group M_{11} .

I can't say much about each of these but I hope to give you a little taste. The topics will be separated by pictures of bridges; these tell you that, if you have lost the thread, you can now pick up a new thread.

Forth bridge, Scotland



1. Graph spectra and root systems

In the 1950s and 1960s, there was a lot of work by Hoffman, Ray-Chaudhuri, Shrikhande, Chang, Seidel, and others about this class of graphs.

1. Graph spectra and root systems

In the 1950s and 1960s, there was a lot of work by Hoffman, Ray-Chaudhuri, Shrikhande, Chang, Seidel, and others about this class of graphs.

Hoffman formulated a conjecture that a “sufficiently large” (in some sense) graph with smallest eigenvalue -2 or greater was a **generalized line graph**, a class of graphs which he devised. He worked long on this problem but was unable to complete the proof.

1. Graph spectra and root systems

In the 1950s and 1960s, there was a lot of work by Hoffman, Ray-Chaudhuri, Shrikhande, Chang, Seidel, and others about this class of graphs.

Hoffman formulated a conjecture that a “sufficiently large” (in some sense) graph with smallest eigenvalue -2 or greater was a **generalized line graph**, a class of graphs which he devised. He worked long on this problem but was unable to complete the proof.

The **line graph** $L(G)$ of a graph G is the graph whose vertices are the edges of G , two vertices of $L(G)$ joined if the corresponding edges of G share a vertex.

1. Graph spectra and root systems

In the 1950s and 1960s, there was a lot of work by Hoffman, Ray-Chaudhuri, Shrikhande, Chang, Seidel, and others about this class of graphs.

Hoffman formulated a conjecture that a “sufficiently large” (in some sense) graph with smallest eigenvalue -2 or greater was a **generalized line graph**, a class of graphs which he devised. He worked long on this problem but was unable to complete the proof.

The **line graph** $L(G)$ of a graph G is the graph whose vertices are the edges of G , two vertices of $L(G)$ joined if the corresponding edges of G share a vertex.

A **cocktail party graph** $CP(n)$ is a graph on $2n$ vertices a_i, b_i with all pairs of vertices joined except a_i and b_i .

1. Graph spectra and root systems

In the 1950s and 1960s, there was a lot of work by Hoffman, Ray-Chaudhuri, Shrikhande, Chang, Seidel, and others about this class of graphs.



Hoffman formulated a conjecture that a “sufficiently large” (in some sense) graph with smallest eigenvalue -2 or greater was a **generalized line graph**, a class of graphs which he devised. He worked long on this problem but was unable to complete the proof.

The **line graph** $L(G)$ of a graph G is the graph whose vertices are the edges of G , two vertices of $L(G)$ joined if the corresponding edges of G share a vertex.



A **cocktail party graph** $CP(n)$ is a graph on $2n$ vertices a_i, b_i with all pairs of vertices joined except a_i and b_i .

A **generalized line graph** is obtained from a line graph by attaching a cocktail party graph associated with each vertex. The next slide shows an example.

A generalized line graph





A generalized line graph



The red part is the line graph $L(G)$; the blue shows the added cocktail party graphs.

A generalized line graph



The red part is the line graph $L(G)$; the blue shows the added cocktail party graphs.



Hoffman's conjecture is true in a strong sense. But the proof (by Jean-Marie Goethals, Jaap Seidel, Ernie Shult, and me) came from a completely different direction, using the theory of **root systems**.

Root systems

Root systems are beautiful geometrical objects, which have been compared to multidimensional crystals.



Root systems

Root systems are beautiful geometrical objects, which have been compared to multidimensional crystals.



Root systems

Root systems are beautiful geometrical objects, which have been compared to multidimensional crystals.



They arose first in the theory of Lie algebras, but now occur everywhere from singularity theory to general relativity, from cluster algebras to finite simple groups.

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

- ▶ if $v \in S$, then $\lambda v \in S$ if and only if $\lambda = \pm 1$;

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

- ▶ if $v \in S$, then $\lambda v \in S$ if and only if $\lambda = \pm 1$;
- ▶ S is closed under reflection in the hyperplane perpendicular to any of its vectors.

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

- ▶ if $v \in S$, then $\lambda v \in S$ if and only if $\lambda = \pm 1$;
- ▶ S is closed under reflection in the hyperplane perpendicular to any of its vectors.

There is another condition often imposed, which makes the root system even more crystal-like by ensuring that the integer combinations of its vectors form a lattice. This will hold in the case of interest here.

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

- ▶ if $v \in S$, then $\lambda v \in S$ if and only if $\lambda = \pm 1$;
- ▶ S is closed under reflection in the hyperplane perpendicular to any of its vectors.

There is another condition often imposed, which makes the root system even more crystal-like by ensuring that the integer combinations of its vectors form a lattice. This will hold in the case of interest here.

The important case for us is when all the vectors in S have the same length. In this case, the angle between any two roots is 60° , 90° , 120° , or 180° .

The definition

A **root system** is a finite set S of vectors spanning the real vector space \mathbb{R}^d and having the properties

- ▶ if $v \in S$, then $\lambda v \in S$ if and only if $\lambda = \pm 1$;
- ▶ S is closed under reflection in the hyperplane perpendicular to any of its vectors.

There is another condition often imposed, which makes the root system even more crystal-like by ensuring that the integer combinations of its vectors form a lattice. This will hold in the case of interest here.

The important case for us is when all the vectors in S have the same length. In this case, the angle between any two roots is 60° , 90° , 120° , or 180° .

We call the root system **indecomposable** if the space is not an orthogonal direct sum of two subspaces which contain all the vectors of S .

The ADE classification

Theorem

An indecomposable root system with all roots of the same length is of one of the types A_d ($d \geq 1$), D_d ($d \geq 4$), E_6 , E_7 , E_8 . (The subscript denotes the dimension.)

The ADE classification

Theorem

An indecomposable root system with all roots of the same length is of one of the types A_d ($d \geq 1$), D_d ($d \geq 4$), E_6 , E_7 , E_8 . (The subscript denotes the dimension.)

The systems are explicitly known and are easy to work with. These particular systems are ubiquitous in mathematics. A forthcoming book “ADE: Patterns in Mathematics” by Pierre Dechant, Yang He, John McKay and me traces some of their many occurrences in different parts of our subject.

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

Here is a sketch proof. Let A be the adjacency matrix. Then $2I + A$ is positive semi-definite, and so is the matrix of inner products of a set of vectors in a real vector space. These vectors all have length $\sqrt{2}$ and any two make angles of 90° or 60° .

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

Here is a sketch proof. Let A be the adjacency matrix. Then $2I + A$ is positive semi-definite, and so is the matrix of inner products of a set of vectors in a real vector space. These vectors all have length $\sqrt{2}$ and any two make angles of 90° or 60° .

A geometric argument shows that such a set can be enlarged to a maximal set with angles 0° , 60° , 90° , 120° or 180° , which is a root system of type A , D or E . So the graph is “represented” in such a root system.

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

Here is a sketch proof. Let A be the adjacency matrix. Then $2I + A$ is positive semi-definite, and so is the matrix of inner products of a set of vectors in a real vector space. These vectors all have length $\sqrt{2}$ and any two make angles of 90° or 60° .

A geometric argument shows that such a set can be enlarged to a maximal set with angles 0° , 60° , 90° , 120° or 180° , which is a root system of type A , D or E . So the graph is “represented” in such a root system.

Now the following observations finish the job:

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

Here is a sketch proof. Let A be the adjacency matrix. Then $2I + A$ is positive semi-definite, and so is the matrix of inner products of a set of vectors in a real vector space. These vectors all have length $\sqrt{2}$ and any two make angles of 90° or 60° .

A geometric argument shows that such a set can be enlarged to a maximal set with angles 0° , 60° , 90° , 120° or 180° , which is a root system of type A , D or E . So the graph is “represented” in such a root system.

Now the following observations finish the job:

- ▶ $A_d \subseteq D_{d+1}$ and $E_6 \subseteq E_7 \subseteq E_8$, so our graph lives in either D_d (for some d) or E_8 ;

The theorem

Theorem

A connected finite graph with smallest eigenvalue -2 or greater is either a generalized line graph or represented by a subset of the root system E_8 .

Here is a sketch proof. Let A be the adjacency matrix. Then $2I + A$ is positive semi-definite, and so is the matrix of inner products of a set of vectors in a real vector space. These vectors all have length $\sqrt{2}$ and any two make angles of 90° or 60° .

A geometric argument shows that such a set can be enlarged to a maximal set with angles 0° , 60° , 90° , 120° or 180° , which is a root system of type A , D or E . So the graph is “represented” in such a root system.

Now the following observations finish the job:

- ▶ $A_d \subseteq D_{d+1}$ and $E_6 \subseteq E_7 \subseteq E_8$, so our graph lives in either D_d (for some d) or E_8 ;
- ▶ a graph represented in D_d is a generalized line graph.

The exceptions

Since E_8 is a finite object, it can only represent finitely many graphs.

The exceptions


Since E_8 is a finite object, it can only represent finitely many graphs.

These include many famous graphs, including the Petersen, Clebsch, Shrikhande, Schläfli and three Chang graphs.

The exceptions

Since E_8 is a finite object, it can only represent finitely many graphs.

These include many famous graphs, including the Petersen, Clebsch, Shrikhande, Schläfli and three Chang graphs.



The Shrikhande graph (thanks to Ambat Vijayakumar)

Howrah bridge, Kolkata



2. The random graph and the Urysohn space

How to choose a random graph? The simplest model: start with the set of vertices, for each pair toss a fair coin to decide whether to join them with an edge.

2. The random graph and the Urysohn space

How to choose a random graph? The simplest model: start with the set of vertices, for each pair toss a fair coin to decide whether to join them with an edge.

In 1959, Erdős and Rényi showed that almost all finite graphs have no non-trivial symmetry, indeed lie “as far as possible” from symmetry.

2. The random graph and the Urysohn space

How to choose a random graph? The simplest model: start with the set of vertices, for each pair toss a fair coin to decide whether to join them with an edge.

In 1959, Erdős and Rényi showed that almost all finite graphs have no non-trivial symmetry, indeed lie “as far as possible” from symmetry.

This began a huge topic, random graphs. But as a tailpiece, they showed that the result is false for countably infinite graphs. With probability 1, a random countable graph has infinitely many automorphisms.

2. The random graph and the Urysohn space

How to choose a random graph? The simplest model: start with the set of vertices, for each pair toss a fair coin to decide whether to join them with an edge.

In 1959, Erdős and Rényi showed that almost all finite graphs have no non-trivial symmetry, indeed lie “as far as possible” from symmetry.

This began a huge topic, random graphs. But as a tailpiece, they showed that the result is false for countably infinite graphs. With probability 1, a random countable graph has infinitely many automorphisms.

The reason for this is even more surprising.

The random graph

Theorem

There exists a countable graph R with the properties:

The random graph

Theorem

There exists a countable graph R with the properties:

- ▶ *R is **universal**: every finite or countable graph is embeddable in R as induced subgraph.*

The random graph

Theorem

There exists a countable graph R with the properties:

- ▶ R is *universal*: every finite or countable graph is embeddable in R as induced subgraph.
- ▶ R is *homogeneous*: every isomorphism between finite induced subgraphs can be extended to an automorphism of R .

The random graph

Theorem

There exists a countable graph R with the properties:

- ▶ R is *universal*: every finite or countable graph is embeddable in R as induced subgraph.
- ▶ R is *homogeneous*: every isomorphism between finite induced subgraphs can be extended to an automorphism of R .
- ▶ With probability 1, a countable random graph is isomorphic to R .

The random graph

Theorem

There exists a countable graph R with the properties:

- ▶ R is *universal*: every finite or countable graph is embeddable in R as induced subgraph.
- ▶ R is *homogeneous*: every isomorphism between finite induced subgraphs can be extended to an automorphism of R .
- ▶ With probability 1, a countable random graph is isomorphic to R .

The Erdős–Rényi proof is a nonconstructive existence proof: if something occurs with probability 1, it certainly occurs. But at almost the same time, Rado constructed a countable universal graph which turned out to be R . His construction was as follows. The vertex set is the set \mathbb{N} of natural numbers (including 0). Given x and y , with $x < y$, write y in base 2: now join x and y if and only if the x -th digit of y is 1.

Fraïssé

In fact, more than ten years earlier, the French logician Roland Fraïssé had given a construction including the graph R as a special case.

Fraïssé

In fact, more than ten years earlier, the French logician Roland Fraïssé had given a construction including the graph R as a special case.

Fraïssé's topic was **relational structures**, sets with a specified collection of relations of prescribed arity. As well as graphs, think digraphs, total or partial orders, hypergraphs, etc.

Fraïssé

In fact, more than ten years earlier, the French logician Roland Fraïssé had given a construction including the graph R as a special case.

Fraïssé's topic was **relational structures**, sets with a specified collection of relations of prescribed arity. As well as graphs, think digraphs, total or partial orders, hypergraphs, etc.

Fraïssé gave a necessary and sufficient condition for a class \mathcal{A} of finite structures to be all the finite structures embeddable (as induced substructure) in a countable homogeneous structure M . Moreover, if M exists, then it is unique. It is now called the **Fraïssé limit** of the class \mathcal{A} .

Fraïssé

In fact, more than ten years earlier, the French logician Roland Fraïssé had given a construction including the graph R as a special case.

Fraïssé's topic was **relational structures**, sets with a specified collection of relations of prescribed arity. As well as graphs, think digraphs, total or partial orders, hypergraphs, etc.

Fraïssé gave a necessary and sufficient condition for a class \mathcal{A} of finite structures to be all the finite structures embeddable (as induced substructure) in a countable homogeneous structure M . Moreover, if M exists, then it is unique. It is now called the **Fraïssé limit** of the class \mathcal{A} .

Thus, the Fraïssé limit of the class of finite graphs is the random graph R ; the Fraïssé limit of the class of finite total orders is $(\mathbb{Q}, <)$.

Fraïssé

In fact, more than ten years earlier, the French logician Roland Fraïssé had given a construction including the graph R as a special case.

Fraïssé's topic was **relational structures**, sets with a specified collection of relations of prescribed arity. As well as graphs, think digraphs, total or partial orders, hypergraphs, etc.

Fraïssé gave a necessary and sufficient condition for a class \mathcal{A} of finite structures to be all the finite structures embeddable (as induced substructure) in a countable homogeneous structure M . Moreover, if M exists, then it is unique. It is now called the **Fraïssé limit** of the class \mathcal{A} .

Thus, the Fraïssé limit of the class of finite graphs is the random graph R ; the Fraïssé limit of the class of finite total orders is $(\mathbb{Q}, <)$.

But even Fraïssé was not the first ...

Urysohn

In 1924, Pavel Alexandrov and Pavel Urysohn, the Soviet pioneers of topology, came to western Europe for discussions with Hilbert, Brouwer, Hausdorff, and others.

Urysohn

In 1924, Pavel Alexandrov and Pavel Urysohn, the Soviet pioneers of topology, came to western Europe for discussions with Hilbert, Brouwer, Hausdorff, and others.

After this, the two went to south-west France for a holiday. They swam in the sea every morning. One morning there was a severe storm and they were advised not to swim, but went in anyway. Alexandrov came back; Urysohn did not. He was 26.

Urysohn

In 1924, Pavel Alexandrov and Pavel Urysohn, the Soviet pioneers of topology, came to western Europe for discussions with Hilbert, Brouwer, Hausdorff, and others.

After this, the two went to south-west France for a holiday. They swam in the sea every morning. One morning there was a severe storm and they were advised not to swim, but went in anyway. Alexandrov came back; Urysohn did not. He was 26. He is known for several results in the foundations of topology, but the theorem I will talk about was less well known; it was published posthumously from the papers he left.

Urysohn

In 1924, Pavel Alexandrov and Pavel Urysohn, the Soviet pioneers of topology, came to western Europe for discussions with Hilbert, Brouwer, Hausdorff, and others.

After this, the two went to south-west France for a holiday. They swam in the sea every morning. One morning there was a severe storm and they were advised not to swim, but went in anyway. Alexandrov came back; Urysohn did not. He was 26. He is known for several results in the foundations of topology, but the theorem I will talk about was less well known; it was published posthumously from the papers he left.

In 2000, I spoke about the random graph at the European Congress of Mathematics in Barcelona. After the talk, I was approached by someone who introduced himself as Anatoly Vershik, and asked if I knew about the Urysohn space. This is a **Polish space** (a complete second-countable metric space) which is universal and homogeneous.

The Urysohn space

We cannot apply Fraïssé directly: there are too many finite metric spaces (uncountably many two-point spaces).

The Urysohn space

We cannot apply Fraïssé directly: there are too many finite metric spaces (uncountably many two-point spaces).

Instead, we define a metric space to be **rational** if all distances are rational numbers. The class of rational metric spaces satisfies Fraïssé's theorem; its Fraïssé limit is the **rational Urysohn space**. Taking its completion in the usual sense (add limits for Cauchy sequences) gives the real Urysohn space.

The Urysohn space

We cannot apply Fraïssé directly: there are too many finite metric spaces (uncountably many two-point spaces).

Instead, we define a metric space to be **rational** if all distances are rational numbers. The class of rational metric spaces satisfies Fraïssé's theorem; its Fraïssé limit is the **rational**

Urysohn space. Taking its completion in the usual sense (add limits for Cauchy sequences) gives the real Urysohn space.

One can replace the rational numbers by other sets such as the non-negative integers (giving the **integral Urysohn space**, a distance-transitive graph) or the set $\{0, 1, 2\}$. Metric spaces with distances $\{0, 1, 2\}$ can be regarded as graphs (distance 1 is adjacency) and conversely. Now the $\{0, 1, 2\}$ Urysohn space is the random graph R !

The Urysohn space

We cannot apply Fraïssé directly: there are too many finite metric spaces (uncountably many two-point spaces).

Instead, we define a metric space to be **rational** if all distances are rational numbers. The class of rational metric spaces satisfies Fraïssé's theorem; its Fraïssé limit is the **rational**

Urysohn space. Taking its completion in the usual sense (add limits for Cauchy sequences) gives the real Urysohn space.

One can replace the rational numbers by other sets such as the non-negative integers (giving the **integral Urysohn space**, a distance-transitive graph) or the set $\{0, 1, 2\}$. Metric spaces with distances $\{0, 1, 2\}$ can be regarded as graphs (distance 1 is adjacency) and conversely. Now the $\{0, 1, 2\}$ Urysohn space is the random graph R !

Vershik and I studied this and found various analogies between the random graph and the Urysohn space.

25 April bridge, Lisbon



3. Graphs on groups

In 1955, Brauer and Fowler published a paper which, with hindsight, was the first step towards the classification of the finite simple groups. They proved that there are only finitely many simple groups of even order containing an involution (an element of order 2) with a prescribed centralizer. The “even order” provision was needed because this predates the famous Feit–Thompson theorem that a non-abelian finite simple group must necessarily have even order.

3. Graphs on groups

In 1955, Brauer and Fowler published a paper which, with hindsight, was the first step towards the classification of the finite simple groups. They proved that there are only finitely many simple groups of even order containing an involution (an element of order 2) with a prescribed centralizer. The “even order” provision was needed because this predates the famous Feit–Thompson theorem that a non-abelian finite simple group must necessarily have even order.

Though the word “graph” does not occur in the paper, the main tool they used was the **commuting graph** of the group, the graph whose vertices are the non-identity elements, two vertices x and y joined if they commute (that is, $xy = yx$).

3. Graphs on groups

In 1955, Brauer and Fowler published a paper which, with hindsight, was the first step towards the classification of the finite simple groups. They proved that there are only finitely many simple groups of even order containing an involution (an element of order 2) with a prescribed centralizer. The “even order” provision was needed because this predates the famous Feit–Thompson theorem that a non-abelian finite simple group must necessarily have even order.

Though the word “graph” does not occur in the paper, the main tool they used was the **commuting graph** of the group, the graph whose vertices are the non-identity elements, two vertices x and y joined if they commute (that is, $xy = yx$).

Since then, several more graphs on groups have been defined and studied, and many interactions found. (I must thank my friend Ambat Vijayakumar for organising a study group in 2021 which led to much of this research, as well as many colleagues, in India and elsewhere, who have collaborated with me on this.)

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

Given a finite group G , the following graphs have vertex set G .

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

Given a finite group G , the following graphs have vertex set G .

- ▶ In the **power graph**, x is joined to y if one of x and y is a power of the other.

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

Given a finite group G , the following graphs have vertex set G .

- ▶ In the **power graph**, x is joined to y if one of x and y is a power of the other.
- ▶ In the **enhanced power graph**, x is joined to y if both x and y are powers of an element z .

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

Given a finite group G , the following graphs have vertex set G .

- ▶ In the **power graph**, x is joined to y if one of x and y is a power of the other.
- ▶ In the **enhanced power graph**, x is joined to y if both x and y are powers of an element z .
- ▶ The power graph is a spanning subgraph of the enhanced power graph; in the **difference graph**, x and y are joined if they are joined in the enhanced power graph but not in the power graph. I denote it $D(G)$.

The difference graph

I will not attempt a survey, but turn immediately to one type of graph, studied by Sucharita Biswas, Angsuman Das, Hiranya Kishore Dey and me.

Given a finite group G , the following graphs have vertex set G .

- ▶ In the **power graph**, x is joined to y if one of x and y is a power of the other.
- ▶ In the **enhanced power graph**, x is joined to y if both x and y are powers of an element z .
- ▶ The power graph is a spanning subgraph of the enhanced power graph; in the **difference graph**, x and y are joined if they are joined in the enhanced power graph but not in the power graph. I denote it $D(G)$.

It may happen that the power graph and enhanced power graph coincide, so that the difference graph is null. Groups with these properties are called **EPPO groups**, those in which every element has prime power order. They were determined by the efforts of Higman, Suzuki and Brandl.

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

A graph Γ is a **cograph** if it contains no induced 4-vertex path. For any graph Γ , the following are equivalent:

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

A graph Γ is a **cograph** if it contains no induced 4-vertex path. For any graph Γ , the following are equivalent:

- ▶ Γ is a cograph;

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

A graph Γ is a **cograph** if it contains no induced 4-vertex path. For any graph Γ , the following are equivalent:

- ▶ Γ is a cograph;
- ▶ Γ can be built from 1-vertex graphs by the operations of disjoint union and complementation;

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

A graph Γ is a **cograph** if it contains no induced 4-vertex path. For any graph Γ , the following are equivalent:

- ▶ Γ is a cograph;
- ▶ Γ can be built from 1-vertex graphs by the operations of disjoint union and complementation;
- ▶ twin reduction converts Γ to a 1-vertex graph.

Twin reduction

Two vertices of a graph are **twins** if they have the same neighbours (possibly excepting each other). The process of **twin reduction** consists of repeatedly choosing a pair of twins and identifying them until no twins remain. The result of twin reduction is unique up to isomorphism, independent of the order of the process.

A graph Γ is a **cograph** if it contains no induced 4-vertex path. For any graph Γ , the following are equivalent:

- ▶ Γ is a cograph;
- ▶ Γ can be built from 1-vertex graphs by the operations of disjoint union and complementation;
- ▶ twin reduction converts Γ to a 1-vertex graph.

It is an open problem which groups have the property that their difference graph is a cograph.

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals.
Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals. Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

One such case is the Mathieu group M_{11} , a simple group of order 7920. Twin reduction yields a graph Γ on 385 vertices which has some remarkable properties:

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals. Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

One such case is the Matheu group M_{11} , a simple group of order 7920. Twin reduction yields a graph Γ on 385 vertices which has some remarkable properties:

- ▶ Γ is bipartite, with bipartite blocks of sizes 165 and 220.

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals. Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

One such case is the Matheu group M_{11} , a simple group of order 7920. Twin reduction yields a graph Γ on 385 vertices which has some remarkable properties:

- ▶ Γ is bipartite, with bipartite blocks of sizes 165 and 220.
- ▶ Γ is semiregular; the valencies of vertices in the two bipartite blocks are 4 and 3 respectively.

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals. Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

One such case is the Matheu group M_{11} , a simple group of order 7920. Twin reduction yields a graph Γ on 385 vertices which has some remarkable properties:

- ▶ Γ is bipartite, with bipartite blocks of sizes 165 and 220.
- ▶ Γ is semiregular; the valencies of vertices in the two bipartite blocks are 4 and 3 respectively.
- ▶ Γ has diameter 10 and girth 10.

The jewel in the lotus

Twin reduction is like the stripping away of lotus petals. Maybe nothing remains; but in interesting cases we may find a beautiful jewel.

One such case is the Matheu group M_{11} , a simple group of order 7920. Twin reduction yields a graph Γ on 385 vertices which has some remarkable properties:

- ▶ Γ is bipartite, with bipartite blocks of sizes 165 and 220.
- ▶ Γ is semiregular; the valencies of vertices in the two bipartite blocks are 4 and 3 respectively.
- ▶ Γ has diameter 10 and girth 10.
- ▶ The automorphism group of Γ is the Mathieu group M_{11} .

What next?

There are many groups, several graphs defined on groups, and a few different ways of “reducing” a graph in addition to twin reduction. So there is plenty more to explore. I invite you all to help explore thus and find more beautiful graphs.

What next?

There are many groups, several graphs defined on groups, and a few different ways of “reducing” a graph in addition to twin reduction. So there is plenty more to explore. I invite you all to help explore thus and find more beautiful graphs.

One curiosity. I have tried three different graphs and various reduction methods on the group M_{11} . All methods seem to give the same example. I do not know why.

What next?

There are many groups, several graphs defined on groups, and a few different ways of “reducing” a graph in addition to twin reduction. So there is plenty more to explore. I invite you all to help explore this and find more beautiful graphs.

One curiosity. I have tried three different graphs and various reduction methods on the group M_{11} . All methods seem to give the same example. I do not know why.

Also, it would be good to explore this beautiful graph further.

What next?

There are many groups, several graphs defined on groups, and a few different ways of “reducing” a graph in addition to twin reduction. So there is plenty more to explore. I invite you all to help explore thus and find more beautiful graphs.

One curiosity. I have tried three different graphs and various reduction methods on the group M_{11} . All methods seem to give the same example. I do not know why.

Also, it would be good to explore this beautiful graph further. But that is the end of my story.



... for your attention.