

# Machine Learning

A. G. Schwing & M. Telgarsky

University of Illinois at Urbana-Champaign, 2018

## L2: Linear Regression

**Last time:**  $k$ -NN.

- Pros: simple (easy to implement and reason about).
- Cons: stores all data; curse of dimension.

**This time:** Linear regression (“ordinary least squares”).

- Also simple!
- Reading: K. Murphy; Machine Learning: A Probabilistic Perspective; Chapter 7.


## Least squares model.

Predict  $y \in \mathbb{R}$  (“label”, “response”)  
from  $\mathbf{x} \in \mathbb{R}^d$  (“features”, “covariate”)  
via  $\mathbf{w}_1^\top \mathbf{x} + w_2$  (where  $\mathbf{w}_1 \in \mathbb{R}^d$  and  $w_2 \in \mathbb{R}$ )


- **Learning:** choose  $(\mathbf{w}_1, w_2)$  from data  $((\mathbf{x}^{(i)}, y^{(i)}))_{i=1}^N$ .
- **Prediction/inference:** obtain  $\mathbf{x}$ , output  $\mathbf{w}_1^\top \mathbf{x} + w_2$ .

**Note.**  $y \in \mathbb{R}$  (“*regression*”) rather than  $\{-1, +1\}$  (“*classification*”).

**Example:** Old faithful eruptions: duration ( $x$ ) vs delay ( $y$ ).




**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).




Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ?

**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ?


**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ?




**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ? If all  $\mathbf{x}^{(i)}$  coincide:


**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ? If all  $\mathbf{x}^{(i)}$  coincide:

$$\hat{y} = \frac{1}{N} \sum_{i=1}^N y^{(i)}$$


**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ? If all  $\mathbf{x}^{(i)}$  coincide:

$$\hat{y} = \frac{1}{N} \sum_{i=1}^N y^{(i)}$$


**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ? If all  $\mathbf{x}^{(i)}$  coincide:

$$\hat{y} = \frac{1}{N} \sum_{i=1}^N y^{(i)} = \arg \min_{y \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} (y - y^{(i)})^2.$$

**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).




Which line  $\mathbf{x} \mapsto \mathbf{w}_1^\top \mathbf{x} + w_2$ ? If all  $\mathbf{x}^{(i)}$  coincide:

$$\hat{y} = \frac{1}{N} \sum_{i=1}^N y^{(i)} = \arg \min_{y \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} (y - y^{(i)})^2.$$

**Remark.** Mean has issues... we'll revisit this...


**Example:** Old faithful eruptions: duration ( $x$ ) vs delay ( $y$ ).



**Ignoring  $x$ :**

$$\arg \min_{w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( w_2 - y^{(i)} \right)^2.$$

**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).



**Ignoring  $\mathbf{x}$ :**

$$\arg \min_{w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( w_2 - y^{(i)} \right)^2.$$

**General form:**

$$\arg \min_{\mathbf{w}_1 \in \mathbb{R}^d, w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( \mathbf{w}_1^\top \mathbf{x}^{(i)} + w_2 - y^{(i)} \right)^2.$$

**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).

**Ignoring  $\mathbf{x}$ :** 
$$\arg \min_{w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( w_2 - y^{(i)} \right)^2.$$

**General form:** 
$$\arg \min_{\mathbf{w}_1 \in \mathbb{R}^d, w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( \mathbf{w}_1^\top \mathbf{x}^{(i)} + w_2 - y^{(i)} \right)^2.$$



**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).

**Ignoring  $\mathbf{x}$ :** 
$$\arg \min_{w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( w_2 - y^{(i)} \right)^2.$$

**General form:** 
$$\arg \min_{\mathbf{w}_1 \in \mathbb{R}^d, w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( \mathbf{w}_1^\top \mathbf{x}^{(i)} + w_2 - y^{(i)} \right)^2.$$

**Simplification:**

$$\mathbf{y} := \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(N)} \end{bmatrix}, \quad \mathbf{X} := \begin{bmatrix} \leftarrow \mathbf{x}^{(1)} \rightarrow & 1 \\ & \vdots \\ \leftarrow \mathbf{x}^{(N)} \rightarrow & 1 \end{bmatrix}, \quad \mathbf{w} := \begin{bmatrix} \uparrow \\ \mathbf{w}_1 \\ \downarrow \\ w_2 \end{bmatrix}.$$

**Example:** Old faithful eruptions: duration ( $\mathbf{x}$ ) vs delay ( $y$ ).

**Ignoring  $\mathbf{x}$ :** 
$$\arg \min_{w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( w_2 - y^{(i)} \right)^2.$$

**General form:** 
$$\arg \min_{\mathbf{w}_1 \in \mathbb{R}^d, w_2 \in \mathbb{R}} \frac{1}{N} \sum_{i=1}^N \frac{1}{2} \left( \mathbf{w}_1^\top \mathbf{x}^{(i)} + w_2 - y^{(i)} \right)^2.$$

**Simplification:** 
$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 \quad \text{where}$$

$$\mathbf{y} := \begin{bmatrix} y^{(1)} \\ \vdots \\ y^{(N)} \end{bmatrix}, \quad \mathbf{X} := \begin{bmatrix} \leftarrow \mathbf{x}^{(1)} \rightarrow & 1 \\ & \vdots \\ \leftarrow \mathbf{x}^{(N)} \rightarrow & 1 \end{bmatrix}, \quad \mathbf{w} := \begin{bmatrix} \uparrow \\ \mathbf{w}_1 \\ \downarrow \\ w_2 \end{bmatrix}.$$

**Goal:** Solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 \quad \text{where } \mathbf{w} \in \mathbb{R}^{d+1}, \mathbf{X} \in \mathbb{R}^{N \times (d+1)}, \mathbf{y} \in \mathbb{R}^N.$$

**Goal:** Solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 \quad \text{where } \mathbf{w} \in \mathbb{R}^{d+1}, \mathbf{X} \in \mathbb{R}^{N \times (d+1)}, \mathbf{y} \in \mathbb{R}^N.$$

**Ordinary least squares (OLS) estimator:**

choose  $\hat{\mathbf{w}} := (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Derivation:** setting derivative to 0, optimal  $\hat{\mathbf{w}}$  satisfies

$$\mathbf{X}^\top (\mathbf{X}\hat{\mathbf{w}} - \mathbf{y}) = 0 \quad \text{and thus} \quad \mathbf{X}^\top \mathbf{X}\hat{\mathbf{w}} = \mathbf{X}^\top \mathbf{y}.$$

*When it exists*, we can write  $\hat{\mathbf{w}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Non-existence** of OLS solution  $\hat{\mathbf{w}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Solution #1:** “Ridge regression”: solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2,$$

giving  $\tilde{\mathbf{w}} := (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Note.** In this course’s homeworks and tests, you may assume  $(\mathbf{X}^\top \mathbf{X})^{-1}$  exists unless otherwise specified.

**Non-existence** of OLS solution  $\hat{\mathbf{w}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Solution #1:** “Ridge regression”: solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2,$$

giving  $\tilde{\mathbf{w}} := (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Remark.** “ $+\frac{\lambda}{2} \|\mathbf{w}\|^2$ ” is *regularization*;  
it affects computation and statistics.

**Note.** In this course’s homeworks and tests, you may assume  $(\mathbf{X}^\top \mathbf{X})^{-1}$  exists unless otherwise specified.

**Non-existence** of OLS solution  $\hat{\mathbf{w}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Solution #1:** “Ridge regression”: solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2,$$

giving  $\tilde{\mathbf{w}} := (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Note.** In this course’s homeworks and tests, you may assume  $(\mathbf{X}^\top \mathbf{X})^{-1}$  exists unless otherwise specified.

**Non-existence** of OLS solution  $\hat{\mathbf{w}} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Solution #1:** “Ridge regression”: solve

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2,$$

giving  $\tilde{\mathbf{w}} := (\mathbf{X}^\top \mathbf{X} + \lambda \mathbf{I})^{-1} \mathbf{X}^\top \mathbf{y}$ .

**Solution #2:** use the *pseudoinverse*:

replace  $(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$  with  $(\mathbf{X}^\top \mathbf{X})^\dagger \mathbf{X}^\top \mathbf{y} = \mathbf{X}^\dagger \mathbf{y}$ .

**Remark.** This still satisfies the “derivative condition”


$$(\mathbf{X}^\top \mathbf{X}) \hat{\mathbf{w}} = \mathbf{X}^\top \mathbf{y}$$

and therefore is optimal!

**Note.** In this course’s homeworks and tests, you may assume  $(\mathbf{X}^\top \mathbf{X})^{-1}$  exists unless otherwise specified.



## Summary so far



### Least squares problem

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2.$$

### OLS solution

$$(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}.$$

### Question:

still “why this line” ?

## Three justifications/interpretations.

- Geometric interpretation.
- Probabilistic model.
- Loss minimization.

## Geometric interpretation.

Focus on **columns** of  $\mathbf{X}$ :

$$\mathbf{X} = \begin{bmatrix} \leftarrow \mathbf{x}^{(1)} \rightarrow & 1 \\ \vdots & \vdots \\ \leftarrow \mathbf{x}^{(N)} \rightarrow & 1 \end{bmatrix} = \begin{bmatrix} \uparrow & & \uparrow \\ \mathbf{z}_1 & \cdots & \mathbf{z}_{d+1} \\ \downarrow & & \downarrow \end{bmatrix}.$$

Then *residual*  $\mathbf{X}\hat{\mathbf{w}} - \mathbf{y}$  is **orthogonal** to  $\text{span}(\{\mathbf{z}_1, \dots, \mathbf{z}_{d+1}\})$ .  
(... since  $\mathbf{X}^\top(\mathbf{X}\hat{\mathbf{w}} - \mathbf{y}) = 0$ .)





figure credit: daniel hsu

## Probabilistic model.

**Suppose** “linear model with Gaussian errors”:  
label  $y$  at point  $\mathbf{x}$  has distribution  $\text{Gaussian}(\bar{\mathbf{w}}^\top \mathbf{x}, \sigma^2)$ :

$$p(y^{(i)}|\mathbf{x}^{(i)}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y^{(i)} - \bar{\mathbf{w}}^\top \mathbf{x}^{(i)})^2\right)$$



## Probabilistic model.

**Suppose** “linear model with Gaussian errors”:

label  $y$  at point  $\mathbf{x}$  has distribution  $\text{Gaussian}(\bar{\mathbf{w}}^\top \mathbf{x}, \sigma^2)$ :

$$p(y^{(i)}|\mathbf{x}^{(i)}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y^{(i)} - \bar{\mathbf{w}}^\top \mathbf{x}^{(i)})^2\right)$$

## Probabilistic model.

**Suppose** “linear model with Gaussian errors”:

label  $y$  at point  $\mathbf{x}$  has distribution  $\text{Gaussian}(\bar{\mathbf{w}}^\top \mathbf{x}, \sigma^2)$ :

$$p(y^{(i)}|\mathbf{x}^{(i)}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y^{(i)} - \bar{\mathbf{w}}^\top \mathbf{x}^{(i)})^2\right)$$

To solve, *maximize likelihood*:

$$\begin{aligned} \arg \max_{\mathbf{w} \in \mathbb{R}^{d+1}} \prod_{i=1}^N p(y^{(i)}|\mathbf{x}^{(i)}) &= (\dots \text{hwk1} \dots) \\ &= \arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2\sigma^2} \sum_{i=1}^N \frac{1}{2} \left( \mathbf{w}^\top \mathbf{x}^{(i)} - y^{(i)} \right)^2. \end{aligned}$$

## Loss minimization

The form  $\|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$  was convenient (OLS solution  $\hat{\mathbf{w}} = \mathbf{X}^\dagger \mathbf{y}$ ).

## Loss minimization

The form  $\|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$  was convenient (OLS solution  $\hat{\mathbf{w}} = \mathbf{X}^\dagger \mathbf{y}$ ).

Again write

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{2} (\mathbf{w}^\top \mathbf{x}^{(i)} - y^{(i)})^2 = \frac{1}{N} \sum_{i=1}^N \ell_{\text{ls}}(y^{(i)}, \mathbf{w}^\top \mathbf{x}^{(i)})$$

where now  $\ell_{\text{ls}}(y, \hat{y}) = \frac{1}{2}(y - \hat{y})^2$  is the *least squares loss*.



## Loss minimization

The form  $\|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$  was convenient (OLS solution  $\hat{\mathbf{w}} = \mathbf{X}^\dagger \mathbf{y}$ ).

Again write

$$\frac{1}{N} \sum_{i=1}^N \frac{1}{2} (\mathbf{w}^\top \mathbf{x}^{(i)} - y^{(i)})^2 = \frac{1}{N} \sum_{i=1}^N \ell_{\text{ls}}(y^{(i)}, \mathbf{w}^\top \mathbf{x}^{(i)})$$

where now  $\ell_{\text{ls}}(y, \hat{y}) = \frac{1}{2}(y - \hat{y})^2$  is the *least squares loss*.

The general form

$$\arg \min_f \frac{1}{N} \sum_{i=1}^N \ell(y^{(i)}, f(\mathbf{x}^{(i)}))$$

is the standard ML idea **Empirical Risk Minimization (ERM)**.

## Three justifications/interpretations.

- Geometric interpretation.
- Probabilistic model.
- Loss minimization.

## Three other questions.

- Classification vs regression.
- How to implement  $\mathbf{X}^\dagger \mathbf{y}$ ?
- Nonlinear least squares.

## Classification vs Regression.

Given  $\mathbf{x}$ , then  $\hat{\mathbf{w}}^\top \mathbf{x} \in \mathbb{R}$  (“regression”)

Alternatively,  $\text{sgn}(\hat{\mathbf{w}}^\top \mathbf{x}) \in \{-1, +1\}$  (“classification”).

## Classification vs Regression.

Given  $\mathbf{x}$ , then  $\hat{\mathbf{w}}^\top \mathbf{x} \in \mathbb{R}$  (“regression”)

Alternatively,  $\text{sgn}(\hat{\mathbf{w}}^\top \mathbf{x}) \in \{-1, +1\}$  (“classification”).

Recall least squares loss  $\ell_{\text{ls}}(y, \hat{y}) = (y - \hat{y})^2/2$ ; if  $y \in \{-1, +1\}$ ,

$$(y - \hat{y})^2/2 = (y^2)(1 - y\hat{y})^2/2 = (1 - y\hat{y})^2/2.$$

Seems weird if our goal is to minimize  $\mathbf{1}[y \neq \hat{y}] \approx \mathbf{1}[y\hat{y} \geq 0]$ ?

## Classification vs Regression.


Given  $\mathbf{x}$ , then  $\hat{\mathbf{w}}^\top \mathbf{x} \in \mathbb{R}$  (“regression”)

Alternatively,  $\text{sgn}(\hat{\mathbf{w}}^\top \mathbf{x}) \in \{-1, +1\}$  (“classification”).

Recall least squares loss  $\ell_{\text{ls}}(y, \hat{y}) = (y - \hat{y})^2/2$ ; if  $y \in \{-1, +1\}$ ,

$$(y - \hat{y})^2/2 = (y^2)(1 - y\hat{y})^2/2 = (1 - y\hat{y})^2/2.$$

Seems weird if our goal is to minimize  $\mathbf{1}[y \neq \hat{y}] \approx \mathbf{1}[y\hat{y} \geq 0]$ ?



## Classification vs Regression.


Given  $\mathbf{x}$ , then  $\hat{\mathbf{w}}^\top \mathbf{x} \in \mathbb{R}$  (“regression”)

Alternatively,  $\text{sgn}(\hat{\mathbf{w}}^\top \mathbf{x}) \in \{-1, +1\}$  (“classification”).

Recall least squares loss  $\ell_{\text{ls}}(y, \hat{y}) = (y - \hat{y})^2/2$ ; if  $y \in \{-1, +1\}$ ,

$$(y - \hat{y})^2/2 = (y^2)(1 - y\hat{y})^2/2 = (1 - y\hat{y})^2/2.$$

Seems weird if our goal is to minimize  $\mathbf{1}[y \neq \hat{y}] \approx \mathbf{1}[y\hat{y} \geq 0]$ ?



**Note.** Even in easy cases, linear classification is **NP**-hard!

## How to solve.

**Question:** are  $(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$  and  $\mathbf{X}^\dagger \mathbf{y}$  in “closed form”?

## How to solve.


**Question:** are  $(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$  and  $\mathbf{X}^\dagger \mathbf{y}$  in “closed form”?

(Libraries will use iterative solvers!)


Since  $\mathbf{w} \mapsto \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|^2$  is convex,  
there are many “efficient” *iterative descent methods*.




## **Nonlinear regression.** Sometimes linear is not enough...



## **Nonlinear regression.** Sometimes linear is not enough...




## **Nonlinear regression.** Sometimes linear is not enough...




Polynomial fit?

## **Nonlinear regression.** Sometimes linear is not enough...



Polynomial fit? (...no thanks...)


**Nonlinear regression.** Sometimes linear is not enough...



Polynomial fit? (...no thanks...)

**How to solve:** replace  $\mathbf{x}^{(i)}$  with *features*  $\tilde{\mathbf{x}}^{(i)} = \phi(\mathbf{x}^{(i)})$ .

## Summary.



### Least squares problem

$$\arg \min_{\mathbf{w} \in \mathbb{R}^{d+1}} \frac{1}{2} \|\mathbf{X}\mathbf{w} - \mathbf{y}\|_2^2.$$

### OLS solution

$$(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}.$$

### Justification.

- Geometric.
- Probabilistic model.
- ERM.

### Concepts.

- Regularization.
- ERM and loss functions.
- Maximum likelihood.