

Proof of the associative law for matrices

Kamila Zdybał

kamilazdybal.github.io/science-docs, kamila.zdybal@gmail.com

Motivation

I was reviewing the MIT course 18.06 on Linear Algebra by Professor Gilbert Strang and the associative law for matrices first appeared on the blackboard in more or less this way:

$$(AB)C = A(BC) \quad (1)$$

Professor Strang said then: "It's not that easy to prove that this is correct. You have to go into the gory details of matrix multiplication, do it both ways and see that you come out the same." So I said: let's do it!

This short note contains the proof of eq.(1).

Proof

Suppose we have three matrices: A of size $(m \times n)$, B of size $(n \times p)$ and C of size $(p \times r)$. We are going to prove that the relation from the eq.(1) holds for these matrices.

We start off with the general equation for the i th- j th element of a matrix that is a result of multiplication of two matrices A and B . This can be found in a linear algebra textbook to be:

$$\gamma_{ij}^{AB} = \sum_{k=1}^n (a_{ik} b_{kj}) \quad (2)$$

Similarly, the general equation for the i th- j th element of a matrix that is a result of multiplication of matrices B and C is:

$$\gamma_{ij}^{BC} = \sum_{k=1}^p (b_{ik} c_{kj}) \quad (3)$$

We then write the general equation for the i th- j th element of a matrix that is a result of the complete multiplication $(AB)C$:

$$\gamma_{ij}^{(AB)C} = \sum_{l=1}^p \left(\left(\sum_{k=1}^n (a_{ik} b_{kl}) \right) c_{lj} \right) \quad (4)$$

and the general equation for the i th- j th element of a matrix that is a result of the complete multiplication $A(BC)$:

$$\gamma_{ij}^{A(BC)} = \sum_{h=1}^n \left(a_{ih} \left(\sum_{k=1}^p (b_{hk} c_{kj}) \right) \right) \quad (5)$$

Now we are going to take eq.(4) and eq.(5) and write the sums element by element.

We start with eq.(4):

$$\begin{aligned} \gamma_{ij}^{(AB)C} &= \sum_{l=1}^p \left(\left(\sum_{k=1}^n (a_{ik} b_{kl}) \right) c_{lj} \right) = \\ &= \sum_{l=1}^p (a_{i1} b_{1l} c_{lj} + \cdots + a_{in} b_{nl} c_{lj}) = \\ &= a_{i1} b_{11} c_{1j} + \cdots + a_{in} b_{n1} c_{1j} + \\ &\quad + a_{i1} b_{12} c_{2j} + \cdots + a_{in} b_{n2} c_{2j} + \\ &\quad \vdots \\ &\quad + a_{i1} b_{1p} c_{pj} + \cdots + a_{in} b_{np} c_{pj} \end{aligned} \quad (6)$$

For eq.(5) we have:

$$\begin{aligned} \gamma_{ij}^{A(BC)} &= \sum_{h=1}^n \left(a_{ih} \left(\sum_{k=1}^p (b_{hk} c_{kj}) \right) \right) = \\ &= \sum_{h=1}^n (a_{ih} b_{h1} c_{1j} + \cdots + a_{ih} b_{hp} c_{pj}) = \\ &= a_{i1} b_{11} c_{1j} + \cdots + a_{i1} b_{1p} c_{pj} + \\ &\quad + a_{i2} b_{21} c_{1j} + \cdots + a_{i2} b_{2p} c_{pj} + \\ &\quad \vdots \\ &\quad + a_{in} b_{n1} c_{1j} + \cdots + a_{in} b_{np} c_{pj} \end{aligned} \quad (7)$$

In eq.(6), we have the final sum written in p "rows" and in each row there is n elements of the sum.

In eq.(7), we have the final sum written in n "rows" and in each row there is p elements of the sum.

Notice now, that the sum of all the first elements from each row of eq.(6) is the same as the sum of all the first's row elements in the eq.(7) (marked in orange in the equations).

The same holds for the second elements from each row of eq.(6) with the second's row elements in eq.(7), and so on.

If we were to write all the elements in an array (that would be size $(p \times n)$ for eq.(6) and $(n \times p)$ for eq.(7)), it would be equivalent to saying that taking the sum of elements from any column q of the case $(AB)C$ is the same as the sum of elements from row q of the case $A(BC)$.

This concludes the proof.¹

¹Especially that I'm running out of letters to index things.