

Least squares regression

a short story on overdetermined linear systems and Moore-Penrose pseudoinverse

This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) license.

Kamila Zdybał

Université libre de Bruxelles, kamila.zdybal@ulb.ac.be
camillejr.github.io/science-docs, kamila.zdybal@gmail.com

Preface

Imagine a linear system of equations with more number equations than the number of unknowns.

Goal of this paper: explain why Moore-Penrose inverse give a least squares regression.

This document is still in preparation. Please feel free to contact me with any suggestions, corrections or comments.

Keywords

overdetermined linear systems, partial least squares regression, Moore-Penrose inverse

Contents

1 Partial Least Squares (PLS) regression

1

1 Partial Least Squares (PLS) regression

2D data example

Have a data set in pairs: (X, Y) . X is a vector of N points and Y is a vector of corresponding N points. Together they make a cloud of points on a 2D-plane.

We now say that: $XA = Y$.

We are interested in finding the coefficients: $y = Cx + D$.




Figure 1: Linear basis function LS regression.




Figure 2: Non-linear (quadratic) basis function LS regression.

References

- [1] Nathan Kutz, *Data Driven Discovery of Dynamical Systems and PDEs*, an online lecture
- [2] Gilbert Strang, *Introduction to Linear Algebra*, Fifth Edition, 2016