

Record 8 of 10



The U.S. government does not review or approve the safety and science of all studies listed on this website.

Read our full [disclaimer](https://clinicaltrials.gov/about-site/disclaimer) (<https://clinicaltrials.gov/about-site/disclaimer>) for details.

Evaluation Of The Efficacy And Safety Of Single Doses Of PF-05089771 In Patients With Primary (Inherited) Erythromelalgia (IEM)

Completed ⓘ

No longer looking for participants

ClinicalTrials.gov ID ⓘ NCT01769274

Sponsor ⓘ Pfizer

Information provided by ⓘ Pfizer (Responsible Party)

Last Update Posted ⓘ 2019-11-19

Results Posted Tab

Results Overview

Conditions ⓘ Inherited Erythromelalgia Intervention/Treatment ⓘ - Drug: PF-05089771 - Drug: Placebo Other Study ID Numbers ⓘ - B3291006 Study Design Allocation ⓘ: Randomized Interventional Model ⓘ: Crossover Assignment Masking ⓘ: Double (Participant, Investigator) Primary Purpose ⓘ: Treatment Results Point of Contact Name/Title: Pfizer ClinicalTrials.gov Call Center Organization: Pfizer Inc. Phone: 1-800-718-1021 Email: ClinicalTrials.gov_Inquiries@pfizer.com	Enrollment (Actual) ⓘ 5 Study Type ⓘ Interventional
--	---



Study Record Dates

These dates track the progress of study record and summary results submissions to ClinicalTrials.gov. Study records and reported results are reviewed by the National Library of Medicine (NLM) to make sure they meet specific quality control standards before being posted on the public website.

Study Registration Dates	Results Reporting Dates
First Submitted ⓘ 2012-10-18 First Submitted that Met QC Criteria ⓘ 2013-01-15 First Posted (Estimated) ⓘ 2013-01-16	Results First Submitted ⓘ 2019-10-25 Results First Submitted that Met QC Criteria ⓘ 2019-11-18 Results First Posted ⓘ 2019-11-19
Certification/Extension Dates	Study Record Updates
Certification/Extension First Submitted ⓘ 2013-07-30 Certification/Extension First Submitted that Met QC Criteria ⓘ 2013-07-30 Certification/Extension First Posted (Estimated) ⓘ 2013-08-06	Last Update Submitted that Met QC Criteria ⓘ 2019-11-18 Last Update Posted ⓘ 2019-11-19 Last Verified ⓘ 2019-11

Participant Flow ⓘ

Recruitment Details	[Not Specified]
Pre-assignment Details	[Not Specified]

Arm/Group Title	Part A	Part B: PF-05089771 Then Placebo Then Placebo Then PF-	Part B: Placebo Then PF-05089771 Then PF-05089771
-----------------	--------	--	---

Baseline Characteristics ⓘ

Arm/Group Title	All Participants
Arm/Group Description	All participants who were enrolled in the study.
Overall Number of Baseline Participants	5
Baseline Analysis Population Description	Safety set included all participants who received at least 1 dose of study medication.

[Expand all](#) / [Collapse all](#)

Age, Continuous

Mean (Standard Deviation) | Unit of measure: years

Number Analyzed	5 participants
	40.2 (18.7)

Sex: Female, Male

Measure Type: Count of Participants | Unit of measure: Participants

Number Analyzed	5 participants
Female	2 40.0%

Outcome Measures

[Expand all](#) / [Collapse all](#)

1. Average Pain Intensity Numerical Rating Scale (PI-NRS) Score - From 0 to 4 Hours Post-dose

Type: Primary | Time Frame: Every 15 minutes from 0 to 4 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10, where 0= no pain and 10= worst possible pain; higher scores signify more pain. The average pain score was calculated as the mean of the pain scores recorded every 15 minutes from 0 to 4 hours post-dose.		
Time Frame	Every 15 minutes from 0 to 4 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Mean (Standard Deviation) Unit of Measure: Units on a scale	2.08 (2.20)	2.73 (2.68)	1.95 (2.29)

2. Average PI-NRS Score - From Post Evoked Pain Time Point 2 (EP2) to 8 Hours Post-dose

Type: Secondary | Time Frame: Post EP2, every 5 minutes for the first hour, then every 15 minutes up to 8 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 2= first time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP2 is approximately a time point 4-5 hours post-dose. The average pain score was calculated as the mean of the pain scores recorded from EP2 to 8 hours post-dose.		
Time Frame	Post EP2, every 5 minutes for the first hour, then every 15 minutes up to 8 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Units on a scale	0.70 (0.64)	0.61 (0.46)	0.55 (0.51)

3. Average PI-NRS Scores - From Post Evoked Pain Time Point 3 (EP3) to 10 Hours Post-dose

Type: Secondary | Time Frame: Post EP3, every 5 minutes for the first hour, then every 15 minutes up to 10 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 3= second time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP3 is approximately a time point 8-9 hours post-dose. The average pain score was calculated as the mean of the pain scores recorded from EP3 to 10 hours post-dose.		
Time Frame	Post EP3, every 5 minutes for the first hour, then every 15 minutes up to 10 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Units on a scale	1.19 (1.22)	1.55 (1.15)	0.82 (0.52)

4. Average PI-NRS Scores - From Post Evoked Pain Time Point 4 (EP4) to 28 Hours Post-dose

Type: Secondary | Time Frame: Post EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 4= third time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP4 is approximately a time point 24-25 hours post-dose. The average pain score was calculated as the mean of the pain scores recorded from EP4 to 28 hours post-dose.		
Time Frame	Post EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	3	3	3
Mean (Standard Deviation) Unit of Measure: Units on a scale	0.99 (0.52)	0.91 (0.53)	0.36 (0.08)

5. Maximum PI-NRS Scores - From 0 Hour to 4 Hours Post-dose

Type: Secondary | Time Frame: From 0 hour to 4 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Maximum pain score in 4 hours period post-dosing is reported.		
Time Frame	From 0 hour to 4 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Mean (Standard Deviation) Unit of Measure: Units on a scale	7.6 (1.7)	8.0 (1.4)	8.2 (2.1)

6. Maximum PI-NRS Scores - From Post EP2 to 8 Hours Post-dose

Type: Secondary | Time Frame: From post EP2 to 8 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 2= first time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP2 is approximately a time point 4-5 hours post-dose. Maximum pain score in period from post EP2 to 8 hours post-dose is reported.		
Time Frame	From post EP2 to 8 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Units on a scale	3.5 (1.7)	5.7 (3.1)	5.5 (3.3)

7. Maximum PI-NRS Scores - From Post EP3 to 10 Hours Post-dose

Type: Secondary | Time Frame: From the end of EP3 to 10 hours post dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 3= second time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP3 is approximately a time point 8-9 hours post-dose. Maximum pain score in period from post EP3 to 10 hours post-dose is reported.		
Time Frame	From the end of EP3 to 10 hours post dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Units on a scale	2.3 (2.6)	6.0 (3.6)	5.3 (2.6)

8. Maximum PI-NRS Scores - From Post EP4 to 28 Hours Post-dose

Type: Secondary | Time Frame: Post EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 4= third time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP4 is approximately a time point 24-25 hours post-dose. Maximum pain score in period from post EP4 to 28 hours post-dose is reported.		
Time Frame	Post EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	3	3	3
Mean (Standard Deviation) Unit of Measure: Units on a scale	7.3 (1.2)	7.3 (2.5)	5.7 (2.1)

9. Duration When PI-NRS Scores Were Greater Than (>) 5 - From 0 Hour to 4 Hours Post-dose

Type: Secondary | Time Frame: From 0 hour to 4 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. The duration of time that participants experienced PI-NRS score greater than 5 from the 0 hours to 4 hours post-dose.		
Time Frame	From 0 hour to 4 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Mean (Standard Deviation) Unit of Measure: Minutes	20.0 (42.0)	29.4 (63.0)	10.4 (23.3)

10. Duration When PI-NRS Scores Were >5 - From Post EP2 to 8 Hours Post-dose

Type: Secondary | Time Frame: Post EP2, every 5 minutes for the first hour, then every 15 minutes up to 8 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 2= first time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP2 is approximately a time point 4-5 hours post-dose. The duration of time that participants experienced PI-NRS score >5 from in period from post EP2 to 8 hours post-dose is reported.		
Time Frame	Post EP2, every 5 minutes for the first hour, then every 15 minutes up to 8 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Minutes	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

11. Duration When PI-NRS Scores Were >5 - From Post EP3 to 10 Hours Post-dose

Type: Secondary | Time Frame: Post EP3, every 5 minutes for the first hour, then every 15 minutes up to 10 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 3= second time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP3 is approximately a time point 8-9 hours post-dose. The duration of time that participants experienced PI-NRS score >5 from the post EP3 to 10 hours post dose.		
Time Frame	Post EP3, every 5 minutes for the first hour, then every 15 minutes up to 10 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	4	3	4
Mean (Standard Deviation) Unit of Measure: Minutes	0.0 (0.0)	0.0 (10.4)	0.0 (0.0)

12. Duration When PI-NRS Scores Were >5 - From Post EP4 to 28 Hours Post-dose

Type: Secondary | Time Frame: After EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose

Description	Participants rated severity of their pain at 11 point PI-NRS, by choosing a number between 0 and 10. Where 0= no pain and 10= worst possible pain; higher scores signify more pain. Evoked pain time point 4= third time point post-dose when pain was evoked using the participant specific heat pain stimulus. EP4 is approximately a time point 24-25 hours post-dose. The duration of time that participants experienced PI-NRS score >5 from post EP4 to 28 hours post-dose.		
Time Frame	After EP4, every 5 minutes for the first hour, then every 15 minutes up to 28 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. "Overall Number of Participants Analyzed" = participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	3	3	3
Mean (Standard Deviation) Unit of Measure: Minutes	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

13. Number of Participants With Participant's Global Satisfaction Score

Type: Secondary | Time Frame: At 4 hour post-dose or at time of first rescue therapy, whichever occurred first

Description	Participant was asked "How would you rate the study medication you received for pain?". The participant was provided the following choices as an answer: excellent=4; good=3; fair=2; poor=1. Response to this question, was participant's overall impression (global evaluation) of the study medication at 4 hour post-dose or at time of first rescue treatment or medication, which ever occurred first.
Time Frame	At 4 hour post-dose or at time of first rescue therapy, whichever occurred first
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication.

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Poor Score *Measure Type: Count of Participants Unit of Measure: Participants	1 20.0%	3 60.0%	0 0.0%
Fair Score *	2 40.0%	1 20.0%	3 60.0%
Good Score *	1 20.0%	0 0.0%	1 20.0%
Excellent Score *	1 20.0%	1 20.0%	1 20.0%
* Measure Type: Count of Participants Unit of Measure: Participants			

14. Time to First Use of Rescue Therapy or Medication

Type: Secondary | Time Frame: Up to maximum of 24 hours post-dose

Description	Time to rescue medication (hour) was calculated as: date/time of rescue medication minus date/time of first dose for each period. If participant who did not receive rescue medication, the time of censoring was cut off at 24 hours or the time of withdrawal, whichever was earlier. Kaplan-Meier method was used for estimation.		
Time Frame	Up to maximum of 24 hours post-dose		
Analysis Population Description	Full analysis set included all participants who successfully completed Part A, were randomized in Part B, and who received at least 1 dose of randomized study medication. Number of Participants Analyzed = number of participants evaluable for this outcome measure.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	2	2	2
Mean (90% Confidence Interval) Unit of Measure: Hours	14.9 ^[1] (1.6 to NA)	2.3 ^[1] (1.8 to NA)	15.3 ^[1] (1.6 to NA)
[1] Upper limit of 95% CI could not be estimated due to low number of participants.			

Statistical Analysis 1

Statistical Analysis Overview

Comparison Group Selection	Treatment Session 1: PF-05089771 1600 mg, Treatment Session 1: Placebo
----------------------------	--

Comments	[Not Specified]
Type of Statistical Test	Other
Comments	[Not Specified]

Statistical Test of Hypothesis

P-Value	1.0000
Comments	[Not Specified]
Method	Hierarchical rank test
Comments	[Not Specified]

Statistical Analysis 2

Statistical Analysis Overview

Comparison Group Selection	Treatment Session 2: PF-05089771 1600 mg, Treatment Session 2: Placebo
Comments	[Not Specified]
Type of Statistical Test	Other
Comments	[Not Specified]

Statistical Test of Hypothesis

P-Value	1.0000
Comments	[Not Specified]
Method	Hierarchical rank test
Comments	[Not Specified]

15. Area Under the Plasma Concentration-Time Curve From Time Zero to the 24 Hour Post-dose (AUC24) of PF-05089771

Type: Secondary | Time Frame: Predose, 0.5, 2, 4, 6, and 24 hours post-dose

Description	[Not Specified]	
Time Frame	Predose, 0.5, 2, 4, 6, and 24 hours post-dose	
Analysis Population Description	Pharmacokinetic (PK) parameter analysis set included all participants who were randomized, received study medication, and had at least 1 of the PK parameters of interest in at least 1 treatment session.	
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5
Geometric Mean (Geometric Coefficient of Variation) Unit of Measure: Nanogram*hour per milliliter (ng*hr/mL)	652100 (19%)	657500 (26%)

16. Area Under the Curve From Time Zero to Last Quantifiable Concentration (AUClast) of PF-05089771

Type: Secondary | Time Frame: Predose and 0.5, 2, 4, 6, and 24 hours post dose

Description	[Not Specified]	
Time Frame	Predose and 0.5, 2, 4, 6, and 24 hours post dose	
Analysis Population Description	PK parameter analysis set included all participants who were randomized, received study medication, and had at least 1 of the PK parameters of interest in at least 1 treatment session.	
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5
Geometric Mean (Geometric Coefficient of Variation) Unit of Measure: Nanogram*hour per milliliter (ng*hr/mL)	652400 (19%)	658000 (26%)

17. Maximum Observed Plasma Concentration (Cmax) of PF-05089771

Type: Secondary | Time Frame: Predose and 0.5, 2, 4, 6, and 24 hours post dose

Description	[Not Specified]	
Time Frame	Predose and 0.5, 2, 4, 6, and 24 hours post dose	
Analysis Population Description	PK parameter analysis set included all participants who were randomized, received study medication, and had at least 1 of the PK parameters of interest in at least 1 treatment session.	
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5
Geometric Mean (Geometric Coefficient of Variation) Unit of Measure: Nanogram per milliliter (ng/mL)	66760 (22%)	59080 (21%)

18. Time to Reach Maximum Observed Plasma Concentration (Tmax) of PF-05089771

Type: Secondary | Time Frame: Predose and 0.5, 2, 4, 6, and 24 hours post dose

Description	[Not Specified]	
Time Frame	Predose and 0.5, 2, 4, 6, and 24 hours post dose	
Analysis Population Description	PK parameter analysis set included all participants who were randomized, received study medication, and had at least 1 of the PK parameters of interest in at least 1 treatment session.	

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5
Median (Full Range) Unit of Measure: Hours	3.95 (2.00 to 6.07)	6.00 (4.00 to 6.13)

19. Number of Participants With Treatment Emergent Adverse Events (AEs) and Serious Adverse Events (SAEs)

Type: Secondary | Time Frame: Baseline up to a maximum of Day 83

Description	An AE was any untoward medical occurrence in a participant who received study drug without regard to possibility of causal relationship. SAE was an AE resulting in any of the following outcomes or deemed significant for any other reason: death; initial or prolonged inpatient hospitalization; life-threatening experience (immediate risk of dying); persistent or significant disability/incapacity; congenital anomaly. A treatment emergent AE was defined as an event that emerged during the treatment period that was absent before treatment, or worsened during the treatment period relative to the pretreatment state.		
Time Frame	Baseline up to a maximum of Day 83		
Analysis Population Description	Safety analysis set included all participants who received at least 1 dose of study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
AEs *Measure Type: Count of Participants Unit of Measure: Participants	5 100.0%	5 100.0%	5 100.0%
SAEs *	0 0.0%	0 0.0%	0 0.0%

* Measure Type: Count of Participants | Unit of Measure: Participants

20. Number of Participants With Laboratory Abnormalities

Type: Secondary | Time Frame: Baseline up to a maximum of Day 73

Description	Laboratory abnormalities: Hemoglobin (Hgb), hematocrit, red blood cell count: less than(<)0.8*lower limit of normal(LLN), platelet: <0.5*LLN/greater than (>)1.75*upper limit of normal (ULN), white blood cell: <0.6*LLN/>1.5*ULN, lymphocyte, neutrophil (absolute, %):<0.8*LLN/>1.2*ULN, total neutrophil <0.8*LLN;basophil, eosinophil, monocyte (absolute, %):>1.2*ULN; mean corpuscular (MC) volume, mean cell hemoglobin, MC hemoglobin concentration, mean platelet volume: <0.9*LLN/>1.1*ULN; total bilirubin >1.5*ULN, aspartate aminotransferase (AT), alanine AT, gammaglutamyl transferase, alkaline phosphatase:> 3.0*ULN, total protein, albumin: <0.8*LLN/>1.2*ULN; blood urea nitrogen, creatinine:>1.3*ULN, uric acid >1.2*ULN; sodium <0.95*LLN/>1.05*ULN, potassium, chloride, calcium, magnesium, bicarbonate: <0.9*LLN/>1.1*ULN; glucose <0.6*LLN/>1.5*ULN; urine (specific gravity <1.003/>1.030, pH <4.5/>8, glucose, ketone, protein, blood/Hgb, urobilinogen, bilirubin, nitrite, leukocyte esterase >=1).		
Time Frame	Baseline up to a maximum of Day 73		
Analysis Population Description	Safety analysis set included all participants who received at least 1 dose of study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Measure Type: Count of Participants Unit of Measure: Participants	2 40.0%	2 40.0%	2 40.0%

21. Number of Participants With Clinically Significant Changes From Baseline in Core Body Temperature

Type: Secondary | Time Frame: Baseline up to a maximum of Day 83

Description	The minimum starting temperature to measure core body temperature used was 33 degree Celsius. Clinically significant changes from baseline in core body temperature was judged by investigator.		
Time Frame	Baseline up to a maximum of Day 83		
Analysis Population Description	Safety analysis set included all participants who received at least 1 dose of study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Measure Type: Count of Participants Unit of Measure: Participants	0 0.0%	0 0.0%	0 0.0%

22. Number of Participants With Clinically Significant Changes From Baseline in Blood Pressure

Type: Secondary | Time Frame: Baseline up to a maximum of Day 83

Description	Criteria for clinically significant blood pressure abnormalities: systolic blood pressure >=30 millimetre of Mercury (mmHg) change from baseline in same posture, systolic blood pressure <90 mmHg, diastolic blood pressure >=20 mmHg change from baseline in same posture, diastolic blood pressure <50 mmHg.		
Time Frame	Baseline up to a maximum of Day 83		
Analysis Population Description	Safety analysis set included all participants who received at least 1 dose of study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Measure Type: Count of Participants Unit of Measure: Participants	00.0%	00.0%	00.0%

23. Number of Participants With Clinically Significant Changes From Baseline in Electrocardiogram (ECG)

Type: Secondary | Time Frame: Baseline up to a maximum of Day 83

Description	Criteria for clinically significant abnormalities in ECG : PR interval >=300 millisecond (msec) and 25 percent (%) increase when baseline >200 msec, 50% increase when baseline less than or equal to (<=) 200 msec; QRS interval >=140 msec, >=50% increase from baseline; QT interval >=500 msec; QT interval corrected using the Fridericia formula (QTcF) 450 msec to <480 msec, >=480 msec, 30 to <60 msec increase from baseline, >=60 msec increase from baseline.		
Time Frame	Baseline up to a maximum of Day 83		
Analysis Population Description	Safety analysis set included all participants who received at least 1 dose of study medication.		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Overall Number of Participants Analyzed	5	5	5
Measure Type: Count of Participants Unit of Measure: Participants	00.0%	00.0%	00.0%

Adverse Events ⓘ

Time Frame	[Not Specified]
Adverse Event Reporting Description	The same event may appear as both an AE and a SAE. However, what is presented are distinct events. An event may be categorized as serious in 1 participant and as nonserious in another participant, or 1 participant may have experienced both a serious

and nonserious event during the study. Adverse events were collected only for Part B of the study.

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.

[Expand all](#) / [Collapse all](#)

All-Cause Mortality

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
	Affected / at Risk (%)	Affected / at Risk (%)	Affected / at Risk (%)
Total	--/--	--/--	--/--

Serious Adverse Events

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
	Affected / at Risk (%)	Affected / at Risk (%)	Affected / at Risk (%)
Total	0/5 (0.00%)	0/5 (0.00%)	0/5 (0.00%)

Other (Not Including Serious) Adverse Events

Frequency Threshold for Reporting Other Adverse Events	1%		
Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
	Affected / at Risk (%)	Affected / at Risk (%)	Affected / at Risk (%)
Total	5/5 (100.00%)	5/5 (100.00%)	5/5 (100.00%)

Gastrointestinal disorders

Abdominal pain * 1	0/5 (0.00%)	0/5 (0.00%)	0/5 (0.00%)
Chapped lips * 1	1/5 (20.00%)	0/5 (0.00%)	0/5 (0.00%)
Constipation * 1	1/5 (20.00%)	0/5 (0.00%)	1/5 (20.00%)
Diarrhoea * 1	0/5 (0.00%)	0/5 (0.00%)	1/5 (20.00%)
Paraesthesia oral * 1	4/5 (80.00%)	0/5 (0.00%)	3/5 (60.00%)

General disorders

Fatigue * 1	0/5 (0.00%)	0/5 (0.00%)	2/5 (40.00%)
Feeling cold * 1	0/5 (0.00%)	0/5 (0.00%)	1/5 (20.00%)
Feeling hot * 1	0/5 (0.00%)	1/5 (20.00%)	0/5 (0.00%)

Infections and infestations

Arm/Group Title	Treatment Session 1: PF-05089771 1600 mg	Treatment Session 1: Placebo	Treatment Session 2: PF-05089771 1600 mg
Arm/Group Description	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 1.	Participants received oral dispersion of placebo matched to PF-05089771 in Treatment session 1.	Participants received oral dispersion of PF-05089771 1600 mg in Treatment session 2.
Chronic sinusitis * 1	0/5 (0.00%)	0/5 (0.00%)	1/5 (20.00%)
Upper respiratory tract infection * 1	1/5 (20.00%)	1/5 (20.00%)	0/5 (0.00%)
Urinary tract infection * 1	1/5 (20.00%)	1/5 (20.00%)	0/5 (0.00%)

Investigations

Hepatic enzyme increased * 1	1/5 (20.00%)	0/5 (0.00%)	1/5 (20.00%)
------------------------------	--------------	-------------	--------------

Musculoskeletal and connective tissue disorders

Back pain * 1	1/5 (20.00%)	0/5 (0.00%)	1/5 (20.00%)
Pain in extremity * 1	1/5 (20.00%)	1/5 (20.00%)	0/5 (0.00%)

Neoplasms benign, malignant and unspecified (incl cysts and polyps)

Skin papilloma * 1	1/5 (20.00%)	1/5 (20.00%)	0/5 (0.00%)
--------------------	--------------	--------------	-------------

Nervous system disorders

Dizziness * 1	2/5 (40.00%)	0/5 (0.00%)	1/5 (20.00%)
Dysgeusia * 1	1/5 (20.00%)	0/5 (0.00%)	0/5 (0.00%)
Headache * 1	1/5 (20.00%)	0/5 (0.00%)	0/5 (0.00%)
Hyperaesthesia * 1	1/5 (20.00%)	0/5 (0.00%)	1/5 (20.00%)
Paraesthesia * 1	2/5 (40.00%)	2/5 (40.00%)	3/5 (60.00%)

Reproductive system and breast disorders

Dysmenorrhoea * 1	0/5 (0.00%)	0/5 (0.00%)	1/5 (20.00%)
-------------------	-------------	-------------	--------------

Respiratory, thoracic and mediastinal disorders

Nasal dryness * 1	0/5 (0.00%)	1/5 (20.00%)	0/5 (0.00%)
-------------------	-------------	--------------	-------------

Skin and subcutaneous tissue disorders

Ecchymosis * 1	1/5 (20.00%)	1/5 (20.00%)	1/5 (20.00%)
Pruritus * 1	2/5 (40.00%)	0/5 (0.00%)	0/5 (0.00%)

Vascular disorders

Flushing * 1	4/5 (80.00%)	0/5 (0.00%)	1/5 (20.00%)
--------------	--------------	-------------	--------------

* Indicates events were collected by non-systematic assessment

1 Term from vocabulary, MedDRA version 16.0

Limitations and Caveats

[Not Specified]

Collaborators and Investigators

This is where you will find people and organizations involved with this study.

Sponsor ⓘ

Pfizer

Investigators ⓘ

- Study Director: Pfizer CT.gov Call Center, Pfizer

Publications

From PubMed

These publications are automatically filled in from PubMed, a public database of scientific and medical articles, and may or may not be about the study.

- [Cao L, McDonnell A, Nitzsche A, Alexandrou A, Saintot PP, Loucif AJ, Brown AR, Young G, Mis M, Randall A, Waxman SG, Stanley P, Kirby S, Tarabar S, Gutteridge A, Butt R, McKernan RM, Whiting P, Ali Z, Bilsland J, Stevens EB. Pharmacological reversal of a pain phenotype in iPSC-derived sensory neurons and patients with inherited erythromelalgia. Sci Transl Med. 2016 Apr 20;8\(335\):335ra56. doi: 10.1126/scitranslmed.aad7653. \(https://pubmed.ncbi.nlm.nih.gov/27099175\).](#)

More Information

Record History

Certain Agreements ⓘ

Principal Investigators are NOT employed by the organization sponsoring the study.

There IS an agreement between Principal Investigators and the Sponsor (or its agents) that restricts the PI's rights to discuss or publish trial results after the trial is completed

Pfizer has the right to review disclosures, requesting a delay of less than 60 days. Investigator will postpone single center publications until after disclosure of pooled data (all sites), less than 12 months from study completion/termination at all participating sites. Investigator may not disclose previously undisclosed confidential information other than study results.