

*Isolation and characterization of 29  
microsatellite markers for the bumphead  
parrotfish, *Bolbometopon muricatum*, and  
cross amplification in 12 related species*


**Mark A. Priest, Glenn R. Almany,  
Camrin D. Braun, Richard J. Hamilton,  
Diego F. Lozano-Cortés, Pablo Saenz-  
Agudelo, et al.**

**Marine Biodiversity**

ISSN 1867-1616

Mar Biodiv

DOI 10.1007/s12526-014-0278-4



**Your article is protected by copyright and all rights are held exclusively by Senckenberg Gesellschaft für Naturforschung and Springer-Verlag Berlin Heidelberg. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at [link.springer.com](http://link.springer.com)".**

# Isolation and characterization of 29 microsatellite markers for the bumphead parrotfish, *Bolbometopon muricatum*, and cross amplification in 12 related species

Mark A. Priest · Glenn R. Almany · Camrin D. Braun ·  
Richard J. Hamilton · Diego F. Lozano-Cortés ·  
Pablo Saenz-Agudelo · Michael L. Berumen

Received: 30 June 2014 / Revised: 30 September 2014 / Accepted: 7 October 2014  
© Senckenberg Gesellschaft für Naturforschung and Springer-Verlag Berlin Heidelberg 2014

**Abstract** We isolated and characterized 29 microsatellite loci for the bumphead parrotfish, *Bolbometopon muricatum*, a wide-ranging parrotfish listed as vulnerable by the International Union for Conservation of Nature (IUCN). The 29 loci were tested on 95 individuals sampled from the Solomon Islands. The number of alleles ranged from two to ten. Evidence of linkage disequilibrium was found for only one pair of loci (Bm54 and Bm112). Two loci (Bm20 and Bm119) showed significant departure from Hardy-Weinberg equilibrium. We also tested each locus for amplification and polymorphism on 11 other scarine labrid species and one labrid species. Amplification success ranged from zero to ten loci per species. These microsatellite loci are the first specific

set for *B. muricatum* and will be a useful tool for assessing genetic population structure, genetic diversity, and parentage in future studies.

**Keywords** Bumphead parrotfish · Scarine Labridae · Microsatellites · Population genetics

## Introduction

The bumphead parrotfish, *Bolbometopon muricatum*, is the world's largest parrotfish species with a geographic range encompassing coral reefs from the Red Sea to French Polynesia (Hamilton and Choat 2012a). Where they are present, *B. muricatum* are responsible for a large amount of bioerosion on coral reefs, removing, on average, five tonnes per individual of carbonate annually, half of which is live coral (Bellwood et al. 2003). In addition to this vital ecosystem function, this species is also important in both subsistence and small scale commercial fisheries (Aswani and Hamilton 2004), yet *B. muricatum* is highly susceptible to overfishing due to life history characteristics of large size, slow growth, and the fact that this species aggregates in shallow water for feeding, spawning and sleeping activities (Hamilton et al. 2008). Consequently, continued declines in abundance throughout its range (Dulvy and Polunin 2004; Bellwood and Choat 2011), have resulted in *B. muricatum* being listed as vulnerable to extinction by the International Union for Conservation of Nature (IUCN) (Chan et al. 2012).

Previous genetic studies on parrotfishes have generally focused on phylogeography to gain insights into the evolutionary history of species (e.g., Bay et al. 2004; Winters et al. 2010; Fitzpatrick et al. 2011). However, effective management of exploited populations requires knowledge of demographically relevant connectivity over ecological time scales,

M. A. Priest (✉) · C. D. Braun · D. F. Lozano-Cortés ·  
P. Saenz-Agudelo · M. L. Berumen  
Red Sea Research Center, King Abdullah University of Science and  
Technology, Thuwal 23955, Kingdom of Saudi Arabia  
e-mail: m.priest@uq.edu.au

M. A. Priest  
Marine Spatial Ecology Lab, School of Biological Sciences,  
University of Queensland, St. Lucia Campus, Brisbane QLD 4072,  
Australia

G. R. Almany  
Laboratoire d'Excellence "CORAIL," USR 3278 CNRS-EPHE  
CRILOBE, Perpignan, France

C. D. Braun  
Biology Department, Woods Hole Oceanographic Institution,  
Woods Hole, Massachusetts 02540, USA

R. J. Hamilton  
The Nature Conservancy, Asia Pacific Division, 245 Riverside Drive,  
West End QLD 4101, Australia

P. Saenz-Agudelo  
Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral  
de Chile, Valdivia, Chile

such as the number of migrants exchanged between spatially segregated habitat patches (see review Jones et al. 2007). These direct measurements of dispersal can now be revealed using highly polymorphic co-dominant microsatellite markers in conjunction with recent statistical advances (Manel et al. 2005; Selkoe and Toonen 2006).


Here, we present 29 novel microsatellite loci for *B. muricatum* that will help determine key population parameters (i.e., genetic population structure, larval dispersal pathways) that are essential for the effective conservation and management of this species.

## Materials and methods

All genomic DNA used in this study was obtained from fin clip tissues using DNeasy extraction kits (Qiagen) following the manufacture's protocol. A genomic library was created using 454 GS FLX titanium shotgun sequencing on one individual sample by the Bioscience Core Lab at the King Abdullah University of Science and Technology. Over 150,000 reads with an average length of 230 bp were generated. Microsatellites were mined from the generated 454 data using the software MSATCOMMANDER v 1.0.8 (Faircloth 2008). Specifically, we searched for dinucleotide and trinucleotide motifs with a minimum of eight perfect repeats. We found 1,748 sequences that contained dinucleotide (1,501) or trinucleotide (248) repeats. Among those, 187 had flanking regions for which primer pairs could be designed. Overall, 120 microsatellite loci were selected for polymerase chain

reaction (PCR) trials (108 dinucleotide and 12 trinucleotide repeats). PCR reactions were set up following protocols associated with the Multiplex PCR kit (Qiagen). Primers were tested at annealing temperatures ranging from 55 to 63 °C. PCR reaction volume was 10 µl, consisting of 5 µL of Multiplex Mix (Qiagen), 1 µL of primers (2 µM), 3 µL of water, and 1 µL of genomic DNA (30–100 ng/µl). PCR reactions were performed using the following parameters: 15 min at 95 °C followed by 30 cycles of 30 s at 94 °C, 90 s at 55–63 °C and 90 s at 72 °C, with a final extension of 10 min at 72 °C. The 120 designed primer pairs were first tested on eight samples to identify polymorphic loci. PCR products were run on a QIAxcel genetic analyzer (Qiagen) using a high-resolution cartridge to identify polymorphic loci. Of the 120 loci tested, 29 exhibited variation and were labeled with ABI fluorescent tags: 6-FAM, PET, VIC, and NED (Fig. 1). PCR reactions were performed as described previously for 95 individuals sampled from one location in the Solomon Islands (Hamilton and Choat 2012b; Hamilton et al. 2013).

PCR products were analyzed with an ABI 3730xl genetic analyzer (Applied Biosystems) and sizes were determined with GENEMAPPER 4.0 (Applied Biosystems). Number of alleles and expected heterozygosities under Hardy-Weinberg equilibrium were calculated using GENALEX v6.5 (Peakall and Smouse 2012). Tests for Hardy-Weinberg and linkage disequilibrium were conducted using genepop on the web v4.2 (Raymond and Rousset 1995; Rousset 2008), with significance levels adjusted for multiple comparisons using false discovery rates (FDR; Benjamini et al. 2006).



**Fig. 1** Schematic of four microsatellite multiplexes designed for *Bolbometopon muricatum* representing allelic size range and fluorescent dyes used for each locus

**Table 1** Primer sequences, repeat motifs and characteristics of 29 microsatellites loci developed for *Bolbometopon muricatum*: annealing temperature (Ta), number of alleles (Na), observed (Ho) and expected (He) heterozygosities, and probability value of Hardy-Weinberg equilibrium test (*p*)

Locus	Primer sequence (5'-3')	Dye	Repeat motif	Ta (°C)	Size range (bp)	Multiplex	Na	Ho	He	<i>p</i> *	Genbank accession number
Bm20	F: TTTCCTGTCTGGAGGGAGCAG R: GAGGGCTTGTGGATGGTG	NED	(AC)9	60	94–102	A	4	0.616	0.649	<b>0.002</b>	KJ489028
Bm42	F: AACCACATACAAACACGGCG R: TCGTGTCTCTCTGTAAACCTG	6-FAM	(AC)17	60	113–121	A	5	0.463	0.401	0.986	KJ489029
Bm29	F: ACATGGCTCTTTGGCTCAC R: GAGTGGGTAGGCTTGTGTG	PET	(AC)15	60	116–130	A	5	0.406	0.369	0.832	KJ489030
Bm43	F: AGTTTGTGCATACGTGGC R: TTCCAGTGTCCAGGGTTTC	VIC	(AC)10	60	130–144	A	3	0.235	0.244	0.462	KJ489031
Bm37	F: AATCCTGGAGTGACAGCAG R: TGAGAGTTTCTGTGGAGGGC	NED	(AC)9	60	176–180	A	3	0.455	0.455	0.507	KJ489032
Bm72	F: CAAGCAGTGGTGTGTGGTGC R: GTGTGCATGTCCCTGTTTAC	6-FAM	(AC)11	60	201–209	A	4	0.136	0.130	1.000	KJ489033
Bm98	F: GTCTGAGGCGAAGTGTGTG R: GCTCAGCTTCATTCGGACAG	PET	(AC)8	60	206–208	A	2	0.049	0.048	1.000	KJ489034
Bm75	F: GTTGGGTTTCAGTGCCATCC R: TGGCGTCTTGATCGTGGTC	VIC	(AC)14	60	200–214	A	5	0.651	0.583	0.917	KJ489035
Bm105	F: TTGGGATCAGGTACCGAGC R: TGTGTGTAATTGCCCGTGAC	VIC	(AC)8	60	346–354	A	2	0.395	0.396	0.576	KJ489036
Bm88	F: CCTTCATGTCTTTGGCAGG R: CAACACAGCCCTATGTAGCTG	6-FAM	(AC)12	60	127–135	B	3	0.474	0.484	0.632	KJ489037
Bm118	F: AACAACTACAGAGGAGAGCAC R: TGGTATGTCTTTGTATCCGCC	NED	(AAT)17	60	104–128	B	8	0.782	0.790	0.353	KJ489038
Bm8	F: TAAGAAAGGAGAGTGCAGGC R: TGTGAGTCATGTAGGGCCTG	NED	(AC)12	60	205–213	B	5	0.699	0.699	0.500	KJ489039
Bm108	F: CATCTGTCAAGGGCTCCAAC R: TATCTCTCGTGCAGGTGTG	PET	(AT)8	60	242–244	B	2	0.397	0.431	0.312	KJ489040
Bm14	F: CAGTCCAGTGTCTGCTTTGG R: GGAGTCCAGTACCAGCAGTC	VIC	(AC)13	60	219–229	B	5	0.714	0.725	0.332	KJ489041
Bm76	F: TGTCTGCCACCTACATGAC R: TTGCCCAAGAAGTCGACTGC	NED	(AC)9	60	271–277	C	4	0.154	0.155	0.466	KJ489042
Bm25	F: AACACTGTGACAAACACCCG R: TCCTGTCTTGCCTTGATGAG	NED	(AC)10	60	150–166	C	3	0.211	0.193	1.000	KJ489043
Bm32	F: GAGTCTTCGTGCTTGATGG R: AGGGAGCTGACACAACATCC	6-FAM	(AC)9	60	167–169	C	2	0.110	0.104	1.000	KJ489044
Bm87	F: AACAGTGTCAATGTGGCTGAC	PET	(AC)12	60	168–178	C	5	0.658	0.675	0.471	KJ489045

**Table 1** (continued)

Locus	Primer sequence (5'-3')	Dye	Repeat motif	Ta (°C)	Size range (bp)	Multiplex	Na	Ho	He	p*	Genbank accession number
Bm109	R: TGTTCGCTCTTCTAGGCTTGG F: GCTTCTGCCATGATAACAACTC	VIC	(AAG)11	60	143–158	C	4	0.159	0.167	0.429	KJ489046
Bm45	R: TGGAAATAGGGACCTCTCGC F: GAGGAGAACGAAGAGAGCCCC	6-FAM	(AC)10	60	234–248	C	5	0.620	0.575	0.862	KJ489047
Bm54	R: GTGTGTGTACTCTGCAGCTG F: CTTGAGCCGGCTGTGTTAAG	PET	(AG)12	60	250–260	C	4	0.187	0.233	0.040	KJ489048
Bm17	R: TATCTCCAGTCCAGCACAGC F: AACTGAGGGTACTGGTGCTG	VIC	(AC)11	60	260–264	C	3	0.527	0.487	0.811	KJ489049
Bm79	R: TCGGATAGAAAGTCAGCCTGG F: AACACACACATTTCTCTGCATG	PET	(AC)9	60	275–291	D	5	0.717	0.674	0.574	KJ489050
Bm81	R: GAGACTGCCATCTAGAGGCG F: CGGCTGTTCCAGTAGATTCC	6-FAM	(AC)11	60	278–294	D	6	0.244	0.245	0.552	KJ489051
Bm112	R: CTGCTTCATCCCTCACCC F: AGATGCCAGTATTATGCAGGTG	NED	(AAT)12	60	256–289	D	8	0.670	0.653	0.592	KJ489052
Bm110	R: TGGGATTCTGTGTACAACTACG F: GGTCCTGTCTGTTTATCAAAAGC	VIC	(AAT)11	60	177–192	D	5	0.298	0.303	0.439	KJ489053
Bm116	R: CTGGGCAGACACACTTCAAC F: GGGTCTGGGATTAGGGTAGG	NED	(AAC)14	60	137–152	D	6	0.613	0.671	0.323	KJ489054
Bm119	R: ATGAGGTCAGAGGTCAGAGC F: GAAGATGACGTGACGCTGAG	PET	(AAT)17	60	181–211	D	10	0.274	0.795	<b>0.000</b>	KJ489055
Bm120	R: TGAACAGAGGGATTACAGCG F: GCAATTTCTTAAGCCTCTCAGC	6-FAM	(AAT)19	60	180–204	D	8	0.761	0.728	0.955	KJ489056
	R: CTTACTGTACTCAAGTCTCTGCC										

\*Numbers in bold indicate significant departures from expected Hardy-Weinberg equilibrium after correction for multiple tests



Finally, all 29 loci were tested for amplification and polymorphism on 11 other scarine labrid species (three individuals per species), and *Cheilinus undulatus*, using the same PCR and genotyping conditions described above.

## Results and discussion

Allelic diversity was generally low, ranging from two to ten (mean allelic diversity 4.621). Observed and expected heterozygosities ranged from 0.049 to 0.782, and 0.048 to 0.795, respectively (mean observed and expected heterozygosities were 0.437 and 0.450, respectively) (Table 1). Two loci

(Bm20, and Bm119) showed significant departure from Hardy-Weinberg equilibrium after corrections for multiple testing [FDR,  $q < 0.05$ ] due to heterozygote deficiency. Only one pair of loci showed evidence of linkage based on the genotypic disequilibrium test using the log likelihood ratio statistic implemented in Genepop (Bm54 and Bm112,  $p < 0.001$  after FDR correction,  $q < 0.05$ ). Allelic diversity was generally low compared to microsatellites identified for some other coral reef fishes (e.g., *Chaetodon vagabundus*, Almany et al. 2009; *Dascyllus marginatus*, Aluana et al. 2012; but see also *Coris bulbifrons*, van der Meer et al. 2013). This could be a result of testing our markers on individuals from only one geographic location, and further testing may reveal greater

**Table 2** Cross-amplification success of 29 microsatellite loci developed for *Bolbometopon muricatum* when tested on 11 other scarine labrid species (three individuals per species)

Locus	Species										
	<i>Calatomus viridescens</i>	<i>Cetoscarus bicolor</i>	<i>Chlorurus sordidus</i>	<i>Scarus arabicus</i>	<i>Scarus collana</i>	<i>Scarus fuscopurpureus</i>	<i>Scarus ghobban</i>	<i>Scarus niger</i>	<i>Scarus rubroviolaceus</i>	<i>Chlorurus gibbus</i>	<i>Hipposcarus harid</i>
Bm20	–	++	–	–	–	–	–	–	+	–	–
Bm42	++	+	–	–	–	–	–	–	+	+	–
Bm29	–	–	–	–	–	–	–	–	–	+	–
Bm43	+	–	–	–	–	–	–	+	–	+	–
Bm37	+	+	–	–	–	–	–	–	–	–	–
Bm72	++	++	+	+	+	+	++	++	++	++	++
Bm98	+	++	–	+	++	–	++	+	++	++	++
Bm75	+	–	–	–	–	–	–	–	–	+	+
Bm105	–	–	–	–	–	–	–	–	–	–	–
Bm88	–	–	–	–	–	–	–	–	–	–	–
Bm118	–	–	–	–	–	–	–	–	–	–	–
Bm8	–	+	–	–	–	–	–	–	–	–	–
Bm108	–	–	–	–	–	–	–	–	–	–	–
Bm14	–	–	–	–	–	–	–	–	–	–	+
Bm76	++	++	–	–	–	–	–	–	–	–	–
Bm25	–	–	–	–	–	–	–	–	–	–	–
Bm32	++	+	++	–	–	++	++	+	++	–	++
Bm87	–	–	–	–	–	–	–	–	–	–	–
Bm109	–	–	–	–	–	–	–	–	–	–	–
Bm45	–	++	+	–	–	–	–	–	++	–	–
Bm54	–	–	–	–	–	–	–	–	+	–	–
Bm17	–	++	+	–	–	+	–	–	+	–	+
Bm79	–	–	–	–	–	–	–	–	–	–	–
Bm81	–	–	–	–	–	–	–	–	–	–	–
Bm112	–	–	–	–	–	–	–	–	–	–	–
Bm110	–	–	–	–	–	–	–	–	–	–	–
Bm116	–	–	–	–	–	–	–	–	–	–	–
Bm119	–	–	–	–	–	–	–	–	–	–	–
Bm120	–	–	–	–	–	–	–	–	–	–	–

– denotes no amplification; + denotes one observed allele; ++ denotes multiple alleles

range-wide allelic diversity. Nonetheless, given the large number of markers presented here, parentage and kinship, and thus dispersal, can still be confidently estimated (Harrison et al. 2013).

Cross-amplification tests revealed contrasting results among several other species. Amplification success per species in the scarine labrids ranged from two loci (in *Scarus arabicus* and *Scarus collana*) to ten loci (in *Cetoscarus bicolor*), with successful amplification of locus Bm72 in all species (Table 2). Thus, these markers may also be useful for further studies on other scarine labrid species. We also tested all loci on the humphead wrasse, *Cheilinus undulatus*, as this species is more closely related to large parrotfishes than many of the smaller wrasse families (Kazancıoğlu et al. 2009), and *C. undulatus* is also listed as endangered by the IUCN (Russell 2004). However, no amplification success was observed at any locus.

The microsatellite loci presented here are the first specific set for *B. muricatum* and will be a useful tool for evaluating many key population parameters, such as stock structure, and larval dispersal distances. This information is essential for improved management and effective conservation of this functionally and commercially important coral reef fish.

**Acknowledgments** The authors wish to thank the Biosciences Core Lab of the King Abdullah University of Science and Technology (KAUST) for technical assistance, Hugo Harrison for the repurposing of Fig. 1, and three anonymous reviewers for helpful comments. Funding was provided in part by KAUST through baseline research funding and through award number CRG-1-BER-002 to MLB.

## References

- Almany GR, De Arruda MP, Arthofer W, Atallah ZK, Beissinger SR, Berumen ML, Bogdanowicz SM et al (2009) Permanent genetic resources added to molecular ecology resources database 1 May 2009–31 July 2009. *Mol Ecol Resour* 9(6):1460–1466
- Aluana AG, Albaina A, Alpermann TJ, Apkenas VE, Bankhead-Dronnet S, Bergek S, Berumen ML et al (2012) Permanent genetic resources added to molecular ecology resources database 1 October 2011–30 November 2011. *Mol Ecol Resour* 12(2):374–376
- Aswani S, Hamilton RJ (2004) Integrating indigenous ecological knowledge and customary sea tenure with marine and social science for conservation of bumphead parrotfish (*Bolbometopon muricatum*) in the Roviana Lagoon, Solomon Islands. *Environ Conserv* 31(1):69–83
- Bay LK, Choat JH, van Herwerden L, Robertson DR (2004) High genetic diversities and complex genetic structure in an Indo-Pacific tropical reef fish (*Chlorurus sordidus*): evidence of an unstable evolutionary past? *Mar Biol* 144(4):757–767
- Bellwood DR, Choat JH (2011) Dangerous demographics: the lack of juvenile humphead parrotfishes *Bolbometopon muricatum* on the Great Barrier Reef. *Coral Reefs* 30:549–554
- Bellwood DR, Hoey AS, Choat JH (2003) Limited functional redundancy in high diversity systems: resilience and ecosystem function on coral reefs. *Ecol Lett* 6:281–285
- Benjamini Y, Krieger AM, Yekutieli D (2006) Adaptive linear step-up procedures that control the false discovery rate. *Biometrika* 93(3):491–507
- Chan T, Sadovy Y, Donaldson TJ (2012) *Bolbometopon muricatum*. In IUCN 2013. IUCN red list of threatened species. Version 2013.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 03 January 2014
- Dulvy NK, Polunin NVC (2004) Using informal knowledge to infer human-induced rarity of a conspicuous reef fish. *Anim Conserv* 7(4):365–374
- Faircloth BC (2008) MSATCOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Mol Ecol Resour* 8(1):92–94
- Fitzpatrick JM, Carlon DB, Lippe C, Robertson DR (2011) The West Pacific diversity hotspot as a source or sink for new species? Population genetic insights from the Indo Pacific parrotfish *Scarus rubroviolaceus*. *Mol Ecol* 20:219–234
- Hamilton RJ, Choat JH (2012a) Bumphead Parrotfish—*Bolbometopon muricatum*. In: Sadovy de Mitcheson Y, Colin PL (eds) Reef fish spawning aggregations: biology, fisheries and management. Springer, Dordrecht
- Hamilton RJ, Choat JH (2012b) Establishing the sustainability of the bumphead parrotfish (*Bolbometopon muricatum*) spear fishery in Isabel Province, Solomon Islands. *Sci Conserv Fish Aggregations Newsl* 16:10
- Hamilton RJ, Adams S, Choat JH (2008) Sexual development and reproductive demography of the green humphead parrotfish (*Bolbometopon muricatum*) in the Solomon Islands. *Coral Reefs* 27:153–163
- Hamilton RJ, Almany G, Stevens D, Pita J (2013) Establishing the population size of adult bumphead parrotfish (*Bolbometopon muricatum*) in Isabel Province, Solomon Islands. *Sci Conserv Fish Aggregations Newsl* 17:13
- Harrison HB, Saenz-Agudelo P, Planes S, Jones GP, Berumen ML (2013) Relative accuracy of three common methods of parentage analysis in natural populations. *Mol Ecol* 22:1158–1170
- Jones GP, Srinivasan M, Almany GR (2007) Population connectivity and conservation of marine biodiversity. *Oceanography* 20(3):100–111
- Kazancıoğlu E, Near TJ, Hanel R, Wainwright PC (2009) Influence of sexual selection and feeding functional morphology on diversification rate of parrotfishes (Scaridae). *Proc R Soc B* 276(1672):3439–3446
- Manel S, Gaggiotti OE, Waples RS (2005) Assignment methods: matching biological questions with appropriate techniques. *Trends Ecol Evol* 20(3):136–142
- Peakall R, Smouse PE (2012) GenA1Ex 6.5: genetic analysis in excel. Population genetic software for teaching and research—an update. *Biodivers Inform* 28(19):2537–2539
- Raymond M, Rousset F (1995) GENPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J Hered* 86(3):248–249
- Rousset F (2008) Genepop'007: a complete re-implementation of the GENEPop software for windows and linux. *Mol Ecol Resour* 8(1):103–106
- Russell B (Grouper & Wrasse Specialist Group) (2004) *Cheilinus undulatus*. In IUCN 2013. IUCN red list of threatened species. Version 2013.2. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 11 March 2014
- Selkoe KA, Toonen RJ (2006) Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecol Lett* 9(5):615–629
- van der Meer MH, Gardner MG, Berumen ML, Hobbs J-PA, van Herwerden L (2013) Identification of seventeen microsatellite loci for conservation genetic studies of the endemic wrasse *Coris bulbifrons*. *Conserv Genet Resour* 5(2):363–366
- Winters KL, van Herwerden L, Choat JH, Robertson DR (2010) Phylogeography of the indo-pacific parrotfish *Scarus psittacus*: isolation generates distinctive peripheral populations in two oceans. *Mar Biol* 157(8):1679–1691