

Math 2000 Tutorial Worksheet

December 2nd, 2015

1. (Section 7.1 #2) Let $A = \{a, b, c\}$ and let $R = \{(a, a), (a, c), (b, b), (b, c), (c, a), (c, b)\}$ define a relation on A . Are the following statements true or false? Explain.
 - (a) For each $x \in A$, $x R x$.
 - (b) For every $x, y \in A$, if $x R y$, then $y R x$.
 - (c) For every $x, y, z \in A$, if $x R y$ and $y R z$, then $x R z$.
 - (d) R is a function from A to A .

2. (Section 7.1 #4) Let U be a nonempty set, and let R be the “subset relation” on $\mathcal{P}(U)$. That is,

$$R = \{(S, T) \in \mathcal{P}(U) \times \mathcal{P}(U) \mid S \subseteq T\}.$$

- (a) Write the open sentence $(S, T) \in R$ using standard subset notation.
 - (b) What is the domain of the relation R ?
 - (c) What is the range of the relation R ?
 - (d) Is R a function from $\mathcal{P}(U)$ to $\mathcal{P}(U)$? Explain.
3. (Section 7.1#5) Repeat parts (b)-(d) of Problem 2 for the “element of” relation

$$R = \{(x, S) \in U \times \mathcal{P}(U) \mid x \in S\}.$$

4. (Section 7.1 #6 and 7) Let $S = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid x^2 + y^2 = 100\}$.
 - (a) Determine the set of all values of x such that $(x, 6) \in S$, and determine the set of all values of x such that $(x, 9) \in S$.
 - (b) Determine the domain and range of the relation S , and write each set using set builder notation.
 - (c) Is the relation on S a function from \mathbb{R} to \mathbb{R} ? Explain.
 - (d) Since S is a relation on \mathbb{R} , its elements can be graphed in the coordinate plane. Describe the graph of the relation S . Is the graph consistent with your answers in parts (a) - (c)? Explain.
 - (e) Repeat parts (a) - (d) for the relation $T = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid y = \sqrt{100 - x^2}\}$. What is the connection between the relations S and T ?

5. (Section 7.1 #8) Determine the domain and range of each of the following relations on \mathbb{R} and sketch the graph of each relation.
- (a) $R = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid x^2 + y^2 = 100\}$
 - (b) $S = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid y^2 = x + 10\}$
 - (c) $T = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid |x| + |y| = 10\}$
 - (d) $W = \{(x, y) \in \mathbb{R} \times \mathbb{R} \mid x^2 = y^2\}$.
6. (Section 7.2 #2) Let $A = \{a, b, c\}$. For each of the following, draw a directed graph that represents a relation on A with the specified properties:
- (a) A relation on A that is symmetric but not transitive.
 - (b) A relation on A that is transitive but not symmetric.
 - (c) A relation on A that is symmetric and transitive but not reflexive.
 - (d) A relation on A that is not reflexive on A , is not symmetric, and is not transitive.
 - (e) A relation on A , other than the identity relation, that is an equivalence relation on A .
7. (Section 7.2 #5) A relation R is defined on \mathbb{Z} as follows: For all $a, b \in \mathbb{Z}$, $a R b$ if and only if $|a - b| \leq 3$. Is R an equivalence relation on \mathbb{Z} ? If not, is R reflexive? Symmetric? Transitive? Justify all conclusions.
8. (Section 7.2 #9) Define the relation \sim on \mathbb{Q} as follows: For $a, b \in \mathbb{Q}$, $a \sim b$ if and only if $a - b \in \mathbb{Z}$.
- (a) Show that \sim is an equivalence relation. (See Progress Check 7.9.)
 - (b) List four different elements of the set $C = \left\{x \in \mathbb{Q} \mid x \sim \frac{5}{7}\right\}$.
 - (c) Use set builder notation, without using the symbol \sim , to specify the set C .
 - (d) use the roster method to specify the set C .
9. (Section 7.2 #13) Let \sim and \approx be relations on \mathbb{Z} defined as follows;
- For $a, b \in \mathbb{Z}$, $a \sim b$ if and only if $2a + 3b \equiv 0 \pmod{5}$.
 - For $a, b \in \mathbb{Z}$, $a \approx b$ if and only if $a + 3b \equiv 0 \pmod{5}$.
- (a) Is \sim an equivalence relation on \mathbb{Z} ? If not, is this relation reflexive? Symmetric? Transitive?
 - (b) Is \approx an equivalence relation on \mathbb{Z} ? If not, is this relation reflexive? Symmetric? Transitive?