

Practice problems for change of variables, including
Quizzes 14 and 15
Math 2580
Spring 2016

Sean Fitzpatrick

March 1st, 2016

For Quiz 14 on Tuesday, you should be able to do the following:

1. Let D be the region bounded by the polar curve $r = 2 \sin \theta$.
 - (a) Find the area of D . (Caution: first identify the curve, and choose your limits of integration accordingly.)
 - (b) Find the volume of the solid that lies above the region D and below the graph of the function $f(x, y) = 4 - \sqrt{x^2 + y^2}$.
2. Evaluate $\iint_D e^{x^2+y^2} dA$, where D is the region bounded by the circles $x^2 + y^2 = 4$ and $x^2 + y^2 = 9$.
3. The volume of a solid T is given in cylindrical coordinates by $\int_0^{\pi/2} \int_0^2 \int_0^{4-r^2} r \, dz \, dr \, d\theta$. Sketch the solid and compute its volume.
4. Evaluate the integral $\iiint_W \frac{1}{\sqrt{x^2+y^2}} dV$, where W is the region given by $0 \leq x \leq \sqrt{9-y^2}$, $0 \leq y \leq 3$, $0 \leq z \leq \sqrt{9-(x^2-y^2)}$, using cylindrical coordinates.
5. Find the volume that the cylinder $r = a \cos \theta$ cuts out of the sphere $x^2 + y^2 + z^2 = a^2$, where $a > 0$.
6. Calculate the Jacobian of the transformation $T(u, v) = (u^2v, uv^2)$.

For Quiz 15 on Thursday, you should be able to do the following:

1. The integral $\int_{-2}^2 \int_0^{\sqrt{4-x^2}} \int_{x^2+y^2}^{\sqrt{8-x^2-y^2}} dz dy dx$ represents the volume of a solid. Sketch the solid, and compute its volume using either cylindrical or spherical coordinates, whichever you prefer.
2. Given the region R in the xy -plane bounded by the hyperbolas $y = 1/x$, $y = 4/x$, and the lines $y = x$, $y = 4x$, find equations for a transformation T that maps a rectangular region S in the uv -plane onto R , where the sides of S are parallel to the u - and v -axes.
3. Evaluate the integral $\iint_R (x+y)e^{x^2-y^2} dA$, where R is the rectangle defined by the inequalities $0 \leq x-y \leq 2$, $0 \leq x+y \leq 3$ by making an appropriate change of variables.
4. Evaluate the integral $\iint_D (6x-4y) dA$, where D is the region bounded by the parallelogram with vertices $(0,0)$, $(2,3)$, $(4,1)$, and $(6,4)$.
5. Use the transformation $x = u^2$, $y = v^2$, $z = w^2$ to express the volume of the solid bounded by the surface $\sqrt{x} + \sqrt{y} + \sqrt{z} = 1$ and the coordinate planes as a triple integral in terms of u , v , and w . (You do not have to evaluate the integral. (Unless you want to.))

Extra fun: Let f be continuous on $[0,1]$ and let R be the triangular region with vertices $(0,0)$, $(1,0)$, and $(0,1)$. Show that

$$\iint_R f(x+y) dA = \int_0^1 u f(u) du.$$

Additional practice problems:

1. Let D be the region in the first quadrant bounded by the curves $y = x$, $y = 2x$, $xy = 1$, and $xy = 2$.
 - (a) Sketch the region, carefully noting all points of intersection of these curves.
 - (b) Determine a change of variables that transforms a rectangle in the (u, v) -plane into the region D .
 - (c) Use this change of variables to evaluate the integral $\iint_D \left(\frac{(x-y)^2}{x^2} - 1 \right) dA$.
2. Repeat the problem above, if D is bounded by the curves $y = x^2$, $y = 2x^2$, $x = y^2$, and $x = 4y^2$, for the integral $\iint_D \left(\frac{x^2}{y^4} + \frac{y^2}{x^4} \right) dA$.
3. Using an appropriate change of variables, show that the area enclosed by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is πab .
4. Evaluate $\iint_D \sqrt{x^2 + y^2} dA$, if D is bounded by the y -axis and the curve $y = \sqrt{4 - x^2}$.
5. Evaluate the following integrals, either by converting to polar coordinates, or by reversing the order of integration:
 - (a) $\int_0^4 \int_{\sqrt{x}}^2 \frac{1}{y^3 + 1} dy dx$.
 - (b) $\int_0^1 \int_y^{\sqrt{2-y^2}} (x + y) dx dy$.
 - (c) $\int_{-1}^1 \int_0^{\sqrt{1-x^2}} \sqrt{1-y^2} dy dx$.