

Math 3500 Assignment #2

University of Lethbridge, Fall 2014

Sean Fitzpatrick

September 12, 2014

Due date: Friday, September 19, by 6 pm.

Please submit solutions to the problems below. For this and the remaining assignments, you are responsible for attempting the exercises in your textbook as practice problems – I won't be providing a list.

Don't forget that you can use Piazza to discuss hints to assigned problems and share solutions to practice problems.

1. Let S and T be nonempty bounded subsets of \mathbb{R} .
 - (a) Prove that if $S \subseteq T$, then $\inf T \leq \inf S \leq \sup S \leq \sup T$.
 - (b) Prove that $\sup S \cup T = \max\{\sup S, \sup T\}$.
(Do not assume that $S \subseteq T$ for part (b).)
2. Let $\mathcal{B}[a, b]$ denote the set of all bounded functions defined on the interval $[a, b]$. (That is, for each $f \in \mathcal{B}[a, b]$, there exist constants $k, l \in \mathbb{R}$ such that $k \leq f(x) \leq l$ for all $x \in [a, b]$.) The *norm* of a function $f \in \mathcal{B}[a, b]$ is defined by

$$\|f\| = \sup\{|f(x)| : x \in [a, b]\}.$$

Prove that $\|f + g\| \leq \|f\| + \|g\|$ for any $f, g \in \mathcal{B}[a, b]$.

Hint: we know that the absolute value function on \mathbb{R} satisfies $|f(x) + g(x)| \leq |f(x)| + |g(x)|$ for all x . The remainder of the proof consists mainly of following the two parts of the definition of a least upper bound to the desired conclusion.

Note: one can check that the set of all bounded functions on an interval has the structure of a vector space, and a norm on a vector space is a generalization of the notion of the length of a vector in \mathbb{R}^n . Now you'll be ready in case anyone ever asks you how long a function is.

3. Prove that if A is any nonempty open subset of \mathbb{R} , then $A \cap \mathbb{Q} \neq \emptyset$.

Hint: use the fact that \mathbb{Q} is dense in \mathbb{R} .

Note: For problems 4-6, you only need to submit **two** of the three problems – you're allowed to skip one.

4. For any set $S \subseteq \mathbb{R}$, let \overline{S} denote the intersection of all the closed sets containing S .
- (a) Prove that \overline{S} is a closed subset of \mathbb{R} .
 - (b) Prove that \overline{S} is the *smallest* closed set containing S . That is, show that $S \subseteq \overline{S}$, and if C is any closed set containing S , then $\overline{S} \subseteq C$.
 - (c) Prove that \overline{S} is equal to the closure of S .
 - (d) Prove that if S is bounded, then \overline{S} is bounded as well.
5. The Nested Intervals Theorem (from the September 10th worksheet, and also mentioned on Piazza) states that if $\{A_n : n \in \mathbb{N}\}$ is a collection of closed bounded intervals (of the form $[a, b]$), and we have $A_{n+1} \subseteq A_n$ for all $n \in \mathbb{N}$, then the intersection $\bigcap A_n$ is nonempty.
- Show that the intervals A_n need to be **both** closed and bounded by giving examples where the theorem fails (that is, where $\bigcap A_n = \emptyset$), if
- (a) The intervals A_n are closed, but not bounded.
 - (b) The intervals A_n are bounded, but not closed.
6. An important theorem regarding compact sets is that if $S \subseteq \mathbb{R}$ is compact, and T is a closed subset of S , then T is compact. Prove this fact using:
- (a) The definition of compactness.
 - (b) The Heine-Borel theorem.