

Solutions to Quiz 12 Practice
Math 2580
Spring 2016


Sean Fitzpatrick

March 1st, 2016

1. For the following double integral, sketch the region of integration, change the order of integration, and evaluate:

$$\int_1^4 \int_1^{\sqrt{x}} (x^2 + y^2) dy dx.$$

Our region of integration is bounded above by $y = \sqrt{x}$ and below by $y = 1$, for $1 \leq x \leq 4$, which gives us the following Type 1 region:



Reversing the order, we have the Type 2 region $y^2 \leq x \leq 4$, for $1 \leq y \leq 2$. Thus, we

have

$$\begin{aligned}
 \int_1^4 \int_1^{\sqrt{x}} (x^2 + y^2) dy dx &= \int_1^2 \int_{y^2}^4 (x^2 + y^2) dx dy \\
 &= \int_1^2 \left(\frac{1}{3}x^3 + xy^2 \right) \Big|_{x=y^2}^{x=4} dy \\
 &= \int_1^2 \left(\frac{64}{3} - \frac{y^6}{3} + 4y^2 - y^4 \right) dy \\
 &= \frac{64y}{3} - \frac{y^7}{21} + \frac{4y^3}{3} - \frac{y^5}{5} \Big|_1^2 \\
 &= \frac{64}{3}(2-1) - \frac{1}{21}(2^7-1) + \frac{4}{3}(2^3-1) - \frac{1}{5}(2^5-1).
 \end{aligned}$$

2. Evaluate the integral $\iiint_B x^2 dV$, where $B = [0, 1] \times [-1, 1] \times [0, 1]$.

We have

$$\iiint_B x^2 dV = \int_0^1 \int_{-1}^1 \int_0^1 x^2 dx dy dz = \int_0^1 \int_{-1}^1 \frac{1}{3} dy dz = \int_0^1 \frac{2}{3} dz = \frac{2}{3}.$$

3. Write the integral $\iiint_W f(x, y, z) dV$, where W is the region between the cone $z = \sqrt{x^2 + y^2}$ and the paraboloid $z = x^2 + y^2$, as an iterated integral. (Start by describing W using inequalities of the form $g_1(x, y) \leq z \leq g_2(x, y)$, where $(x, y) \in D$, and then describe D as either a Type 1 or Type 2 region.)

The cone and parabola intersect when $\sqrt{x^2 + y^2} = x^2 + y^2$. Letting $r = \sqrt{x^2 + y^2}$, this requires $r = r^2$, so $r = 0$ or $r = 1$. (Notice that the region of integration can be obtained by taking the area above the parabola $z = y^2$ and below the line $z = y$, and revolving about the z -axis.) So the two surfaces intersect at the origin, and again along the circle $x^2 + y^2 = 1$ in the plane $z = 1$.

We can thus describe our region via the inequalities $x^2 + y^2 \leq z \leq \sqrt{x^2 + y^2}$, where $x^2 + y^2 \leq 1$. We can describe the disc $x^2 + y^2 \leq 1$ as the Type 1 region $-1 \leq x \leq 1$, $-\sqrt{1-x^2} \leq y \leq \sqrt{1-x^2}$. Thus,

$$\iiint_W f(x, y, z) dV = \int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_{x^2+y^2}^{\sqrt{x^2+y^2}} f(x, y, z) dz dy dx.$$

4. Evaluate the integral $\iiint_W z dV$, where W is the region bounded by the cylinder $x^2 + y^2 = 4$ and the planes $z = 0$ and $z = 1$.

Here, we have $0 \leq z \leq 1$, where $x^2 + y^2 \leq 4$. Carrying out the integral over z , we have

$$\iiint_W z \, dV = \iint_D \left(\int_0^1 z \, dz \right) dA = \iint_D \frac{1}{2} dA = \frac{A(D)}{2},$$

where D is the disc $x^2 + y^2 \leq 4$. We know that this disc has area $A(D) = \pi(2)^2 = 4\pi$, so the result of the integral must be 2π .

5. Describe the surfaces given in cylindrical coordinates by (i) $r = 3$, (ii) $\theta = \pi/4$, and (iii), $z = 2$.

(i) This is the cylinder $x^2 + y^2 = 9$.

(ii) This is the portion of the plane $x = y$ for which $x, y \geq 0$. (It is a vertical plane obtained by extending horizontally from the z -axis along the line $x = y$ for each z -value.)

(iii) This is just the horizontal plane $z = 2$, as usual.

6. Express the surface $z = x^2 + y^2$ in spherical coordinates.

In spherical coordinates we have $x = \rho \cos \theta \sin \varphi$ and $y = \rho \sin \theta \sin \varphi$, so

$$x^2 + y^2 = \rho^2 \cos^2 \theta \sin^2 \varphi + \rho^2 \sin^2 \theta \sin^2 \varphi = \rho^2 \sin^2 \varphi,$$

and $z = \rho \cos \varphi$. Equating the two gives $\rho \cos \varphi = \rho^2 \sin^2 \varphi$, which we can simplify to $\rho = \cot \varphi \csc \varphi$. (Note that this last equation still has $\rho = 0$ as a solution, so we haven't lost the solution $\rho = 0$ by cancelling ρ from both sides.)