

Practice for Quizzes 23 and 24

Math 2580

Spring 2016

Sean Fitzpatrick

April 12th, 2016

For Quiz 23 on Tuesday, you should make sure you can do the problems from Section 2 of the handout on surface integrals. For Quiz 24 on Thursday, you should be able to do the following problems:

1. Use Stokes' theorem to evaluate $\iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$, where $\mathbf{F}(x, y, z) = \langle yz, xz, xy \rangle$, and S is the part of the paraboloid $z = 9 - x^2 - y^2$ that lies above the plane $z = 5$, oriented upward.
2. Evaluate $\iint_S (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$, directly, where $\mathbf{F}(x, y, z) = \langle yz, xz, xy \rangle$, and S is the disk $x^2 + y^2 \leq 4$ in the plane $z = 5$.
3. How are Problems 1 and 2 related?
4. Use Stokes' theorem to evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F}(x, y, z) = \langle xy, 2x, 3y \rangle$, and C is the curve of intersection of the plane $x + z = 5$ and the cylinder $x^2 + y^2 = 9$.
5. Use Stokes' theorem to show that if \mathbf{F} is C^1 vector field defined on all of \mathbb{R}^3 such that $\nabla \times \mathbf{F} = \mathbf{0}$, then \mathbf{F} is conservative.
6. Use the Divergence theorem to evaluate $\iiint_S \mathbf{F} \cdot d\mathbf{S}$ for the vector field $\mathbf{F}(x, y, z) = \langle x^4, -x^3z^2, 4xy^2z \rangle$, where S is the boundary of the region bounded by the cylinder $x^2 + y^2 = 1$ and the planes $z = 0$ and $z = x + 2$.
7. Use the Divergence theorem to evaluate $\iint_S (2x + 2y + z^2) dS$, where S is the sphere $x^2 + y^2 + z^2 = 1$.

Hint: This is the surface integral of a scalar field, but you can re-write it as a the integral of a vector field. (What is the unit normal vector for the given sphere?)