

Math 3410 Assignment #2

University of Lethbridge, Spring 2015

Sean Fitzpatrick

May 21, 2015

Due date: Wednesday, February 11, by 5 pm.

Please provide solutions to the problems below, using the same guidelines as for Assignment #1:

1. Let U be a subspace of a vector space V , and let $S : U \rightarrow W$ be a linear transformation. Prove that the function $T : V \rightarrow W$ given by

$$Tv = \begin{cases} Sv, & \text{if } v \in U \\ 0, & \text{if } v \notin U \end{cases}$$

is **not** a linear transformation.

Hint: if $u \in U$ and $v \notin U$, can $u + v$ be an element of U ? What about $-v$?

Solution: Let $u \in U$ be an element of U such that $Su \neq 0$, and choose some $v \in V$ with $v \notin U$. It follows that $u + v \notin U$, since otherwise we'd have that

$$v = 0 + v = (-u + u) + v = -u + (u + v) \in V,$$

since U is closed under addition. (And if $u \in U$, then we must have $-u \in U$ as well.) Since $u + v \notin U$, on the one hand we have

$$T(u + v) = 0,$$

by definition of T . On the other hand, since $u \in U$ and $v \notin U$, we have

$$T(u) + T(v) = S(u) + 0 = S(u) \neq 0.$$

Thus, T cannot be linear, since $T(u + v) \neq Tu + Tv$.

2. Suppose V is a finite-dimensional vector space, and let $U \subseteq V$ be a subspace. Prove that any linear transformation $S : U \rightarrow W$ can be extended to a linear transformation $T : V \rightarrow W$.

Hint: any basis of U can be extended to a basis for V .

Solution: Let $U \subseteq V$ be a subspace of V , and let $B_U = \{u_1, \dots, u_k\}$ be a basis for U . We know that B_U can be extended to a basis

$$B_V = \{u_1, \dots, u_k, v_1, \dots, v_m\}$$

of V . We know from class (or the textbook) that we can uniquely define a linear transformation by specifying its value on each basis vector. Thus, for example, we can define $T : V \rightarrow W$ by setting $Tu_i = Su_i$ for $i = 1, 2, \dots, k$, and $Tv_i = 0$ for $i = 1, 2, \dots, m$.

Note: We don't have to define T to be zero on all the v_i , but it is a convenient choice. It also provides an opportunity to point out that this is the *correct* way to "extend by zero" from a subspace. Make sure you understand why the method in this question works, but the method in the first problem does not. Basically, we know that the extension by the vectors v_1, \dots, v_k defines a complementary subspace W such that $V = U \oplus W$. We can define T by setting $Tu = Su$ for all $u \in U$, and $Tw = 0$ for all $w \in W$. But this is **not** the same as defining $Tv = 0$ for all $v \notin U$, since if we take nonzero vectors $u \in U$ and $w \in W$, then $v = u + w$ belongs to *neither* U *nor* W .

3. Suppose that V is a finite-dimensional vector space, and $T : V \rightarrow W$ is a linear transformation. Prove that there exists a subspace $U \subseteq V$ such that:

- (a) $U \cap \text{null } T = \{0\}$, and
- (b) $\text{range } T = \{Tu : u \in U\}$.

Solution: We know that $\text{null } T$ is a subspace of V , and since V is finite dimensional, so is $\text{null } T$. Thus, we can choose a basis $B_0 = \{x_1, \dots, x_k\}$ of $\text{null } T$. (Note that $Tx_i = 0$ for $i = 1, 2, \dots, k$.)

Now, since B_0 is a basis for a subspace of V , we can extend it to a basis of V . Let's say that $B_V = \{x_1, \dots, x_k, y_1, \dots, y_n\}$ is the extension. Since B_V is a basis for V , we know that any $v \in V$ can be written in the form

$$v = (a_1x_1 + \dots + a_kx_k) + (b_1y_1 + \dots + b_ny_n).$$

Thus, if we define $U = \text{span}\{y_1, \dots, y_n\}$, it follows that

$$V = U \oplus \text{null } T,$$

and thus $U \cap \text{null } T = \{0\}$, and for any $v \in V$, we can write $v = x + y$, where $x \in \text{null } T$ and $y \in U$. It follows that

$$Tv = T(x + y) = Tx + Ty = Ty,$$

and since v was arbitrary, any element of $\text{range } T$ is of the form Ty , with $y \in U$.

4. Suppose V and W are finite-dimensional vector spaces.

- (a) Prove that there exists an injective (one-to-one) linear transformation $T : V \rightarrow W$ if and only if $\dim V \leq \dim W$.

The Fundamental Theorem of Linear Maps tells us that

$$\dim V = \dim \text{null } T + \dim \text{range } T.$$

If T is one-to-one, then $\dim \text{null } T = 0$, so $\dim V = \dim \text{range } T \leq \dim W$, since $\text{range } T$ is a subspace of W . Conversely, suppose $\dim V \leq \dim W$, and choose bases $\{v_1, \dots, v_k\}$ of V and $\{w_1, \dots, w_n\}$ of W , with $k \leq n$. Let us define $T : V \rightarrow W$ by setting

$$Tv_1 = w_1, Tv_2 = w_2, \dots, Tv_k = w_k,$$

and extending by linearity. (As noted above, T is completely determined by its values on a basis of V .) Then T is an injection, as follows: suppose $v \in \text{null } T$, and write $v = c_1v_1 + \dots + c_kv_k$ in terms of the given basis. Then

$$0 = Tv = T(c_1v_1 + \dots + c_kv_k) = c_1Tv_1 + \dots + c_kTv_k = c_1w_1 + \dots + c_kw_k,$$

and the vectors w_1, \dots, w_k are independent, since they're part of a basis. It follows that $c_1 = 0, \dots, c_k = 0$, and thus $v = 0$, which implies that $\text{null } T = \{0\}$ and that T is an injection.

- (b) Prove that there exists a surjective (onto) linear transformation $T : V \rightarrow W$ if and only if $\dim V \geq \dim W$.

If T is onto, then $\text{range } T = W$, so

$$\dim V = \dim \text{null } T + \dim \text{range } T = \dim \text{null } T + \dim W \geq \dim W.$$

Conversely, suppose $\dim V \geq \dim W$, and choose bases $\{v_1, \dots, v_k\}$ and $\{w_1, \dots, w_n\}$ of V and W respectively, with $k \geq n$. As above, we can define $T : V \rightarrow W$ in terms of the basis for V . We set

$$Tv_1 = w_1, Tv_2 = w_2, \dots, Tv_n = w_n, \text{ and } Tv_j = 0 \text{ for } n+1 \leq j \leq k.$$

It follows that T is a surjection, since given any

$$w = c_1w_1 + \dots + c_nw_n \in W,$$

we can let $v = c_1v_1 + \dots + c_nv_n$, and then $Tv = w$, as required.