

Measuring the forces between DNAs in solution

Xiangyun Qiu*, Lisa Kwok, Hye Yoon Park, Jessica S. Lamb, Kurt Andresen, Lois Pollack

School of Applied and Engineering Physics, Cornell University, Ithaca, New York 14853, USA

Contact: xiangyun@mail.nih.gov

Quantifying the forces between DNAs is essential to improve our understanding of many fundamental biological processes. Using the small angle X-ray scattering (SAXS) technique, we have probed the effective forces between isolated DNAs (25 base pair, 48 bare charge) in solution by measuring the inter-DNA interference, from which the inter-DNA pair potentials are obtained following the generalized one-component method (GOCM). We report the measured effective interaction charges under varying DNA concentrations, counterion concentrations and valences. Notably, we observed the onset of an intriguing attractive force at rather low divalent Mg^{2+} concentrations $>16\text{mM}$, which was successfully modelled as a short range attraction. Our quantitative measurement of the effective inter-DNA forces should provide valuable comparisons to future theoretical studies.

- Present address: National Institutes of Health, Bethesda, MD, 20892

References

1. X. Qiu, K. Andresen, L. W. Kwok, J. S. Lamb, H.-Y. Park, and L. Pollack, *Inter-DNA Attraction Mediated by Divalent Counterions*, Physical Review Letters, 99, 038104, (2007).
2. X. Qiu L. W. Kwok, H.-Y. Park, J. S. Lamb, K. Andreson, and L. Pollack. *Measuring Inter-DNA Potentials in Solution* , Physical Review Letters, 96, 138101, (2006)