

a
Precision–recall (ITT; descriptive export)
zavg_BEF (AP 0.333)
B (AP 0.310)
GLS (AP 0.212)
Spearman (AP 0.204)
Pearson (AP 0.197)
PC-rm Pearson (AP 0.239)
prevalence 0.0055
Precision
Recall
b
Paired differences in AP
zavg_BEF − GLS
zavg_BEF − Spearman
zavg_BEF − Pearson
B − GLS
B − Spearman
B − Pearson
B − zavg_BEF
PC-rm − zavg_BEF
ITT (primary metric)
S1 (locked verdict set)
Δ AP (95% cluster-bootstrap CI)
c
Calibration (ITT, log-spaced bins)
zavg_BEF: Brier skill 0.222
calibration-in-the-large 0.893
slope 1.007
zavg_BEF
GLS
Spearman
Pearson
Observed co-complex rate
Mean predicted probability (isotonic, fitted on train)
d
Pooled vs cluster-structured AP (ITT)
zavg_BEF
GLS
Spearman
Pearson
zavg_BEF − GLS, macro-AP:
+0.005 [-0.014, +0.047]
AP (ITT)
AP (pooled)
AP without largest comp.
AP without 10 largest
macro per-cluster AP
e
Generalisation to proteins not seen among train positives
zavg_BEF
B
GLS
Spearman
Pearson
AP (95% cluster-bootstrap CI)
C1: both proteins in train positives
positives = 265
Δ AP vs GLS:
+0.081 [+0.026, +0.140]
C2: one protein
positives = 738
Δ AP vs GLS:
+0.050 [+0.017, +0.092]
C3: neither protein
positives = 2,309
Δ AP vs GLS:
+0.148 [+0.093, +0.199]
all (ITT)
positives = 3,312
Δ AP vs GLS:
+0.121 [+0.079, +0.159]
f
Macro per-cluster AP by positive pairs per test cluster (ITT; descriptive, non-confirmatory)
zavg_BEF
GLS
Macro per-cluster AP
(within-bin resample CI)
1 positive pair
91 clusters, 91 pos.
Δ macro AP vs GLS:
-0.033 [-0.090, +0.023]
2-5 positive pairs
67 clusters, 213 pos.
Δ macro AP vs GLS:
-0.024 [-0.061, +0.014]
6-20 positive pairs
43 clusters, 413 pos.
Δ macro AP vs GLS:
+0.041 [+0.008, +0.080]
21-100 positive pairs
25 clusters, 1,141 pos.
Δ macro AP vs GLS:
+0.126 [+0.075, +0.182]
101+ positive pairs
6 clusters, 1,454 pos.
Δ macro AP vs GLS:
+0.142 [+0.078, +0.198]