
a
Map overview
zavg_BEF: 367 complexes, 3,642 genes
GLS: 379 complexes, 2,862 genes
Clusters
Level-1 cluster size (genes)
3-4
5-8
9-16
17-60
b
Coverage
Fraction of S genes clustered (S = 6,022 genes)
594
410
1,722
1,494
1,326
958
pan-essential
selective
other
c
Selection
selected z0 = 6, k = 10
Split-half recovery statistic (mean ± SD over 4 splits)
zavg_BEF, k = 10
zavg_BEF, k = 20
GLS (edge-matched), k = 10
GLS (edge-matched), k = 20
rewired baseline = 0
Edge threshold z0
d
Tiers
zavg_BEF L1 (n = 367)
zavg_BEF L1 non-locus (n = 263)
zavg_BEF L2 (n = 132)
GLS L1 (n = 379)
GLS L1 non-locus (n = 295)
GLS L2 (n = 108)
159
74
124
155
60
47
59
28
42
159
81
131
151
69
69
42
24
38
Fraction of clusters
T1
T2
T3-neg
T3-nc
e
Variability
Median best-match Jaccard in 20 bootstrap maps
Complex size bin
pairwise ARI (median, 190 map pairs):
zavg_BEF 0.754, GLS 0.747
f
Bridge endpoint: no difference claimed (the 95% CI of the primary endpoint includes 0)
PRIMARY: BioPlex precision
no coverage matching
locus clusters excluded
secondary: co-abundance r
zavg 0.157 vs GLS 0.182; Δ -0.025 [-0.075, +0.031]
zavg 0.164 vs GLS 0.184; Δ -0.020 [-0.054, +0.017]
zavg 0.212 vs GLS 0.208; Δ +0.003 [-0.060, +0.074]
zavg 0.211 vs GLS 0.272; Δ -0.061 [-0.101, -0.018]
pair level, S × S (13,868,011 pairs): AP vs BioPlex zavg_BEF 0.0345, GLS 0.0215, Pearson 0.0223 (prevalence 0.0026)
Difference zavg_BEF map − GLS map (95% protein-node bootstrap CI; ← favours GLS | favours zavg_BEF →)