








a**T** $p = 5.0e-04$ **b****N** $p = 3.8e-04$ **c****stage** $p < 1e-4$ 

PHLDA3 copy number vs expression
Spearman rho=0.21, Kruskal P=1.8e-05






PHLDA3 co-expression functional enrichment (TCGA-THCA tumors, n=502)


a

GO Biological Process




b

KEGG Pathway




c

GSEA Hallmark



AGO2-CLIP-supported miRNAs targeting PHLDA3
(blue = inverse → candidate repressors)




PHLDA3 candidate ceRNA network (exploratory)
lncRNA — miRNA — PHLDA3
(TCGA-THCA + ENCORI AGO2-CLIP)

lncRNA (ceRNA)
+corr with PHLDA3


miRNA
(AGO2-CLIP, repressor)

target




Compounds reversing the PHLDA3-high signature (L1000CDS2)


red = HDAC inhibitor · teal = HSP90 inhibitor



PHLDA3-high cell lines are sensitive to MEK/ERK inhibitors
 (left = more sensitive · teal = MAPK-pathway inhibitor; FDR<0.05)




clue.io CMap (MOA-class): HDAC/HSP90 inhibitors reverse the
PHLDA3-high program (red), consistent with L1000CDS2




PHLDA3 protein (HPA IHC, antibody HPA017385)

Normal thyroid gland



Thyroid cancer (papillary)



a

Nomogram for predicting lymph-node metastasis (N1)

