



İTÜ Computer Engineering Dep.
 Assoc. Prof. Feza BUZLUCA
 Asst. Prof. Sanem KABADAYI
 Prof. Mustafa KAMAŞAK
 November 24, 2023

BLG 231E DIGITAL CIRCUITS MIDTERM SOLUTIONS

QUESTION 1 (30 Points):

Note that Parts (a) and (b) below are not related.

a) [15 points]

i. After an addition operation, overflow can occur in the following two cases:

1. negative + negative = positive
2. positive + positive = negative

Based on the given information, only the second case is possible.

$r_3: 1$	$a_3: 0$	$b_3: 0$	Case 1: both -, result + $1\ 1\ a_1\ a_0$ $1\ 1\ b_1\ b_0$ $0\ r_3\ r_1\ r_0$ <u>not possible</u>	Case 2: both +, result - $0\ 0\ a_1\ a_0$ $0\ 0\ b_1\ b_0$ $1\ r_3\ r_1\ r_0$ possible
----------	----------	----------	--	---

ii.

Answer: No

Reason: Overflow occurring in their addition means they have the same sign, so overflow cannot occur in their subtraction.

b) [15 Points]

i.

$$29/2 = 2 \times 14 + 1 \text{ (least significant bit)}$$

$$14/2 = 2 \times 7 + 0$$

$$7/2 = 2 \times 3 + 1$$

$$3/2 = 2 \times 1 + 1$$

$$1/2 = 2 \times 0 + 1$$

$$\begin{array}{rcl}
 29(\text{decimal}) & = & 11101 \text{ (5-bit binary)} \\
 & = & 0001\ 1101 \text{ (8-bit binary)} \\
 1\text{'s complement} & = & 1110\ 0010 \\
 +1 & & 0000\ 0001 \\
 2\text{'s complement} & = & \underline{1110\ 0011} = -29
 \end{array}$$

ii.

$$Y = 1110\ 0011$$

$$X = \$9A = 1001\ 1010$$

$$X' = 0110\ 0101$$

$$X'+1 = 0110\ 0110$$

$$Y = 1110\ 0011$$

$$X'+1 = \begin{array}{r} +\ 0110\ 0110 \\ \hline 10100\ 1001 \end{array}$$

$$10100\ 1001$$

ignore 9th bit

Since negative – negative = positive, there is **no overflow**.

Since the result is **positive** and there is **no overflow**, **Y > X**.

Notes:

1. Only checking the sign of the result to compare signed integers is not sufficient. You must also **consider the overflow**. The sign is valid only if there is **no overflow**.

2. As the integers provided are **signed** numbers, comparison between them using **borrow is not possible**.

3. \$9A is **not** 154 (decimal). You don't need to convert a hexadecimal number to decimal before converting to binary. The hexadecimal representation is only used for documentation (to write and read easily). Converting hexadecimal to binary, and vice versa, can be done easily. (Slide 1.21)

QUESTION 2 (35 Points):

Note that Parts (a), (b), and (c) below are not related.

a) [15 points]

$$(a+f(0,b))(\bar{a}+f(1,b))$$

$$\begin{aligned} &= a\bar{a} + af(1,b) + \bar{a}f(0,b) + f(0,b)f(1,b) && \text{(multiply out (distributive law))} \\ &= 0 + af(1,b) + \bar{a}f(0,b) + f(0,b)f(1,b) && \text{(since } a\bar{a}=0 \text{ (inverse law))} \\ &= af(1,b) + \bar{a}f(0,b) && \text{(consensus theorem)} \end{aligned}$$

Note: $f(0,b)f(1,b)$ is the consensus term of $af(1,b) + \bar{a}f(0,b)$ with respect to a .

b) [10 points]

Yes.




- NOT: $f(0,x) = 0 + \bar{x} = \bar{x}$.
- OR : $f(x,f(0,y)) = f(x,\bar{y}) = x + \overline{(\bar{y})} = x + y$.

Since {OR, NOT} is a complete set, AND can also be implemented. Specifically,

- AND: $xy = (\bar{x} + \bar{y})' = f(0, f(f(0,x), y))$
 $= f(0, f(\bar{x}, y))$
 $= f(0, \bar{x} + y)$
 $= (\bar{x} + y)'$

c) [10 points]

$$\begin{aligned} r = f(w,x,y,z) &= w\bar{x}y + xyz \\ &= (w\bar{x} + xz)y \end{aligned}$$



Six NAND gates are sufficient.

QUESTION 3 (35 Points):

Note that Parts (a), (b), and (c) below are not related.

a) [10 points]

abc		
(0)	000	✓
(1)	001	✓
(2)	010	✓
(4)	100	✓
(6)	110	✓

abc		
(0,1)	00-	
(0,2)	0-0	✓
(0,4)	-00	✓
(2,6)	-10	✓
(4,6)	1-0	✓

abc		
(0,2,4,6)	--0	

The set of all prime implicants = $\{\bar{c}, \bar{a}\bar{b}\}$

b) [10 points]

	$\bar{C}\bar{D}$	$\bar{C}D$	$C\bar{D}$	CD
$\bar{A}\bar{B}$	1	0	1	1
$\bar{A}B$	1	0	x	1
$A\bar{B}$	0	0	0	x
AB	1	1	0	1

<input type="checkbox"/>	$\bar{A}C$
<input type="checkbox"/>	$\bar{A}\bar{D}$
<input type="checkbox"/>	$\bar{B}\bar{D}$
<input type="checkbox"/>	$C\bar{D}$

☐ $A\bar{B}\bar{C}$

When we are finding the set of all prime implicants of a function in SOP form, we assign 1 to Φ 's.

The set of all prime implicants = $\{\bar{a}c, \bar{a}\bar{d}, \bar{b}\bar{d}, c\bar{d}, a\bar{b}\bar{c}\}$

c) [15 points]

	i	j	k	l	Cost
A		X		X	4
B	X	X			3
C	X	X	X		5
D		X	X		2
E			X	X	2

Since column j covers i, we remove column j.

	i	k	l	Cost
A			X	4
B	X			3
C	X	X		5
D		X		2
E		X	X	2

Prime implicant E covers more points than A, and its cost is lower. Prime implicant E covers more points than D, and their cost is the same. Therefore, we remove rows A and D.

	i	k	l	Cost
B	X			3
C	X	X		5
E		X	X	2

To cover l, we must take the prime implicant E. We mark E and remove it from the table.

	i	Cost
B	X	3
C	X	5

B is cheaper than C. To cover i, we take the prime implicant B. We mark B. We covered all the true points of the function.

The minimal covering sum: $f = B + E$

Note: Since the given prime implicant chart is simple, the solution is obvious (readily apparent). However, you are expected to simplify the chart in a **systematic way** and provide **all the steps** of the process.