

## VV570

### Numerical Analysis

#### Assignment 3

Jing and Manuel — UM-JI

#### Reminders

- Write in a neat and legible handwriting or use L<sup>A</sup>T<sub>E</sub>X
- Clearly explain the reasoning process
- Write in a complete style (subject, verb, and object)
- Be critical on your results

*Questions preceded by a \* are optional. Although they can be skipped without any deduction, it is important to know and understand the results they contain.*

#### Ex. 1 — Cantor's set

In this exercise we investigate a few properties of Cantor's set.

Calling  $C_0$  the compact  $[0, 1]$  we trisect it and remove the middle open part and define  $C_1$  as  $[0, 1/3] \cup [2/3, 1]$ . Recursively repeating the process we construct an infinite sequence  $C_0 \supset C_1 \supset \dots \supset C_i \supset \dots$ . We define the *Cantor set* as the intersection of all the nested  $C_i$ ,  $C = \bigcap_{i=0}^{\infty} C_i$ .

- \* 1. Prove that  $C$  is compact.
- 2. Show that for any two elements  $x < y$  of  $C$ , there exists  $z \notin C$  such that  $x < z < y$ .
- 3. We estimate how "large" the set  $C$  is with respect to  $\lambda$ , the Lebesgue measure.
  - a) What is the Lebesgue measure of a countable set?
  - b) Show that for  $n \in \mathbb{N}$ ,  $\lambda(C_n) = \left(\frac{2}{3}\right)^n$ , and conclude that  $C$  has Lebesgue measure 0.
- 4. We estimate how "large" the set  $C$  is with respect to cardinality.
  - a) Show that  $C$  is not empty.
  - \* b) We write any  $x \in [0, 1]$  using the ternary expansion

$$x = \sum_{i=1}^{\infty} \frac{a_i}{3^i}, \quad \text{with } a_i \in \{0, 1, 2\}.$$

Describe the form of the  $x$  belonging to  $C_i$ ,  $i \in \mathbb{N}$ .

- c) Clearly explain Cantor diagonal argument.
- \* d) Using Cantor diagonal argument show that Cantor's set is uncountable.
- 5. From questions 3, Cantor's set has Lebesgue measure 0, and from question 4 it is uncountable. Explain how this is possible.

#### Ex. 2 — Cantor's function

We now define the Cantor function following the construction process of the Cantor set. Let  $f_1$  be 1/2 over the "removed interval"  $(1/3, 2/3)$  and linear on  $C_1$ . Then define  $f_2$  to be 1/4 and 3/4 on the two removed intervals, to coincide with  $f_1$  in  $1/3$  and  $2/3$ , while being linear on the remaining four intervals. The process is carried on as the Cantor set is built, defining the Cantor function  $f_C$ .

- 1. Show that the  $(f_n)_{n \in \mathbb{N}}$  define a sequence of monotonically increasing continuous functions.
- 2. Show that  $(f_n)_{n \in \mathbb{N}}$  converges uniformly to  $f_C : [0, 1] \rightarrow [0, 1]$ .
- 3. Prove that Cantor function is
  - a) Uniformly continuous;

*Hint: prove (or assume) that a continuous function on a compact is uniformly continuous.*

- b) Monotonically increasing;
- \* c) Differentiable almost everywhere, with  $f'(x) = 0$ ;

4. Prove that  $f_C$  is not absolutely continuous.

*Note:* the Cantor function is one of the most simple function to be uniformly continuous but not absolutely continuous

**Ex. 3 — Taylor's theorem**

Reasoning by induction and applying the fundamental theorem of calculus, prove

1. Taylor's theorem as given in the slides (theorem 2.36).
2. Taylor's theorem with the remainder in the Lagrange form; Calling  $P_n(x)$  the polynomial part of  $f(x)$  in Taylor's theorem, show the existence of  $c \in [a, x]$  such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x-a)^{n+1}.$$

**Ex. 4 — Convergence of rationals to irrationals**

Intuitively a *complete space* has “no point missing” anywhere. In particular it means that any Cauchy sequence converges inside the space. In this exercise we show that  $e$  is not rational while we find a Cauchy sequence of rationals converging to  $e$ .

1. Show that  $e$  is irrational.
2. Show that the sequence  $(u_n)_{n \in \mathbb{N}}$  defined by  $u_n = \left(1 + \frac{1}{n}\right)^n$  is a Cauchy sequence converging to  $e$ .
3. Is  $\mathbb{Q}$  complete? Explain.