

Introduction to Numerical Analysis

HW9

Yu Cang
018370210001

July 30, 2018

1 QUESTION 1

- (a) *Proof.* As \mathcal{A} is convex, the mean value theorem can be applied in terms of y s.t.

$$\Phi(t, y_2) - \Phi(t, y_1) = (y_2 - y_1) \frac{\partial \Phi(t, y)}{\partial y} \Big|_{y=\xi} \quad (1.1)$$

where $\xi \in (y_1, y_2)$.

Since there exists $c > 0$ s.t. for all $(t, y) \in \mathcal{A}$

$$\left| \frac{\partial \Phi(t, y)}{\partial y} \right| \leq c \quad (1.2)$$

Thus

$$\begin{aligned} & |\Phi(t, y_2) - \Phi(t, y_1)| \\ &= |y_2 - y_1| \left| \frac{\partial \Phi(t, y)}{\partial y} \Big|_{y=\xi} \right| \\ &\leq c |y_2 - y_1| \end{aligned} \quad (1.3)$$

which implies that $\Phi(t, y)$ satisfies Lipschitz condition in y on \mathcal{A} . \square

- (b) *Proof.* Let $P_1 = (t_1, y_1)$ and $P_2 = (t_2, y_2)$. Then any point P' lies on the line segment joining P_1 and P_2 can be expressed as

$$\begin{aligned} P &= (1 - \alpha)P_1 + \alpha P_2 \\ &= ((1 - \alpha)t_1 + \alpha t_2, (1 - \alpha)y_1 + \alpha y_2) \\ &\triangleq (t', y') \end{aligned} \quad (1.4)$$

where $\alpha \in [0, 1]$.

It's clear that t_1 and t_2 lies between t_0 and T , and t' lies between t_1 and t_2 . Thus, t' also lies between t_0 and T .

Further, it's also clear that $-\infty < y' < +\infty$.

Hence $P \in \mathcal{D}$, which implies that \mathcal{D} is convex. \square

(c) *Proof.* Let

$$\Phi(t, y) = \frac{4t^3 y}{1 + t^4} \quad (1.5)$$

Then

$$\frac{\partial \Phi(t, y)}{\partial y} = \frac{4t^3}{1 + t^4} = \frac{4t}{t^2 + \frac{1}{t^2}} < \frac{4}{t^2 + \frac{1}{t^2}} < \frac{4}{2\sqrt{t^2 \frac{1}{t^2}}} = 2 \quad (1.6)$$

as $t \in (0, 1)$.

which implies that $\Phi(t, y)$ satisfies a Lipschitz condition in y .

Thus, the given IVP problem has a unique solution. \square

(d) Definitely not recommended.

As $\Phi(t, y) = 1 + y^2$, then

$$\frac{\partial \Phi}{\partial y} = 2y \triangleq \lambda y \quad (1.7)$$

Here $\lambda > 0$, and the Euler's method is not stable as the error will be amplified at each iteration step. Finally the calculation will diverge.

2 QUESTION 2

(a)

3 QUESTION 4

4 QUESTION 6