

Advanced Convection

HW1

Yu Cang
018370210001

June 29, 2018

1 PROBLEM 1

When $u_2 = 0$ and the flow is entirely in the x-direction, the governing equations can be simplified as follows

$$u_1 \frac{\partial u_1}{\partial x_1} + \frac{1}{\rho} \frac{\partial p}{\partial x_1} = 0 \quad (1.1)$$

$$\frac{\partial p}{\partial x_2} = 0 \quad (1.2)$$

where the body-force are neglected.

It can be concluded that pressure doesn't change in y-direction when x is fixed.

Also, pressure gradient in x-direction is the driven force of the fluids.

2 PROBLEM 2

When it comes to constant property

$$\frac{\partial}{\partial y} \left(k \frac{\partial T}{\partial y} \right) = k \frac{\partial^2 T}{\partial y^2} \quad (2.1)$$

Since $i = C_p T$ when considering ideal gas

$$\rho u \frac{\partial i}{\partial x} + \rho v \frac{\partial i}{\partial y} = \rho C_p \left(u \frac{\partial T}{\partial x} + v \frac{\partial T}{\partial y} \right) \quad (2.2)$$

As $(\frac{\partial u}{\partial y})^2$ is a small term compared to other terms, it can be seen as 0.
(Sorry, haven't figured out how to eliminate $\frac{\partial p}{\partial x}...$)

3 PROBLEM 3