

Problem 8

Problem 1. $\forall L1, L2 \in \text{NatList}, \text{diff}(L1, L2) = \text{diff}(L1, \text{rev}(L2))$

Proof. By structural induction on $L1$.

(1) Base case

What to show: $\text{diff}(\text{nil}, l2) = \text{diff}(\text{nil}, \text{rev}(l2))$.
 where $l2 \in \text{NatList}$. Note that $l2$ is a fresh constant¹.

$$\begin{array}{ll} \text{diff}(\text{nil}, l2) \longrightarrow \text{nil} & \text{(by diff1)} \\ \underline{\text{diff}(\text{nil}, \text{rev}(l2))} \longrightarrow \text{nil} & \text{(by diff1)} \end{array}$$

(2) Induction case

What to show: $\text{diff}(x \mid l1, l2) = \text{diff}(x \mid l1, \text{rev}(l2))$
 Induction hypothesis: $\text{diff}(l1, l2) = \text{diff}(l1, \text{rev}(l2))$
 where $x \in \text{PNat}$ and $l1, l2 \in \text{NatList}$. Note that $x, l1, l2$ are fresh constants.

$$\begin{array}{l} \underline{\text{diff}(x \mid l1, l2)} \longrightarrow \text{if has}(l2, x) \text{ then } \underline{\text{diff}(l1, l2)} \text{ else } (x \mid \underline{\text{diff}(l1, l2)}) \text{ fi} \\ \hspace{15em} \text{(by diff2)} \\ \hspace{10em} \longrightarrow \text{if has}(l2, x) \text{ then } \text{diff}(l1, \text{rev}(l2)) \text{ else } (x \mid \text{diff}(l1, \text{rev}(l2))) \text{ fi} \\ \hspace{15em} \text{(by IH)} \\ \underline{\text{diff}(\text{nil}, \text{rev}(l2))} \longrightarrow \text{if } \underline{\text{has}(\text{rev}(l2), x)} \text{ then } \text{diff}(l1, \text{rev}(l2)) \text{ else } (x \mid \text{diff}(l1, \text{rev}(l2))) \text{ fi} \\ \hspace{15em} \text{(by diff2)} \\ \hspace{10em} \longrightarrow \text{if has}(l2, x) \text{ then } \text{diff}(l1, \text{rev}(l2)) \text{ else } (x \mid \text{diff}(l1, \text{rev}(l2))) \text{ fi} \\ \hspace{15em} \text{(by Problem 6)} \end{array}$$

□

¹A fresh constant of a sort denotes an arbitrary value of the sort, and has never been used before.