

## Problem 12

**Problem 1.**  $\forall L \in \text{NatList}, \text{sum}(L) = \text{sum}(\text{rev}(L))$ .

*Proof.* By structural induction on  $L$ .

**(1) Base case**

What to show:  $\text{sum}(\text{nil}) = \text{sum}(\text{rev}(\text{nil}))$ .

$$\begin{aligned} \frac{}{\text{sum}(\text{nil})} &\longrightarrow 0 && \text{(by sum1)} \\ \text{sum}(\text{rev}(\text{nil})) &\longrightarrow \frac{}{\text{sum}(\text{nil})} && \text{(by rev1)} \\ &\longrightarrow 0 && \text{(by sum1)} \end{aligned}$$

**(2) Induction case**

What to show:  $\text{sum}(x \mid l) = \text{sum}(\text{rev}(x \mid l))$

Induction hypothesis:  $\text{sum}(l) = \text{sum}(\text{rev}(l))$

where  $x \in \text{PNat}$  and  $l \in \text{NatList}$ . Note that  $x, l$  are fresh constants<sup>1</sup>.

$$\begin{aligned} \frac{}{\text{sum}(x \mid l)} &\longrightarrow x + \text{sum}(l) && \text{(by sum2)} \\ &\longrightarrow x + \text{sum}(\text{rev}(l)) && \text{(by IH)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + x && \text{(by comm+)} \\ \text{sum}(\text{rev}(x \mid l)) &\longrightarrow \text{sum}(\text{rev}(l) @ (x \mid \text{nil})) && \text{(by rev2)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + \text{sum}(x \mid \text{nil}) && \text{(by Lemma 1)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + (x + \text{sum}(\text{nil})) && \text{(by sum2)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + (x + 0) && \text{(by sum1)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + (0 + x) && \text{(by comm+)} \\ &\longrightarrow \text{sum}(\text{rev}(l)) + x && \text{(by +1)} \end{aligned}$$

□

**Lemma 1.**  $\forall L1, L2 \in \text{NatList}, \text{sum}(L1 @ L2) = \text{sum}(L1) + \text{sum}(L2)$ .

<sup>1</sup>A fresh constant of a sort denotes an arbitrary value of the sort, and has never been used before.

*Proof.* By structural induction on  $L1$ .

**(1) Base case**

What to show:  $\text{sum}(\text{nil} @ l2) = \text{sum}(\text{nil}) + \text{sum}(l2)$   
 where  $l2 \in \text{NatList}$ . Note that  $l2$  is a fresh constant.

$$\begin{aligned} \text{sum}(\text{nil} @ l2) &\longrightarrow \text{sum}(l2) && \text{(by @1)} \\ \underline{\text{sum}(\text{nil})} + \text{sum}(l2) &\longrightarrow \underline{0 + \text{sum}(l2)} && \text{(by sum1)} \\ &\longrightarrow 0 && \text{(by +1)} \end{aligned}$$

**(2) Induction case**

What to show:  $\text{sum}((x | l1) @ l2) = \text{sum}(x | l1) + \text{sum}(l2)$

Induction hypothesis:  $\text{sum}(l1 @ l2) = \text{sum}(l1) + \text{sum}(l2)$

where  $x \in \text{PNat}$  and  $l1, l2 \in \text{NatList}$ . Note that  $x, l1, l2$  are fresh constants.

$$\begin{aligned} \text{sum}(\underline{(x | l)}) @ l2 &\longrightarrow \underline{\text{sum}(x | (l1 @ l2))} && \text{(by @2)} \\ &\longrightarrow x + \underline{\text{sum}(l1 @ l2)} && \text{(by sum2)} \\ &\longrightarrow x + (\underline{\text{sum}(l1) + \text{sum}(l2)}) && \text{(by IH)} \\ &\longrightarrow (x + \text{sum}(l1)) + \text{sum}(l2) && \text{(by assoc+)} \\ \underline{\text{sum}(x | l1)} + \text{sum}(l2) &\longrightarrow (x + \text{sum}(l1)) + \text{sum}(l2) && \text{(by sum2)} \end{aligned}$$

□