



Day 0 — HST 0 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
0

1,000 L TANK

**plain water — feed
starts tomorrow**

EC · pH

— · **5,8–6,3**

SOP / TANK

—

TIMER · 0.5 L

**3 × 5 min · 8 am · 12 pm · 4
pm**

1 Transplant + Trichoderma

Plant seedlings gently into the buffered bags. **Plain water only today**, no feed. Water the bags first, then **one cup (250 ml)** of Trichoderma at the stem base: 10 g per 10 L bucket, stand 15–30 min, use the same day. **Never** through the drippers or tank.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves
Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 1 — HST 1 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
1

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4 pm

1 Feed starts today

Stock A + Stock B S1 → 1,000 L tank · EC 2.1 · pH 5.8–6.3. No SOP.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 2 — HST 2 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
2

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4 pm **No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 3 — HST 3 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
3

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4
pm

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 4 — HST 4 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
4

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4 pm

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 5 — HST 5 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
5

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4 pm

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 6 — HST 6 jobs

S1 · Grow — establish · Sparmanik 4.0

HST
6

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4 pm**No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.5 L

3 × 5 min · 8 am · 12 pm · 4
pm**1 Start removing side shoots**

Every 2–3 days until node 8. Pinch the **shoot** in the leaf armpit on nodes 1–8, **keep the leaf**. Never the main tip. Leave nodes 9–12 — fruit comes from there.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 75–100 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Seedlings wilting at noon · rotting at the base (damping-off) · yellow leaves

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 8 — HST 8 jobs

S1 · Grow — vegetative · Sparmanik 4.0

HST
8

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5 pm

1 Silicon drench (Kalsika)

MANUAL switch. 400 L clean water → run 3 min → one 100 g sachet into ~300 L, stir until clear → run 8 min → top up to 100 L only → run 3 min until empty → make the normal feed tank. Today only the **2 pm and 5 pm** pulses.

2 Change the timer

From tomorrow (day 9): **4 × 6 min · 8 · 11 am · 2 · 5 pm**. Set it once and leave it until day 25.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 9 — HST 9 jobs

S1 · Grow — vegetative · Sparmanik 4.0

HST
9

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm**

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 11 — HST 11 jobs

S1 · Grow — vegetative · Sparmanik 4.0

HST
11

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm** **No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****1 Silicon drench (Kalsika)**

MANUAL switch. 400 L clean water → run 3 min → one 100 g sachet into ~300 L, stir until clear → run 8 min → top up to 100 L only → run 3 min until empty → make the normal feed tank. Today only the **2 pm and 5 pm** pulses.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****1 Trichoderma cup drench**

10 g per 10 L bucket of plain water, stand 15–30 min. Water the bags first, then one cup (250 ml) at the stem base. Morning or late afternoon. Stir every 10 bags. 2 days away from silicon.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 18 — HST 18 jobs

S1 · Grow — vegetative · Sparmanik 4.0

HST
18

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm**

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 20 — HST 20 jobs

S1 · Grow — vegetative · Sparmanik 4.0

HST
20

1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm**

1 Node 8 — stop removing shoots

From node 9 up, leave the shoots. Female flowers and fruit grow from here.

2 Tri-Pholate spray after 4 pm

2 g per 16 L spray pack (0.125 g/L). Leaves only. After 4 pm.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****1 Silicon drench (Kalsika)**

MANUAL switch. 400 L clean water → run 3 min → one 100 g sachet into ~300 L, stir until clear → run 8 min → top up to 100 L only → run 3 min until empty → make the normal feed tank. Today only the **2 pm and 5 pm** pulses.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm** **No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S1 stock

EC · pH

2.1 · 5,8–6,3

SOP / TANK

no SOP

TIMER · 0.8 L

**4 × 6 min · 8 · 11 am · 2 · 5
pm****1 Tonight: make the FLOWER stock**

Boyd's kit (with Boroni). See book pages 23 / 53. Never boric acid.

Every day

- 06:30 tank check: EC 2.1 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below $\rightarrow$ mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 120–160 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Aphids · thrips · whitefly · spider mites · caterpillars · first mildew spots · pale new leaves (run-off pH)

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 CHANGE TO FLOWER

Tank: FLOWER stock + **SOP 310 g** → EC 2.4. Timer **5 × 7 min**. Start hand pollination every morning. Hang fruit-fly traps.

2 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Tri-Pholate spray after 4 pm**2 g per 16 L spray pack** (0.125 g/L). Leaves only. After 4 pm.**2 Hand pollination 06:00–10:00**

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Silicon drench (Kalsika)

MANUAL switch. 400 L clean water → run 3 min → one 100 g sachet into ~300 L, stir until clear → run 8 min → top up to 100 L only → run 3 min until empty → make the normal feed tank. Today only the **2 pm and 5 pm** pulses.

2 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S2 stockEC · pH
2.4 · 5,8–6,3SOP / TANK
310 gTIMER · 1.2 L
5 × 7 min · 8 · 10 · 12 · 2 · 4**1 Hand pollination 06:00–10:00**

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S2 stockEC · pH
2.4 · 5,8–6,3SOP / TANK
310 gTIMER · 1.2 L
5 × 7 min · 8 · 10 · 12 · 2 · 4**1 Tri-Pholate spray after 4 pm — LAST ONE****2 g per 16 L spray pack** (0.125 g/L). Leaves only. After 4 pm. Stop after HST 35 — spray marks the net.**2 Hand pollination 06:00–10:00**Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.**3 Calcinit leaf spray****80 g per 16 L spray pack**. Early morning or after 4 pm, never in strong sun. Leaves only.**Every day**

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Silicon drench (Kalsika) — LAST ONE

MANUAL switch. 400 L clean water → run 3 min → one 100 g sachet into ~300 L, stir until clear → run 8 min → top up to 100 L only → run 3 min until empty → make the normal feed tank. Today only the **2 pm and 5 pm** pulses. **No silicon after today** — it splits fruit.

2 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Hand pollination 06:00–10:00

Open female flowers on nodes 9–12 → pollen from a male flower on the **same plant**. None? Nearest plant of the same variety and generation. Write date + cross on the tag — **this date times the harvest**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Trichoderma cup drench

10 g per 10 L bucket of plain water, stand 15–30 min. Water the bags first, then one cup (250 ml) at the stem base. Morning or late afternoon. Stir every 10 bags. 2 days away from silicon.

2 Keep one fruit + top

Keep **ONE** best fruit per plant. Top just above **leaf 25** (count green leaves, don't guess). No fruit → **don't top**. From today: **no plain water, glass watering or silicon**.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon**. Extra water = split fruit.

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Finish fruit selection & topping

Finish every row. Send Boyd: topped ____ · not yet ____ · no fruit ____.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Finish fruit selection & topping

Finish every row. Send Boyd: topped ____ · not yet ____ · no fruit ____.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S2 stock

EC · pH

2.4 · 5,8–6,3

SOP / TANK

310 g

TIMER · 1.2 L

5 × 7 min · 8 · 10 · 12 · 2 · 4

1 Tonight: make the SIZING stock

No Boroni (4.0). See book pages 27 / 54.

Every day

- 06:30 tank check: EC 2.4 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 175–235 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Thrips on flowers · dead flowers / deformed fruit (boron) · brown gummy ooze at the stem base (Gummy Stem Blight) · fruit fly

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 42 — HST 42 jobs

S3 · Fruit sizing · Sparmanik 4.0

HST

42

1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

1 Calcinit leaf spray

80 g per 16 L spray pack. Early morning or after 4 pm, never in strong sun. Leaves only. **Last one** — no leaf sprays HST 45–59.

2 CHANGE TO SIZING

Tank: SIZING stock + **SOP 535 g** → EC **2.55**. Timer **6 × 8 min**. Support fruit in slings. Mondays: weigh 10 fruit per variety.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity ≥ temperature × 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S3 stockEC · pH
2.55 · 5,8–6,3SOP / TANK
535 gTIMER · 1.6 L
6 × 8 min · 7 · 9 · 11 · 1 · 3 · 5 **No special job today**

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S3 stockEC · pH
2.55 · 5,8–6,3SOP / TANK
535 gTIMER · 1.6 L
6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S3 stockEC · pH
2.55 · 5,8–6,3SOP / TANK
535 gTIMER · 1.6 L
6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5**1 Netting starts****No leaf sprays until HST 59.** Remove old lower leaves with no green left, for airflow.**Every day**

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S3 stockEC · pH
2.55 · 5,8–6,3SOP / TANK
535 gTIMER · 1.6 L
6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S3 stockEC · pH
2.55 · 5,8–6,3SOP / TANK
535 gTIMER · 1.6 L
6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

✓ No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5



No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·

5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5

No special job today

Do the daily routine below carefully.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S3 stock

EC · pH

2.55 · 5,8–6,3

SOP / TANK

535 g

TIMER · 1.6 L

6 × 8 min · 7 · 9 · 11 · 1 · 3 ·
5**1 Trichoderma cup drench**

10 g per 10 L bucket of plain water, stand 15–30 min. Water the bags first, then one cup (250 ml) at the stem base. Morning or late afternoon. Stir every 10 bags. 2 days away from silicon.

2 Tonight: make the RIPEN stock

Its own Ripen drums: Calcinit 8.95 kg (A) · MKP 1.85 kg + MAG-S 6.25 kg + Vitaflex + Fe (B). See book pages 30 / 55.

Every day

- 06:30 tank check: EC 2.55 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 240–320 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

BER (sunken brown patch on the fruit base) · fresh wet splits · a dry bag · fruit fly · mildew

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 CHANGE TO RIPEN**

Tank: RIPEN stock + **SOP 645 g** → EC **2.7**. Timer **4 × 6 min**. Run-off 10–15%. **To the last fruit** — no switch to plain water.

2 4 pm leaf spray — MKP

MKP 48 g per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.**

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 66 — HST 66 jobs

S4 · Ripen & harvest · Sparmanik 4.0

HST
66

1,000 L TANK
S4 stock

EC · pH
2.7 · 5,8–6,3

SOP / TANK
645 g

TIMER · 0.8 L
**4 × 6 min · 8 · 11 am · 2 · 5
pm**

1 4 pm leaf spray — MKP

MKP 48 g per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.**

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit
Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP**MKP 48 g per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**3 Get ready to harvest**

Get scissors, crates, scale and a shaded area ready. Start picking ripe fruit each morning.

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**3 Get ready to harvest**

Get scissors, crates, scale and a shaded area ready. Start picking ripe fruit each morning.

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP**MKP 48 g per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Brix check**Use fruit that **split today**, flesh still clean. Juice from mid-flesh. **Never cut a good fruit.** Send the numbers to Boyd.**3 Get ready to harvest**

Get scissors, crates, scale and a shaded area ready. Start picking ripe fruit each morning.

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



Day 80 — HST 80 jobs

S4 · Ripen & harvest · Sparmanik 4.0

HST

80

1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm

1 4 pm leaf spray — MKP

MKP 48 g per 16 L pack. Leaves only. MKP and Calcinit are never mixed.

2 Morning harvest 06:00–09:30

Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.

Every day

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials



1,000 L TANK

S4 stock

EC · pH

2.7 · 5,8–6,3

SOP / TANK

645 g

TIMER · 0.8 L

4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — MKP****MKP 48 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials

1,000 L TANK
S4 stockEC · pH
2.7 · 5,8–6,3SOP / TANK
645 gTIMER · 0.8 L
4 × 6 min · 8 · 11 am · 2 · 5
pm**1 4 pm leaf spray — Calcinit****Calcinit 80 g** per 16 L pack. Leaves only. **MKP and Calcinit are never mixed.****2 Morning harvest 06:00–09:30**Ripe = full net · ground spot yellow · aroma · slips with a gentle twist · Brix 15+. Cut with scissors, T-stalk, into the shade. **Don't cut anything else.** Grade A–D, weigh, log by plant ID.**Every day**

- 06:30 tank check: EC 2.7 · pH 5.8–6.3 · photo to Boyd before the pump goes on
- Walk every row in the morning: split / rotten fruit → sack → out of the greenhouse, count
- Krisbow meter 8 · 10 · 12 · 2 — humidity $\geq$ temperature $\times$ 2; below → mist leaves 5–10 min (last 3:30 pm)
- Run-off 2–3× a week (not on a silicon day): target 80–120 ml per bag
- Monday: measure the 5 marked plants (height, leaves, fruit)
- **No plain water, glass watering or silicon.** Extra water = split fruit.

Watch for at this stage

Split fruit · fruit fly (CRITICAL) · fruit rot · rodents & birds · Brix under 12 on ripe-looking fruit

Not sure? Stop and call Boyd.

Tank EC / pH

Run-off ml / EC

Fruit picked / binned

Name & initials