

Constraining the geometry of the Universe using Type Ia supernovae

Jacky Cao, Group C3, Physics Problem Solving

Date of report: 28/02/2018

Type Ia supernovae have the unique trait of being standard candles, their light curves can be used in cosmology to calculate and constrain cosmological parameters. In observing Type Ia supernovae and fitting model light curves to such data one can attempt to derive such values. We have monitored, collected, and analysed data for supernova explosions over a period of 34 days. A 16'' and a 0.5 m telescope situated in Durham and La Palma was used for this project. After calculating the magnitudes for a Type Ia (2017hhz) and Type Ia-91bg (2017hle) supernova object, we fitted template light curves with the Python program, *SNooPy*. The quality of fit for the program's `fit()` function was deemed to be acceptable in accordance to the average reduced χ^2 values calculated for the B and V photometric bands, $\chi^2_{\nu,B} \approx 1.38$ and $\chi^2_{V,\nu} \approx 2.95$ - a good fit requiring $\chi^2_{\nu} \approx 1$. The distance modulus to the supernova 2017hhz was calculated by *SNooPy* to be 36.121 ± 0.106 mag, using this value we were able to compute $H_0 = 70 \pm 20 \text{ km s}^{-1} \text{ Mpc}^{-1}$. However, the quoted error negates the meaning of H_0 as it is too large of an uncertainty. In improving the accuracy and uncertainties we suggest that more observations of the supernovae were required, and constraining values should be used for the parameters in *SNooPy*'s templates.

I. INTRODUCTION

a. Supernovae

In cosmology, we can argue that one of the most important objects that we can observe are supernovae.

[1]

b. Cosmological Parameters

sdfsdf

II. RESULTS

asdasd

III. DISCUSSION

asdasd

IV. CONCLUSIONS

sdfsdf

V. ACKNOWLEDGEMENTS

sdfsadf

References

- [1] A. J. Norton, editor. *Observing the Universe*. The Open University, 2004.

Appendix A: Observation Logs