

Producing light curves for supernova explosions and their usage in determining a value for Hubble's Constant

Jacky Cao, AstroLabs, Lab Partner: Duncan Middlemiss

Dates of experiment: 19/10/2017 to 17/03/2017, Date of report: 19/03/2017

We have measured the magnitude of supernova explosions over an extended period of 47 days using 0.5 m and 1.2 m telescopes situated in Durham and La Palma. We have plotted several light curves and identified Type Ia, Type II, and II supernovae. Our fittings have had χ^2 analysis has performed, and it has produced values of χ^2 , χ^2 , χ^2 . We expect our biggest source of uncertainty arose due to the conditions and data analysis that we performed. Using a Type Ia supernova of brightness 00.00 mag, we have managed to produce a value for Hubble's Constant, $H_0 = 74 \text{ kms}^{-1} \text{ Mpc}^{-1}$. We attempted to calculate Einstein's coefficient, Λ , but this was unsuccessful due to redshift of something.

I. INTRODUCTION

In astronomy, one of the most violent and luminous events which can occur is a supernova explosion. At the end of a massive star's lifetime, there is a possibility that the equilibrium configurations for a star will cease to exist after it has ran out of nuclear fuel to burn. This eventually leads to the final explosion, the luminosity of which when at it's peak can be as bright as a small galaxy [4].

Observing these events and measuring their magnitude over a period of time allows us to then plot light curves (magnitude is displayed as a function of time). From visualising this evolution we can draw the conclusion that there is some order in the explosions - they can be grouped together into multiple types as their light curves have similar shapes and features.

Once we have our different groups we can then use them in cosmology for example. One type of supernova explosion called Type Ia's can be used in calculations which leads to a value for Hubble's Constant. The methodology of which will be discussed later in section Ic.

In studying and using supernovae as a tool, we must understand the underlying physics which creates these awesome cosmic events. (??)

a. Supernovae Classification

The two basic classifications for supernovae are Type I and Type II explosions. The main distinction between them both is that the Type I supernova explosion (SNe) has an absence of the Balmer series of hydrogen in their optical spectrum (at maximum light), and Type II does have this hydrogen feature [2, 4]. (??)

In the following sections we will discuss the stages which led to the different types of supernovae. (??)=

With these two groups there are further possible sub-classifications. This is possible as there are visible differences their light curves and also in their spectral features [4]. In Table I, I have highlighted the subtypes which we will explore through experimentation and analysis. (??)

Type	Characteristic
Type Ia	Si II line at 616.0 nm <i>Type Ib and Type Ic also exist</i>
Type IIP	Reaches a plateau in it's light curve
Type IIL	Displays a linear decrease in it's light curve <i>Type IIn and Type IIb also exist</i>

TABLE I: Some of the subclassifications of supernova [4].

1. Type I Supernovae

With the case of Type I supernovae, the general consensus is that presupernova, a white dwarf within a binary system accretes matter from a donor star. The white dwarf increases in it's mass from [ADD VALUE GENERAL STAR HERE] until it reaches a critical point [4, 6]. This limit is thought to be the Chandrasekhar mass, $1.4M_\odot$, after accreting the required matter, the object will be gravitationally unstable [4]. As the material is continually compressed and heated to a temperature of $T \geq 10^9 \text{ K}$, the fusion of carbon and oxygen begins in the core which releases thermonuclear energy. The composition of the star at this point is [...]. This energy that is produced disrupts the star at velocities of the order [give a value and cite it] and prevents a collapse into a neutron star, this disruption leads to a supernova explosion [6].

As we observe the supernova event at a distance away, say from our position on Earth, from it we can perform spectral analysis to find out the chemical composition of what the SNe contains. [??]

One of the subgroups of Type I which are especially important for cosmology are Type Ia SNe. When plotting their magnitudes against time passed we see that their light curves are generally homogeneous. These objects are the most luminous supernovae known, their absolute magnitudes in the B band are typically $M_B = -19.5 \pm 0.1$ [6]. Using light curve relationships and models we can determine precisely the magnitudes of very distant SNe, this has allowed astronomers to determine the redshift-distance relation for redshifts high redshift objects ($z > 1$). Then, with these calculated values it is then possible to estimate the cosmological parameters Ω_0 and Ω_Λ . In doing so, it has been found that using Type Ia SNe has produced agreement with the literature value. so and so (cite paper) have done this and produced values of.

2. Type II Supernovae

For a star to end it's life as a Type II supernova explosion, it's core must collapse to produce the enormous amounts of energy that we associate with a SNe. For stars more massive than $8M_\odot$, their post-main-sequence evolution relies on the continual burning of carbon, oxygen, and silicon burning. As this process continues to heat up the core to larger temperatures, photodisintegration will start to occur. Photons have enough energy to breakup heavier elements such as iron (^{56}Fe) and helium (^4He), this process is endothermic so energy is removed from the gas that would have created the pressure required to support the core of the star [2].

The free electrons which had been supporting the star through degeneracy pressure are now captured by heavy nuclei and by the protons that were produced through photodisintegration. Through the photodisintegration of iron, and combined with electron capture by other elements, most of the core's support disappears and then the core begins to collapse extremely rapidly.

b. Supernova Discovery

When observing the universe the magnitudes that we observe using our instruments are not

In discovering supernovae it is important that the search is not focussed on just one part of the sky as there is potential for SNe to occur in any region of the sky. What is required is a survey which covers the entire visible sky, data can then be compared day-to-day and any new objects that appear can be verified to see if it is a new supernova or not. [reference]

One such survey that employs this is the *All-Sky Automated Survey for Supernovae* (or ASAS-SN). Their setup (??) involves two stations based in Halekalo Observatory in Hawaii, and Cerro Tololo International Observatory in Chile. Together they can produce a survey of the entire visible sky up to a depth of ~ 17 mag [3].

With this case specifically it has been set so that the data is taken automatically, but it requires humans to go over it. Another method that can be used is to observe a location such as a galactic cluster and do essentially the same method.

In the classification of supernovae and in determining their redshifts, spectroscopic observations are required on the objects after they have been discovered [1]. While we can use light curves to find the type of a supernova, it is more effective to use their spectra as that will allow us to be more certain [??].

c. Application to Cosmology

In discovering supernovae we can calculate the rates at which supernovae occur.

Type Ia supernovae are useful to us as they can be classed as standard candles [cite]. With this we can use them to calculate a value for Hubble's Constant, H_0 , and thus calculate the expansion rate of the Universe.

asdasd [5]

$$v = H_0 d, \quad (1)$$

with v as the velocity of the receding object, H_0 as the value for Hubble's find a reference for this.

d. Project Aims

Studying the evolution of the magnitudes of different supernovae will allow us to produce a light curve which can be fitted with known models. Through this we can then discover the type of supernova that we are observing.

The main focus in our experimentation is attempting to collect the data which would form our light curves.

Could we confirm the expansion rate of the universe through our observations of supernovae?

II. OBSERVATIONS

a. Data Collection

Observing supernova explosions requires instruments which are sensitive enough to receive the photons which

have travelled for vast distances before being recorded. In performing such observations (?) our main equipment are telescopes - they come in a wide variety of configurations and each can produce different results.

In collecting the data for our supernova we used three different types of telescopes. Two of them based in Durham on the Department of Physic's Roof, and one in La Palma.

As we were observing objects which were faint we required that we either have a large telescope to collect as much light as possible, or have good seeing conditions which allowed us to collect as much light as possible over a large observation time.

b. Observations Made

Over a period of (?) days we took images of SNe, however this also depended on the conditions and whether it was worth observing if the seeing was bad e.g. obscured by bad weather.

c. Data Analysis

As we collected our supernova day by day we performed various photometry techniques to try and calculate the magnitudes of the explosions.

d. Data Uncertainties

There are uncertainties which are prevalent within our experiment which required to be quantified.

e. Final Data

asdasd

III. ANALYSIS

a. Supernovae Models

The fitting of the templates/models to our observations.

b. Resultuts

sdfsd fsa

IV. DISCUSSION

It is highly unlikely that we will see supernova remnants for the supernova that we have been studying, in the shape of the Crab Nebula. Much longer time periods would be required for this.

[neutrino studying]?

asdasd

V. CONCLUSIONS

sdfsd f sadas

Acknowledgements

I would finally like to thank the coffee shops of Durham for putting up with my buy-one-drink-and-stay-in-there-for-six-hours mentality. It has allowed me to complete my lab report in a slightly buzzed but also happy state.

References

- [1] T. M. Brown et al. Las Cumbres Observatory Global Telescope Network. *Publications of the Astronomical Society of Pacific*, 125:1031–1055, 2013.

- [2] B. W. Carroll and D. A. Ostlie. *An Introduction to Modern Astrophysics*. Pearson, 2nd edition, 2007.
- [3] C. S. Kochanek et al. The All-Sky Automated Survey for Supernovae (ASAS-SN) Light Curve Server V1.0. *Publications of the Astronomical Society of Pacific*, 129(980):104592, 2017.
- [4] M. Longair. *High Energy Astrophysics*. Cambridge University Press, 3rd edition, 2011.
- [5] M. M. Phillips. The Absolute Magnitudes of Type Ia Supernovae. *The Astrophysics Journal*, 413(2):105–108, 1993.
- [6] S. E. Woosley and T. A. Weaver. The Physics of Supernova Explosions. *Annual Review of Astronomy and Astrophysics*, 24:205–253, 1986.

Appendix A - Objects Log

A list of the objects that were chosen to be observed, and then the subsequent notes on them. Not all objects were chosen to be observed for an extended period, the ones noted were observed for a couple of nights to ensure suitability. The subsequent observation logs can be found in Appendix B.

Object	RA	Dec	Magnitude	First Discovered	Type	Notes
*ASASSN-17mz	23:56:21.82	32:27:24.08	14.6	2017/09/30.500	Ia	Too close to galactic nucleus, cannot see
*AT2017hld	22:18:22.849	34:45:08.46	16.1	2017/10/17.339	CV	Cataclysmic Variable, stopped observing
*AT2017hky	11:23:30.514	63:21:59.43	16.2	2017/10/16.640	II	Not viewable from Durham or La Palma
2017hhz	01:44:16.75	12:15:18.00	16.83	2017/10/16.140	Ia	A measured redshift, $z = 0.0392$
AT2017gvb	08:04:42.34	61:31:41.50	17.33	2017/09/26.59	unk	-
*ASASSN-17nb	07:27:37.32	35:36:28.30	17.31	2017/09/25.59	II	Object is dwarfed by brightness of the galaxy
2017hle	01:07:36.060	32:24:30	18.0	2017/10/18.684	Ia-91bg	-
2017hou	04:09:02.140	-01:09:36.40	17.9	2017/10/24.370	Ia	Viewable from La Palma
*AT2017hmw	01:07:16.570	31:25:28.88	17.2	2017/10/19.415	CV	Cataclysmic Variable
2017hpa	04:39:50.750	07:03:54.90	17.9	2017/10/15.346	Ia	Viewable from La Palma
*AT2017hnm	01:42:03.24	42:31:08.50	16.69	2017/10/23.44	unk	Another star in the image dwarfs the SN in brightness
AT2017hpm	08:04:15.100	-00:03:58.03	16.4	2017/10/26.290	unk	-
*AT2017hqa	01:08:59.160	32:38:04.10	17.3	2017/10/26.740	unk	Unobservable, too close to galactic centre
2017hqc	23:23:08.210	10:38:54.63	18.0	2017/10/27.490	Ia	-
*AT2017hrr	11:29:06.490	-08:59:18.56	15.4	2017/10/30.607	unk	Cannot view from Durham or La Palma
*AT2017hhq	00:42:50.230	41:15:27.10	17.7	2017/10/30.599	NV	A nova close to M31
AT2017htb	22:09:38.520	17:39:39.56	15.7	2017/11/02.190	unk	-

TABLE II: Objects that we chose to observe and notes on them. RA is the Right Ascension, given in units of hours : arcminutes : arcseconds. Dec is the Declination, degrees : minutes : seconds. The stated magnitude is the initial magnitude that the object was discovered at (or stated otherwise in the notes), it's units are (??).

Appendix B - Observation Logs

Given below are all the observations and details of those observations made between x and y. With the information provided in the column, **Exposures**, the filters used and the number of exposures taken in the given exposure time.

Date	Object	Time	Exposures	Conditions	Notes
20/10/17	2017hhz	22:25:34 to 22:58:50	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	ASASSN-17nb	02:56:08 to 03:23:36	B: 5, 60s V: 5, 60s	Cloudy	pt5m: Images produced have a lot of noise.
21/10/17	-	-	-	Cloudy	<i>No observations: weather not sufficient in Durham or La Palma for observations.</i>
22/10/17	AT2017hmw	21:52:20 to 22:31:49	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hhz	22:40:20 to 23:19:50	B: 4, 60s V: 12, 60s	Clear	pt5m: -
	ASASSN-17nb	02:42:45 to 03:10:46	B: 5, 120s V: 5, 60s	Clear	pt5m: -
	AT2017gyb	03:12:31 to 03:58:58	B: 5, 180s V: 5, 120s	Clear	pt5m: -
23/10/17	2017hhz	22:53:12 to 23:32:42	B: 5, 120s V: 5, 120s	Cloudy	pt5m: Data produced has FWHM ranging from 1.6 to 9.9.
24/10/17	2017hhz	22:44:01 to 23:23:30	B: 5, 120s V: 5, 120s	Cloudy	pt5m: Data is noisy, potentially another object transited across the frame while observing.
25/10/17	-	-	-	Cloudy	<i>No observations: too cloudy for observations in Durham, and items in pt5m queue were pushed out in favour of other objects.</i>
26/10/17	2017hle	20:54:44 to 21:09:03	B: 4, 60s V: 4, 60s	Clear	FE16: -
	2017hhz	21:18:00 to 21:25:31	B: 4, 60s V: 4, 60s	Clear	FE16: -
	AT2017hmw	21:29:15 to 21:39:36	B: 4, 60s V: 4, 60s	Cloudy	FE16: -
	Messier-7	21:50:23 to 21:54:47	C: 1, 60s	Slightly Cloudy	FE16: Test object for SN discovery.
	Messier-10	21:55:47 to 22:00:13	C: 1, 60s	Cloudy	FE16: Test object for SN discovery.
	AT2017hnm	23:27:22 to 23:41:55	B: 5, 120s V: 3, 120s	Clear	pt5m: -
	AT2017gyb	-	-	-	pt5m: Object pushed out of the queue in favour of others.
27/10/17	AT2017hqa	18:48:12 to 18:52:59	B: 4, 60s	Cloudy	FE16: Object too close to galactic nucleus.
	AT2017hnm	18:56:34 to 19:00:23	C: 1, 60s	Slightly Cloudy	FE16: Object too dim, cloud cover reduced light we were receiving.
	Abell 426	19:04:23 to 19:15:41	C: 1, 60s B: 9, 60s V: 9, 60s	Clear	E14: Object to observe to discover new SN.
	2017hhz	20:11:35 to 20:22:43	B: 4, 60s V: 4, 60s	Cloudy and Windy	E14: Seeing is bad due to the weather.
	2017hle	20:26:18 to 20:37:35	B: 4, 90s V: 4, 90s	Slightly Cloudy	E14: Seeing is bad in these images as well.
	2017hou	23:48:09 to 23:50:14	B: 2, 120s	-	pt5m: Only two frames taken, not sufficient data.

TABLE III: Observing logs for the entire observation period for our experiment.

Date	Object	Time	Exposures	Conditions	Notes
27/10/17	AT2017gyb	02:16:01 to 03:03:05	B: 5, 120s V: 5, 120s	Clear/Cloudy	pt5m: Some images are more noisy due to clouds.
28/10/17	AT2017hnm	21:41:55 to 22:21:25	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hhz	21:28:50 to 21:30:50	B: 1, 120s	Clear	pt5m: -
29/10/17	2017hle	21:22:39 to 21:41:21	B: 10, 120s V: 1, 120s	Clear	pt5m: Mistake on our part to take so many images in B band, and only one in V band.
	AT2017hnm	21:55:34 to 22:35:04	B: 5, 120s V: 5, 120s	Slightly Cloudy	pt5m: -
	2017hhz	23:16:48 to 23:56:18	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hou	00:33:33 to 01:13:02	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hqc	01:16:20 to 01:41:20	B: 4, 120s V: 4, 120s	Clear	pt5m: -
	AT2017gyb	02:18:51 to 02:55:47	V: 13, 180s	Clear	pt5m: Long exposure and large number of exposures chosen as a test to see the quality of stacking them.
30/10/17	2017hle	21:18:56 to 21:58:25	B: 10, 120s V: 10, 120s	Clear	pt5m: -
	2017hhz	22:06:34 to 22:46:04	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	AT2017gyb	02:04:22 to 03:02:50	V: 20, 180s	Clear	pt5m: An human error, we re-selected this object from the previous night instead of the one with the correct number of required frames for each band.
31/10/17	-	-	-	Cloudy	<i>No observations: too cloudy for observations in Durham and La Palma.</i>
01/11/17	2017hle	23:56:53 to 00:15:35	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	AT2017hnm	00:18:23 to 00:57:55	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hhz	01:00:47 to 01:40:17	B: 5, 120s V: 5, 120s	Clear	pt5m: -
	2017hou	01:43:38 to 02:23:08	B: 5, 120s V: 5, 120s	Clear	pt5m: -
02/11/17	-	-	-	Cloudy	<i>No observations: too cloudy for observations in Durham and La Palma.</i>
03/11/17	-	-	-	Cloudy	<i>No observations: too cloudy for observations in Durham and La Palma.</i>
04/11/17	-	-	-	Cloudy	<i>No observations: too cloudy for observations in Durham and La Palma.</i>
05/11/17	ASASSN-17mz	17:53:45 to 18:09:34	B: 8, 60s V: 8, 60s	Clear	FE16: -
	2017hhz	18:50:39 to 19:24:43	B: 8, 120s V: 8, 60s	Clear	FE16: -
	AT2017htb	19:38:14 to 20:02:16	B: 8, 60s V: 8, 60s	Slightly Cloudy	W14: -
	2017hle	20:05:49 to 20:08:28	V: 2, 60s	Cloudy	W14: Stopped taking exposures after viewing the weather.
	2017hqc	20:14:16 to 20:18:25	V: 3, 60s	Cloudy	W14: Seeing if images will be strongly affected by the cloud cover.
	AT2017hhq	20:33:58 to 20:42:26	B: 4, 60s V: 4, 60s	Clear	W14: -

TABLE IV: Observing logs for the entire observation period for our experiment.

Date	Object	Time	Exposures	Conditions	Notes
05/11/17	AT2017hhq	19:38:14 to 20:02:16	B: 3, 30s V: 3, 30s R: 3, 30s	Cloudy	E14: Images taken so that we can produce a colour image.
06/11/17	-	-	-	Cloudy, and Rain	<i>No observations: too cloudy for observations in Durham, and torrential rain in La Palma.</i>
07/11/17	-	-	-	Cloudy and Rain	<i>No observations: too cloudy for observations in Durham, and torrential rain in La Palma.</i>
08/11/17	-	-	-	Cloudy and Rain	<i>No observations: too cloudy for observations in Durham, and torrential rain in La Palma.</i>

TABLE V: Observing logs for the entire observation period for our experiment.

Appendix C - Set of Sample Data