Lecture 14

October 3, 2018

 \blacktriangleright The estimates for β and their confidence intervals etc

The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model.

The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent,

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.
 - 3. We have assumed that all the subjects obey the same linear model.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.
 - We have assumed that all the subjects obey the same linear model. In practice, it may happen that a few subjects do not obey the model.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.
 - We have assumed that all the subjects obey the same linear model. In practice, it may happen that a few subjects do not obey the model. These few observations might change the choice and fit of the model.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.
 - We have assumed that all the subjects obey the same linear model. In practice, it may happen that a few subjects do not obey the model. These few observations might change the choice and fit of the model.

Fortunately, we can do regression diagnostics to check if the data show any evidence of deviating from the assumptions.

- The estimates for β and their confidence intervals etc in the linear model depend on the assumptions underlying the linear model. In particular, note that we have assumed:
 - 1. The errors e_1, \ldots, e_n are independent, have equal variance σ^2 and are normally distributed.
 - 2. We have assumed that the expected value of the response vector Y equals $X\beta$.
 - We have assumed that all the subjects obey the same linear model. In practice, it may happen that a few subjects do not obey the model. These few observations might change the choice and fit of the model.

Fortunately, we can do regression diagnostics to check if the data show any evidence of deviating from the assumptions.

Regression diagnostics should always be performed after regression analysis.

▶ We will first discuss the third part above.

► We will first discuss the third part above. How to detect the presence of unusual observations.

We will first discuss the third part above. How to detect the presence of unusual observations. We will mainly use leverages,

We will first discuss the third part above. How to detect the presence of unusual observations. We will mainly use leverages, jackknife residuals (or standardized predicted residuals)

We will first discuss the third part above. How to detect the presence of unusual observations. We will mainly use leverages, jackknife residuals (or standardized predicted residuals) and Cook's distance for this purpose.

► For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage.

For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?

.

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- It turns out that the ith subject has high leverage if and only if

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- ▶ It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.
- To understand this, we need to know how to define "far from the explanatory variable values of the remaining subjects".

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.
- To understand this, we need to know how to define "far from the explanatory variable values of the remaining subjects". This is done via the notion of *Mahalanobis* distance.

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- ▶ It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.
- ➤ To understand this, we need to know how to define "far from the explanatory variable values of the remaining subjects". This is done via the notion of *Mahalanobis* distance.
- ▶ Given a set of points (also called a point cloud) $v_1, ..., v_n$ on the real line

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.
- To understand this, we need to know how to define "far from the explanatory variable values of the remaining subjects". This is done via the notion of *Mahalanobis* distance.
- Given a set of points (also called a point cloud) v₁,..., v_n on the real line and a test point v, can we quantify how far v is from the set?

- For the bootstrap confidence intervals to be accurate, we need that there are no points with very high leverage. What does high leverage mean?
- It turns out that the ith subject has high leverage if and only if the explanatory variable values for the ith subject are far from the explanatory variable values of the remaining subjects.
- To understand this, we need to know how to define "far from the explanatory variable values of the remaining subjects". This is done via the notion of *Mahalanobis* distance.
- Given a set of points (also called a point cloud) v₁,..., v_n on the real line and a test point v, can we quantify how far v is from the set?
- A natural idea is to look at $|v \bar{v}|^2$ where $\bar{v} = (v_1 + \dots + v_n)/n$.

▶ This makes sense but it does not give a sense of scale.

► This makes sense but it does not give a sense of scale. For example if I get $|v - \bar{v}|^2 = 1$, then does this mean that v is far from the point cloud or near it.

- ► This makes sense but it does not give a sense of scale. For example if I get $|v \bar{v}|^2 = 1$, then does this mean that v is far from the point cloud or near it.
- A better idea is to look at

- ► This makes sense but it does not give a sense of scale. For example if I get $|v \bar{v}|^2 = 1$, then does this mean that v is far from the point cloud or near it.
- A better idea is to look at

$$\frac{(v-\bar{v})^2}{s^2}$$

- ► This makes sense but it does not give a sense of scale. For example if I get $|v \bar{v}|^2 = 1$, then does this mean that v is far from the point cloud or near it.
- A better idea is to look at

$$\frac{(v-\bar{v})^2}{s^2}$$

where

$$s^2 := \frac{1}{n-1} \sum_{j=1}^{n} (v_j - \bar{v})^2$$

▶ This easily generalizes to the multivariate case. Suppose v_1, \ldots, v_n are all in \mathbb{R}^p . Then for a test point $v \in \mathbb{R}^p$, its distance to the point cloud $\{v_1, \ldots, v_n\}$ is measured by the *Mahalanobis distance* defined by

$$(v-\bar{v})^T S^{-1}(v-\bar{v}).$$

where

$$S := \frac{1}{n-1} \sum_{i=1}^{n} (v_i - \bar{v})(v_i - \bar{v})^T$$

Let us now see how this applies to our regression setup with *Y* and *X*.

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the X matrix be denoted by x_i^T .

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- $\blacktriangleright \text{ We can write } x_i^T = [1, z_i^T]$

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- We can write $x_i^T = [1, z_i^T]$ where z_i^T just contains the values of the explanatory variables (without 1) for the *i*th subject.

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- ▶ We can write $x_i^T = [1, z_i^T]$ where z_i^T just contains the values of the explanatory variables (without 1) for the *i*th subject.
- ► The Mahalanobis distance for the *i*th subject is defined as

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- ▶ We can write $x_i^T = [1, z_i^T]$ where z_i^T just contains the values of the explanatory variables (without 1) for the *i*th subject.
- ▶ The Mahalanobis distance for the *i*th subject is defined as

$$\Gamma_i := (z_i - \bar{z})^T S^{-1} (z_i - \bar{z}),$$

- ► Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- ▶ We can write $x_i^T = [1, z_i^T]$ where z_i^T just contains the values of the explanatory variables (without 1) for the *i*th subject.
- ▶ The Mahalanobis distance for the *i*th subject is defined as

$$\Gamma_i := (z_i - \bar{z})^T S^{-1} (z_i - \bar{z}),$$

where

$$S := \frac{1}{n-1} \sum_{i=1}^{n} (z_{i} - \bar{z})(z_{i} - \bar{z})^{T}$$

- Let us now see how this applies to our regression setup with *Y* and *X*.
- Let the *i*th row of the *X* matrix be denoted by x_i^T . Therefore x_i is a $(p+1) \times 1$ vector whose first entry is one and the rest of the entries carry the values of the explanatory variables for the *i*th subject.
- ▶ We can write $x_i^T = [1, z_i^T]$ where z_i^T just contains the values of the explanatory variables (without 1) for the *i*th subject.
- ▶ The Mahalanobis distance for the *i*th subject is defined as

$$\Gamma_i := (z_i - \bar{z})^T S^{-1} (z_i - \bar{z}),$$

where

$$S := \frac{1}{n-1} \sum_{i=1}^{n} (z_j - \bar{z})(z_j - \bar{z})^T$$

and

$$\bar{z}=\sum_{i=1}^{n}z_{i}/n.$$

Γ_i clearly measures how far the ith subject is from the rest of the subjects

 $ightharpoonup \Gamma_i$ clearly measures how far the *i*th subject is from the rest of the subjects in terms of the values of the explanatory variables.

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- ▶ It turns out that the leverage for the *i*th subject

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- lt turns out that the leverage for the *i*th subject, h_{ii} ,

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- ▶ It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- ▶ It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

$$\Gamma_i = (n-1)h_{ii} - \frac{n-1}{n}.\tag{1}$$

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

$$\Gamma_i = (n-1)h_{ii} - \frac{n-1}{n}.$$
 (1)

► This explains the second interpretation of leverage.

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

$$\Gamma_i = (n-1)h_{ii} - \frac{n-1}{n}.$$
 (1)

This explains the second interpretation of leverage. For the above formula to hold, it is necessary that there be an intercept term in the model.

- Γ_i clearly measures how far the ith subject is from the rest of the subjects in terms of the values of the explanatory variables.
- ▶ It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

$$\Gamma_i = (n-1)h_{ii} - \frac{n-1}{n}.$$
 (1)

- This explains the second interpretation of leverage. For the above formula to hold, it is necessary that there be an intercept term in the model.
- In other words, this formula does not hold without an intercept term.

- $ightharpoonup \Gamma_i$ clearly measures how far the *i*th subject is from the rest of the subjects in terms of the values of the explanatory variables.
- ▶ It turns out that the leverage for the *i*th subject , h_{ii} , is related to Γ_i by the following simple expression:

$$\Gamma_i = (n-1)h_{ii} - \frac{n-1}{n}.$$
 (1)

- ► This explains the second interpretation of leverage. For the above formula to hold, it is necessary that there be an intercept term in the model.
- In other words, this formula does not hold without an intercept term. We will not go over the proof of this formula. It can be easily verified in R.

Note that one consequence of (1) is the fact that $h_{ii} \ge 1/n$ for every i.

Note that one consequence of (1) is the fact that $h_{ii} \ge 1/n$ for every i. This is because Γ_i is always nonnegative.

Note that one consequence of (1) is the fact that $h_{ii} \ge 1/n$ for every i. This is because Γ_i is always nonnegative. The leverages therefore always lie between 1/n and 1 whenever there is an intercept term in the linear model.