

Lecture 4

September 4, 2018

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .
- ▶ There are n subjects and data is collected on the variables from these subjects.

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .
- ▶ There are n subjects and data is collected on the variables from these subjects.
- ▶ Data on the response variable is y_1, \dots, y_n .

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .
- ▶ There are n subjects and data is collected on the variables from these subjects.
- ▶ Data on the response variable is y_1, \dots, y_n .
- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .
- ▶ There are n subjects and data is collected on the variables from these subjects.
- ▶ Data on the response variable is y_1, \dots, y_n .
- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data for the i th subject is $(y_i, x_{i1}, \dots, x_{ip})$.

Multiple Regression

- ▶ There is a response variable y and p explanatory variables x_1, \dots, x_p . The goal is understand the relationship between y and x_1, \dots, x_p .
- ▶ There are n subjects and data is collected on the variables from these subjects.
- ▶ Data on the response variable is y_1, \dots, y_n .
- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data for the i th subject is $(y_i, x_{i1}, \dots, x_{ip})$.
- ▶ The linear regression model assumes that these are independent with

$$\mathbb{E}(y_i | x_{i1}, \dots, x_{ip}) = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip}, \quad \text{for } i = 1, \dots, n.$$

Multiple Regression

- Equivalently, we can write the model as

$$y_i = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip} + e_i, \quad \text{with } \mathbb{E}(e_i | x_{i1}, \dots, x_{ip}) = 0.$$

Estimation of $\beta_0, \beta_1, \dots, \beta_p$

- As before, we use least squares: Estimate $\beta_0, \beta_1, \dots, \beta_p$ by the minimizers of

$$s(b) = \sum_{i=1}^n (y_i - b_0 - b_1 x_{i1} - \dots b_p x_{ip})^2 \quad (1)$$

over b_0, \dots, b_p .

Estimation of $\beta_0, \beta_1, \dots, \beta_p$

- ▶ As before, we use least squares: Estimate $\beta_0, \beta_1, \dots, \beta_p$ by the minimizers of

$$s(b) = \sum_{i=1}^n (y_i - b_0 - b_1 x_{i1} - \dots b_p x_{ip})^2 \quad (1)$$

over b_0, \dots, b_p .

- ▶ Like in the simple regression case, we can try to take partial derivatives with respect to b_0, \dots, b_p , set them equal to zero and then solve the resulting equations.

Estimation of $\beta_0, \beta_1, \dots, \beta_p$

- ▶ As before, we use least squares: Estimate $\beta_0, \beta_1, \dots, \beta_p$ by the minimizers of

$$s(b) = \sum_{i=1}^n (y_i - b_0 - b_1 x_{i1} - \dots b_p x_{ip})^2 \quad (1)$$

over b_0, \dots, b_p .

- ▶ Like in the simple regression case, we can try to take partial derivatives with respect to b_0, \dots, b_p , set them equal to zero and then solve the resulting equations.
- ▶ Doing this directly using the expression (1) is painful. Things are greatly simplified if one uses matrices and vectors.

Estimation of $\beta_0, \beta_1, \dots, \beta_p$

- ▶ As before, we use least squares: Estimate $\beta_0, \beta_1, \dots, \beta_p$ by the minimizers of

$$s(b) = \sum_{i=1}^n (y_i - b_0 - b_1 x_{i1} - \dots - b_p x_{ip})^2 \quad (1)$$

over b_0, \dots, b_p .

- ▶ Like in the simple regression case, we can try to take partial derivatives with respect to b_0, \dots, b_p , set them equal to zero and then solve the resulting equations.
- ▶ Doing this directly using the expression (1) is painful. Things are greatly simplified if one uses matrices and vectors.
- ▶ Data on the response variable, y_1, \dots, y_n , are represented by the column vector $Y = (y_1, \dots, y_n)^T$ (the T here stands for transpose).

- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.

- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data on all the explanatory variables is represented by the $n \times (p + 1)$ matrix X whose first column consists of ones, second column has the values $x_{11}, x_{21}, \dots, x_{n1}$ corresponding to the first explanatory variable, third column has values for the second explanatory variable and so on.

- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data on all the explanatory variables is represented by the $n \times (p + 1)$ matrix X whose first column consists of ones, second column has the values $x_{11}, x_{21}, \dots, x_{n1}$ corresponding to the first explanatory variable, third column has values for the second explanatory variable and so on.
- ▶ With this notation, the sum of squares in (1) can be rewritten as

$$s(b) = \|Y - Xb\|_2^2.$$

where the norm of the vector x is defined as

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}. \text{ Note the equality } \|x\|_2^2 = x^T x.$$

- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data on all the explanatory variables is represented by the $n \times (p + 1)$ matrix X whose first column consists of ones, second column has the values $x_{11}, x_{21}, \dots, x_{n1}$ corresponding to the first explanatory variable, third column has values for the second explanatory variable and so on.
- ▶ With this notation, the sum of squares in (1) can be rewritten as

$$s(b) = \|Y - Xb\|_2^2.$$

where the norm of the vector x is defined as

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}. \text{ Note the equality } \|x\|_2^2 = x^T x.$$

- ▶ Using this, we can write

$$s(b) = (Y - Xb)^T (Y - Xb) = Y^T Y - 2b^T X^T Y + b^T X^T X b.$$

prove : <https://math.stackexchange.com/questions/2753210/when-can-we-say-that-atb-bta>

- ▶ Data on the j th explanatory variable x_j is $x_{1j}, x_{2j}, \dots, x_{nj}$.
- ▶ The data on all the explanatory variables is represented by the $n \times (p + 1)$ matrix X whose first column consists of ones, second column has the values $x_{11}, x_{21}, \dots, x_{n1}$ corresponding to the first explanatory variable, third column has values for the second explanatory variable and so on.
- ▶ With this notation, the sum of squares in (1) can be rewritten as

$$s(b) = \|Y - Xb\|_2^2.$$

where the norm of the vector x is defined as

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}. \text{ Note the equality } \|x\|_2^2 = x^T x.$$

- ▶ Using this, we can write

$$s(b) = (Y - Xb)^T (Y - Xb) = Y^T Y - 2Y^T Xb + b^T X^T Xb.$$

- ▶ This can be minimized via calculus. Take partial derivatives with respect to i for $i = 0, 1, \dots, p$ and equate them to 0. It is easy to check that

$$\nabla S(b) = 2X^T Xb - 2X^T y$$

$$\begin{aligned}
S(b) &= \sum e_i^2 = e'e = (y - Xb)'(y - Xb) \\
&= y'y - y'Xb - b'X'y + b'X'Xb.
\end{aligned}
\tag{3.6}$$

Derivation of least squares estimator

The minimum of $S(b)$ is obtained by setting the derivatives of $S(b)$ equal to zero. Note that the function $S(b)$ has scalar values, whereas b is a column vector with k components. So we have k first order derivatives and we will follow the convention to arrange them in a column vector. The second and third terms of the last expression in (3.6) are equal (a 1×1 matrix is always symmetric) and may be replaced by $-2b'X'y$. This is a linear expression in the elements of b and so the vector of derivatives equals $-2X'y$. The last term of (3.6) is a quadratic form in the elements of b . The vector of first order derivatives of this term $b'X'Xb$ can be written as $2X'Xb$. The proof of this result is left as an exercise (see Exercise 3.1). To get the idea we consider the case $k = 2$ and we denote the elements of $X'X$ by c_{ij} , $i, j = 1, 2$, with $c_{12} = c_{21}$. Then $b'X'Xb = c_{11}b_1^2 + c_{22}b_2^2 + 2c_{12}b_1b_2$. The derivative with respect to b_1 is $2c_{11}b_1 + 2c_{12}b_2$ and the derivative with respect to b_2 is $2c_{12}b_1 + 2c_{22}b_2$. When we arrange these two partial derivatives in a 2×1 vector, this can be written as $2X'Xb$. See Appendix A (especially Examples A.10 and A.11 in Section A.7) for further computational details and illustrations.

T

► where

$$\nabla S(b) = \left(\frac{\partial S(b)}{\partial \beta_0}, \dots, \frac{\partial S(b)}{\partial \beta_p} \right).$$

denotes the gradient of $S(b)$ with respect to $b = (b_1, \dots, b_p)^T$.

- ▶ where

$$\nabla S(b) = \left(\frac{\partial S(b)}{\partial \beta_0}, \dots, \frac{\partial S(b)}{\partial \beta_p} \right).$$

denotes the gradient of $S(b)$ with respect to $b = (b_1, \dots, b_p)^T$.

- ▶ It follows therefore that the minimizer of $S(b)$ satisfies the equality

$$X^T X b = X^T y.$$

- ▶ where

$$\nabla S(b) = \left(\frac{\partial S(b)}{\partial \beta_0}, \dots, \frac{\partial S(b)}{\partial \beta_p} \right).$$

denotes the gradient of $S(b)$ with respect to $b = (b_1, \dots, b_p)^T$.

- ▶ It follows therefore that the minimizer of $S(b)$ satisfies the equality

$$X^T X b = X^T y.$$

- ▶ This gives p linear equations for the $p + 1$ unknowns b_0, \dots, b_p .

- ▶ where

$$\nabla S(b) = \left(\frac{\partial S(b)}{\partial \beta_0}, \dots, \frac{\partial S(b)}{\partial \beta_p} \right).$$

denotes the gradient of $S(b)$ with respect to $b = (b_1, \dots, b_p)^T$.

- ▶ It follows therefore that the minimizer of $S(b)$ satisfies the equality

$$X^T X b = X^T y.$$

- ▶ This gives p linear equations for the $p + 1$ unknowns b_0, \dots, b_p .
- ▶ This important set of equations are called normal equations.

- ▶ where

$$\nabla S(b) = \left(\frac{\partial S(b)}{\partial \beta_0}, \dots, \frac{\partial S(b)}{\partial \beta_p} \right).$$

denotes the gradient of $S(b)$ with respect to $b = (b_1, \dots, b_p)^T$.

- ▶ It follows therefore that the minimizer of $S(b)$ satisfies the equality

$$X^T X b = X^T y.$$

- ▶ This gives p linear equations for the $p + 1$ unknowns b_0, \dots, b_p .
- ▶ This important set of equations are called normal equations.
- ▶ Their solution, denoted by $\hat{\beta} = (\hat{\beta}_0, \dots, \hat{\beta}_p)^T$ gives an estimate of β called the least squares estimate.

- ▶ When $X^T X$ is invertible, we have a unique solution to the normal equations that is given by $(X^T X)^{-1} X^T Y$. Thus our least squares estimate is

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

- ▶ When $X^T X$ is invertible, we have a unique solution to the normal equations that is given by $(X^T X)^{-1} X^T Y$. Thus our least squares estimate is

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

- ▶ What happens when $X^T X$ is not invertible?

- ▶ When $X^T X$ is invertible, we have a unique solution to the normal equations that is given by $(X^T X)^{-1} X^T Y$. Thus our least squares estimate is

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

- ▶ What happens when $X^T X$ is not invertible? We shall deal with this later.

- ▶ When $X^T X$ is invertible, we have a unique solution to the normal equations that is given by $(X^T X)^{-1} X^T Y$. Thus our least squares estimate is

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

- ▶ What happens when $X^T X$ is not invertible? We shall deal with this later.
- ▶ Our next goal is to understand the properties of the least squares estimator. Before that, let us take a detour and learn some formulae for dealing with mean and variance of random vectors.

Basic Mean and Covariance Formulae for Random Vectors

- ▶ We next want to explore properties of $\hat{\beta} = (X^T X)^{-1} X^T Y$ as an estimator of β in the linear model. For this we need a few facts about means and covariances.

Basic Mean and Covariance Formulae for Random Vectors

- ▶ We next want to explore properties of $\hat{\beta} = (X^T X)^{-1} X^T Y$ as an estimator of β in the linear model. For this we need a few facts about means and covariances.
- ▶ Let $Z = (Z_1, \dots, Z_k)^T$ be a random vector. Its expectation $\mathbb{E}Z$ is defined as a vector whose i th entry is the expectation of Z_i i.e., $\mathbb{E}Z = (\mathbb{E}Z_1, \mathbb{E}Z_2, \dots, \mathbb{E}Z_k)^T$.

Basic Mean and Covariance Formulae for Random Vectors

- ▶ We next want to explore properties of $\hat{\beta} = (X^T X)^{-1} X^T Y$ as an estimator of β in the linear model. For this we need a few facts about means and covariances.
- ▶ Let $Z = (Z_1, \dots, Z_k)^T$ be a random vector. Its expectation $\mathbb{E}Z$ is defined as a vector whose i th entry is the expectation of Z_i i.e., $\mathbb{E}Z = (\mathbb{E}Z_1, \mathbb{E}Z_2, \dots, \mathbb{E}Z_k)^T$.
- ▶ The covariance matrix of Z , denoted by $\text{Cov}(Z)$, is a $k \times k$ matrix whose (i, j) th entry is the covariance between Z_i and Z_j .

Basic Mean and Covariance Formulae for Random Vectors

- ▶ We next want to explore properties of $\hat{\beta} = (X^T X)^{-1} X^T Y$ as an estimator of β in the linear model. For this we need a few facts about means and covariances.
- ▶ Let $Z = (Z_1, \dots, Z_k)^T$ be a random vector. Its expectation $\mathbb{E}Z$ is defined as a vector whose i th entry is the expectation of Z_i i.e., $\mathbb{E}Z = (\mathbb{E}Z_1, \mathbb{E}Z_2, \dots, \mathbb{E}Z_k)^T$.
- ▶ The covariance matrix of Z , denoted by $\text{Cov}(Z)$, is a $k \times k$ matrix whose (i, j) th entry is the covariance between Z_i and Z_j .
- ▶ If $W = (W_1, \dots, W_m)^T$ is another random vector, the covariance matrix between Z and W , denoted by $\text{Cov}(Z, W)$, is a $k \times m$ matrix whose (i, j) th entry is the covariance between Z_i and W_j . Note then that, $\text{Cov}(Z, Z) = \text{Cov}(Z)$.

Basic Mean and Covariance Formulae for Random Vectors

The following formulae are very important:

Basic Mean and Covariance Formulae for Random Vectors

The following formulae are very important:

- ▶ $\mathbb{E}(AZ + c) = A\mathbb{E}(Z) + c$ for any constant matrix A and any constant vector c .

Basic Mean and Covariance Formulae for Random Vectors

The following formulae are very important:

- ▶ $\mathbb{E}(AZ + c) = A\mathbb{E}(Z) + c$ for any constant matrix A and any constant vector c .
- ▶ $\text{Cov}(AZ + c) = A\text{Cov}(Z)A^T$ for any constant matrix A and any constant vector c .

Basic Mean and Covariance Formulae for Random Vectors

The following formulae are very important:

- ▶ $\mathbb{E}(AZ + c) = A\mathbb{E}(Z) + c$ for any constant matrix A and any constant vector c .
- ▶ $\text{Cov}(AZ + c) = A\text{Cov}(Z)A^T$ for any constant matrix A and any constant vector c .
- ▶ $\text{Cov}(AZ + c, BW + d) = A\text{Cov}(Z, W)B^T$ for any pair of constant matrices A and B and any pair of constant vectors c and d .

Assumptions

- ▶ The linear model is

$$Y = X\beta + e.$$

Assumptions

- ▶ The linear model is

$$Y = X\beta + e.$$

- ▶ The zero conditional mean assumption on e can be written as $\mathbb{E}(e|X) = 0$. For variance calculations, we also make the assumption that $\text{Cov}(e|X) = \sigma^2 I_n$.

Assumptions

- ▶ The linear model is

$$Y = X\beta + e.$$

- ▶ The zero conditional mean assumption on e can be written as $\mathbb{E}(e|X) = 0$. For variance calculations, we also make the assumption that $\text{Cov}(e|X) = \sigma^2 I_n$.
- ▶ Because of the above formulae, we can write

$$\mathbb{E}(Y|X) = X\beta, \quad \text{cov}(Y|X) = \sigma^2 I_n.$$

Properties of the Least Squares Estimator

- ▶ Assume that $X^T X$ is invertible (equivalently, that X has rank $p + 1$) and consider the least squares estimator

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

Properties of the Least Squares Estimator

- ▶ Assume that $X^T X$ is invertible (equivalently, that X has rank $p + 1$) and consider the least squares estimator

$$\hat{\beta} = (X^T X)^{-1} X^T Y.$$

- ▶ What properties does $\hat{\beta}$ have as an estimator of β ?

Linearity

- ▶ An estimator of β is said to be linear if it can be written as AY for some matrix A .

Linearity

- ▶ An estimator of β is said to be linear if it can be written as AY for some matrix A .
- ▶ Clearly $\hat{\beta} = (X^T X)^{-1} X^T Y$ is of this form and hence it is a linear estimator of β .

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

$$\mathbb{E}(\hat{\beta}) = \mathbb{E}((X^T X)^{-1} X^T Y)$$

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

$$\begin{aligned}\mathbb{E}(\hat{\beta}) &= \mathbb{E}((X^T X)^{-1} X^T Y) \\ &= (X^T X)^{-1} X^T \mathbb{E}(Y)\end{aligned}$$

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

$$\begin{aligned}\mathbb{E}(\hat{\beta}) &= \mathbb{E}((X^T X)^{-1} X^T Y) \\ &= (X^T X)^{-1} X^T \mathbb{E}(Y) \\ &= (X^T X)^{-1} X^T X \beta\end{aligned}$$

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

$$\begin{aligned}\mathbb{E}(\hat{\beta}) &= \mathbb{E}((X^T X)^{-1} X^T Y) \\ &= (X^T X)^{-1} X^T \mathbb{E}(Y) \\ &= (X^T X)^{-1} X^T X \beta \\ &= \beta.\end{aligned}$$

Unbiasedness

- ▶ An estimator for a parameter is said to be unbiased if its expectation equals the parameter (for all values of the parameter).
- ▶ The expectation of the least squares estimator is (using the formula for expectation: $\mathbb{E}AZ = A\mathbb{E}Z$)

$$\begin{aligned}\mathbb{E}(\hat{\beta}) &= \mathbb{E}((X^T X)^{-1} X^T Y) \\ &= (X^T X)^{-1} X^T \mathbb{E}(Y) \\ &= (X^T X)^{-1} X^T X \beta \\ &= \beta.\end{aligned}$$

- ▶ In particular, this means that $\mathbb{E}\hat{\beta}_i = \beta_i$ for each i which implies that each $\hat{\beta}_i$ is an unbiased estimator of β_i .

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

$$Cov(\hat{\beta}) = Cov((X^T X)^{-1} X^T Y)$$

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

$$\begin{aligned}Cov(\hat{\beta}) &= Cov((X^T X)^{-1} X^T Y) \\&= (X^T X)^{-1} X^T Cov(Y) X (X^T X)^{-1}\end{aligned}$$

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

$$\begin{aligned}Cov(\hat{\beta}) &= Cov((X^T X)^{-1} X^T Y) \\&= (X^T X)^{-1} X^T Cov(Y) X (X^T X)^{-1} \\&= (X^T X)^{-1} X^T (\sigma^2 I_n) X (X^T X)^{-1}\end{aligned}$$

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)AT$:

$$\begin{aligned}Cov(\hat{\beta}) &= Cov((X^T X)^{-1} X^T Y) \\&= (X^T X)^{-1} X^T Cov(Y) X (X^T X)^{-1} \\&= (X^T X)^{-1} X^T (\sigma^2 I_n) X (X^T X)^{-1} \\&= \sigma^2 (X^T X)^{-1}.\end{aligned}$$

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

$$\begin{aligned}Cov(\hat{\beta}) &= Cov((X^T X)^{-1} X^T Y) \\&= (X^T X)^{-1} X^T Cov(Y) X (X^T X)^{-1} \\&= (X^T X)^{-1} X^T (\sigma^2 I_n) X (X^T X)^{-1} \\&= \sigma^2 (X^T X)^{-1}.\end{aligned}$$

- ▶ In particular, the variance of $\hat{\beta}_i$ equals σ^2 multiplied by the i th diagonal element of $(X^T X)^{-1}$.

Covariance Matrix

- ▶ The Covariance matrix of the estimator $\hat{\beta}$ can be easily calculated using the formula: $Cov(AZ) = ACov(Z)A^T$:

$$\begin{aligned}Cov(\hat{\beta}) &= Cov((X^T X)^{-1} X^T Y) \\&= (X^T X)^{-1} X^T Cov(Y) X (X^T X)^{-1} \\&= (X^T X)^{-1} X^T (\sigma^2 I_n) X (X^T X)^{-1} \\&= \sigma^2 (X^T X)^{-1}.\end{aligned}$$

- ▶ In particular, the variance of $\hat{\beta}_i$ equals σ^2 multiplied by the i th diagonal element of $(X^T X)^{-1}$.
- ▶ Once we learn how to estimate σ^2 , we can use this to obtain standard errors for $\hat{\beta}_i$.

Optimality - The Gauss-Markov Theorem

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator).

Optimality - The Gauss-Markov Theorem

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator).
- ▶ This means that $\hat{\beta}$ is the best estimator among all linear and unbiased estimators of β . Here, best is in terms of variance.

Optimality - The Gauss-Markov Theorem

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator).
- ▶ This means that $\hat{\beta}$ is the best estimator among all linear and unbiased estimators of β . Here, best is in terms of variance.
- ▶ This implies that $\hat{\beta}_i$ has the smallest variance among all linear and unbiased estimators of i for every i .

Optimality - The Gauss-Markov Theorem

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator).
- ▶ This means that $\hat{\beta}$ is the best estimator among all linear and unbiased estimators of β . Here, best is in terms of variance.
- ▶ This implies that $\hat{\beta}_i$ has the smallest variance among all linear and unbiased estimators of i for every i .
- ▶ The Gauss-Markov theorem is actually quite simple to prove. We shall prove it in the next class.