Lecture 2

August 28, 2018

Simple Linear Regression

► Here there is only one explanatory variable x. We denote the data in simple linear regression as $(x_1, y_1), \ldots, (x_n, y_n)$.

Simple Linear Regression

- ► Here there is only one explanatory variable x. We denote the data in simple linear regression as $(x_1, y_1), \ldots, (x_n, y_n)$.
- We assume that these are independent with the relation of y_i to x_i expressed as

$$\mathbb{E}(y_i|x_i) = \beta_0 + \beta_1 x_i, \quad i = 1, \ldots, n.$$

Simple Linear Regression

- ► Here there is only one explanatory variable x. We denote the data in simple linear regression as $(x_1, y_1), \ldots, (x_n, y_n)$.
- We assume that these are independent with the relation of y_i to x_i expressed as

$$\mathbb{E}(y_i|x_i) = \beta_0 + \beta_1 x_i, \quad i = 1, \ldots, n.$$

Why can we also say E(Y)=Xb, ignoring conditional expectation?

We can alternatively write the simple linear regression model as

$$y_i = \beta_0 + \beta_1 x_i + e_i$$
, where $\mathbb{E}(e_i|x_i) = 0$.

▶ How does one estimate the parameters β_0 and β_1 ?

- ▶ How does one estimate the parameters β_0 and β_1 ?
- ➤ The most popular way is to use the method of least squares.

- ▶ How does one estimate the parameters β_0 and β_1 ?
- ► The most popular way is to use the method of least squares.
- ► The least squares estimates of β_0 and β_1 are those values b_0 and b_1 that minimize:

$$Q(b_0,b_1) = \sum_{i=1}^n (y_i - b_0 - b_1 x_i)^2$$


- ▶ How does one estimate the parameters β_0 and β_1 ?
- The most popular way is to use the method of least squares.
- The least squares estimates of β_0 and β_1 are those values b_0 and b_1 that minimize:

$$Q(b_0,b_1) = \sum_{i=1}^n (y_i - b_0 - b_1 x_i)^2$$

► These estimates can be computed by simply differentiating Q with respect to b₀ and b₁ and setting the partial derivatives equal to zero. This gives

$$\frac{\partial Q}{\partial b_0} = -2 \sum_{i=1}^n (y_i - b_0 - b_1 x_i) = 0,$$

and



$$\frac{\partial Q}{\partial b_1} = -2\sum_{i=1}^n x_i(y_i - b_0 - b_1x_i) = 0.$$

► These equations can be equivalently written as

$$nb_0 + b_1 \sum_{i=1}^n x_i = \sum_{i=1}^n y_i$$
, and $b_0 \sum_{i=1}^n x_i + b_1 \sum_{i=1}^n x_i^2 = \sum_{i=1}^n x_i y_i$.

$$\frac{\partial Q}{\partial b_1} = -2\sum_{i=1}^n x_i(y_i - b_0 - b_1x_i) = 0.$$

► These equations can be equivalently written as

$$nb_0 + b_1 \sum_{i=1}^n x_i = \sum_{i=1}^n y_i$$
, and $b_0 \sum_{i=1}^n x_i + b_1 \sum_{i=1}^n x_i^2 = \sum_{i=1}^n x_i y_i$.

► The above two equations are known as the normal equations. These can be easily solved to give

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{\sum_{i=1}^n (x_i - \bar{x})y_i}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

and

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x},$$

where we used the notation $\bar{y} = \frac{1}{n} \sum_{i=1}^{n} y_i$, and $\bar{x} = \frac{1}{n} \sum_{i=1}^{n} x_i$.

 \triangleright β_0 and β_1 are called the Least Squares estimates of β_0 and β_1 respectively. Sometimes they are also called OLS estimates (OLS stands for Ordinary Least Squares).

- β_0 and β_1 are called the Least Squares estimates of β_0 and β_1 respectively. Sometimes they are also called OLS estimates (OLS stands for Ordinary Least Squares).
- estimates (OLS stands for Ordinary Least Squares).

 Note that for these estimates to make sense, we need $\sum_{i=1}^{n} (x_i \bar{x})^2 > 0$ which means that the *x*-variables are

not all the same.

▶ The OLS regression line is given by $y = \hat{\beta}_0 + \hat{\beta}_1 x$. This is the line with slope $\hat{\beta}_1$ and intercept $\hat{\beta}_0$. The regression line can be used to predict y given x.

- ► The OLS regression line is given by $y = \hat{\beta}_0 + \hat{\beta}_1 x$. This is the line with slope $\hat{\beta}_1$ and intercept $\hat{\beta}_0$. The regression line can be used to predict y given x.
- ► There is a very interesting alternative way of writing the regression line. This uses the following notation. Let

$$s_x := \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}, \text{ and } s_y := \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2}.$$

- ► The OLS regression line is given by $y = \hat{\beta}_0 + \hat{\beta}_1 x$. This is the line with slope $\hat{\beta}_1$ and intercept $\hat{\beta}_0$. The regression line can be used to predict y given x.
- ► There is a very interesting alternative way of writing the regression line. This uses the following notation. Let

$$s_x := \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}, \text{ and } s_y := \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2}.$$

► Also let

$$r := \frac{\sum_{i=1}^{n} (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^{n} (x_i - \bar{x})^2 \sum_{i=1}^{n} (y_i - \bar{y})^2}} = \frac{1}{n} \sum_{i=1}^{n} \left(\frac{x_i - \bar{x}}{s_x} \right) \left(\frac{y_i - \bar{y}}{s_y} \right),$$

▶ s_x and s_y denote the sample standard deviations of x and y respectively and r denotes the sample correlation coefficient between x and y. It is always true that $0 \le |r| \le 1$ (why?).

- ▶ s_x and s_y denote the sample standard deviations of x and y respectively and r denotes the sample correlation coefficient between x and y. It is always true that $0 \le |r| \le 1$ (why?).
- With this notation, it is easy to see that $\hat{\beta}_1 = rs_y/s_x$.

- ▶ s_x and s_y denote the sample standard deviations of x and y respectively and r denotes the sample correlation coefficient between x and y. It is always true that $0 \le |r| \le 1$ (why?).
- With this notation, it is easy to see that $\hat{\beta}_1 = rs_y/s_x$. Hence, the regression line equals

$$\frac{y-\bar{y}}{s_y}=r\left(\frac{x-\bar{x}}{s_x}\right).$$

- ▶ s_x and s_y denote the sample standard deviations of x and y respectively and r denotes the sample correlation coefficient between x and y. It is always true that $0 \le |r| \le 1$ (why?).
- With this notation, it is easy to see that $\hat{\beta}_1 = rs_y/s_x$. Hence, the regression line equals

$$\frac{y-\bar{y}}{s_y}=r\left(\frac{x-\bar{x}}{s_x}\right).$$

Note that the regression line passes through the point (\bar{y}, \bar{x}) .

▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.
- ▶ The estimate $\hat{\beta}_1$ is unbiased for β_1 .

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.
- ► The estimate $\hat{\beta}_1$ is unbiased for β_1 . To see this, let us first compute

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.
- ► The estimate $\hat{\beta}_1$ is unbiased for β_1 . To see this, let us first compute

$$\mathbb{E}(\hat{\beta}_1|X) \,=\, \mathbb{E}\left(\tfrac{\sum_{i=1}^n (x_i - \bar{x})y_i}{\sum_{i=1}^n (x_i - \bar{x})^2}|X\right) =$$

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.
- ▶ The estimate $\hat{\beta}_1$ is unbiased for β_1 . To see this, let us first compute

$$\mathbb{E}(\hat{\beta}_1|X) = \mathbb{E}\left(\frac{\sum_{i=1}^{n}(x_i-\bar{x})y_i}{\sum_{i=1}^{n}(x_i-\bar{x})^2}|X\right) = \frac{\sum_{i=1}^{n}(x_i-\bar{x})\mathbb{E}(y_i|X)}{\sum_{i=1}^{n}(x_i-\bar{x})^2}$$

- ▶ Recall the linear regression model described before. The quantities β_0 and β_1 are parameters of the model. These are estimated by $\hat{\beta}_0$ and $\hat{\beta}_1$. How good are these estimators?
- We write X for the set of explanatory variable values $\{x_1, \ldots, x_n\}$. Under the assumptions of the linear model, it is easy to see that $\mathbb{E}(y_i|X) = \beta_0 + \beta_1 x_i$.
- ▶ The estimate $\hat{\beta}_1$ is unbiased for β_1 . To see this, let us first compute

$$\mathbb{E}(\hat{\beta}_{1}|X) = \mathbb{E}\left(\frac{\sum_{i=1}^{n}(x_{i}-\bar{x})y_{i}}{\sum_{i=1}^{n}(x_{i}-\bar{x})^{2}}|X\right) = \frac{\sum_{i=1}^{n}(x_{i}-\bar{x})\mathbb{E}(y_{i}|X)}{\sum_{i=1}^{n}(x_{i}-\bar{x})^{2}} \\
= \frac{\sum_{i=1}^{n}(x_{i}-\bar{x})(\beta_{0}+\beta_{1}x_{i})}{\sum_{i=1}^{n}(x_{i}-\bar{x})^{2}}.$$

▶ I will leave it as an exercise to verify that the last expression equals β_1 .

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$.

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 .

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 . This automatically implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?).

- I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is

also holds for $\hat{\beta}_0$:

- conditionally (on X) unbiased for β_1 . This automatically
- implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?). Unbiasedness (conditionally on X and unconditionally)

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 . This automatically implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?).
- ▶ Unbiasedness (conditionally on X and unconditionally) also holds for $\hat{\beta}_0$:

$$\mathbb{E}(\hat{\beta}_0|X) =$$

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 . This automatically implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?).
- ▶ Unbiasedness (conditionally on X and unconditionally) also holds for $\hat{\beta}_0$:

$$\mathbb{E}(\hat{eta}_0|X) = \mathbb{E}(\bar{y} - \bar{x}\hat{eta}_1|X) =$$

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 . This automatically implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?).
- ▶ Unbiasedness (conditionally on X and unconditionally) also holds for $\hat{\beta}_0$:

$$\mathbb{E}(\hat{\beta}_0|X) = \mathbb{E}(\bar{y} - \bar{x}\hat{\beta}_1|X) = \mathbb{E}(\bar{y}|X) - \bar{x}\beta_1.$$

- ▶ I will leave it as an exercise to verify that the last expression equals β_1 .
- ▶ This gives $\mathbb{E}(\hat{\beta}_1|X) = \beta_1$. This means that $\hat{\beta}_1$ is conditionally (on X) unbiased for β_1 . This automatically implies unconditional unbiasedness of $\hat{\beta}_1$ as well (why?).
- ▶ Unbiasedness (conditionally on X and unconditionally) also holds for $\hat{\beta}_0$:

$$\mathbb{E}(\hat{\beta}_0|X) = \mathbb{E}(\bar{y} - \bar{x}\hat{\beta}_1|X) = \mathbb{E}(\bar{y}|X) - \bar{x}\beta_1.$$

▶ Show that the last expression equals β_0 .

Examples

