

## Lecture 9

September 19, 2018

## Last class

- ▶ In the last class, we started the study of hypothesis testing in linear regression under the assumption of normality.

# Last class

- ▶ In the last class, we started the study of hypothesis testing in linear regression under the assumption of normality. We assumed that  $Y \sim N_n(X\beta, \sigma^2 I)$  (or equivalently that  $e \sim N_n(0, \sigma^2 I)$ ).

## Last class

- ▶ In the last class, we started the study of hypothesis testing in linear regression under the assumption of normality. We assumed that  $Y \sim N_n(X\beta, \sigma^2 I)$  (or equivalently that  $e \sim N_n(0, \sigma^2 I)$ ).
- ▶ Two consequences of the normality assumption are that  $\hat{\beta} \sim N(\beta, \sigma^2(X^T X)^{-1})$  and that  $RSS/\sigma^2 \sim \chi_{n-p-1}^2$ .

# Last class

- ▶ In the last class, we started the study of hypothesis testing in linear regression under the assumption of normality. We assumed that  $Y \sim N_n(X\beta, \sigma^2 I)$  (or equivalently that  $e \sim N_n(0, \sigma^2 I)$ ).
- ▶ Two consequences of the normality assumption are that  $\hat{\beta} \sim N(\beta, \sigma^2(X^T X)^{-1})$  and that  $RSS/\sigma^2 \sim \chi_{n-p-1}^2$ .
- ▶ Moreover,  $\hat{\beta}$  and  $RSS$  are statistically independent.

## Testing $H_0 : \beta_j = 0$

- ▶ We started by testing the hypothesis  $H_0 : \beta_j = 0$ .

## Testing $H_0 : \beta_j = 0$

- ▶ We started by testing the hypothesis  $H_0 : \beta_j = 0$ . We saw the t-test for testing this. The t-test is based on the statistic:

$$\frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)}$$

which is distributed according to the t-distribution with  $n - p - 1$  degrees of freedom.

## Testing $H_0 : \beta_j = 0$

- ▶ We started by testing the hypothesis  $H_0 : \beta_j = 0$ . We saw the t-test for testing this. The t-test is based on the statistic:

$$\frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)}$$

which is distributed according to the t-distribution with  $n - p - 1$  degrees of freedom.

- ▶ The  $p$ -value for testing  $H_0 : \beta_j = 0$  can then be got by

$$\mathbb{P} \left( |t_{n-r}| > \left| \frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)} \right| \right).$$



## Testing $H_0 : \beta_j = 0$

- ▶ We started by testing the hypothesis  $H_0 : \beta_j = 0$ . We saw the t-test for testing this. The t-test is based on the statistic:

$$\frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)}$$

which is distributed according to the t-distribution with  $n - p - 1$  degrees of freedom.

- ▶ The  $p$ -value for testing  $H_0 : \beta_j = 0$  can then be got by

$$\mathbb{P} \left( |t_{n-r}| > \left| \frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)} \right| \right).$$

$R = p+1$

- ▶ Note that when  $n - r$  is large, the  $t$ -distribution is almost the same as a standard normal distribution.

## Testing $H_0 : \beta_j = 0$

- ▶ The p-value above is for the two-sided alternative  $H_1 : \beta_j \neq 0$ .

## Testing $H_0 : \beta_j = 0$

- ▶ The p-value above is for the two-sided alternative  $H_1 : \beta_j \neq 0$ . If you want to test against one-sided alternatives  $H_1 : \beta_j > 0$  or  $\beta_j < 0$ , then just divide the above p-value by 2.

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ There is another way of testing the null hypothesis that  $H_0 : \beta_j = 0$ .

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ There is another way of testing the null hypothesis that  $H_0 : \beta_j = 0$ . This turns out to be identical to the  $t$ -test but can be generalized more easily to other tests.

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ There is another way of testing the null hypothesis that  $H_0 : \beta_j = 0$ . This turns out to be identical to the  $t$ -test but can be generalized more easily to other tests.
- ▶ The  $F$ -test for  $H_0 : \beta_j = 0$  is based on the following idea. The null hypothesis  $H_0$  says that the explanatory variable  $x_j$  can be dropped from the linear model. Let us call this reduced model  $m$ .

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ There is another way of testing the null hypothesis that  $H_0 : \beta_j = 0$ . This turns out to be identical to the  $t$ -test but can be generalized more easily to other tests.
- ▶ The  $F$ -test for  $H_0 : \beta_j = 0$  is based on the following idea. The null hypothesis  $H_0$  says that the explanatory variable  $x_j$  can be dropped from the linear model. Let us call this reduced model  $m$ .
- ▶ Also, let us call the original model  $M$  (this is the full model:  $y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$ ).

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ There is another way of testing the null hypothesis that  $H_0 : \beta_j = 0$ . This turns out to be identical to the  $t$ -test but can be generalized more easily to other tests.
- ▶ The  $F$ -test for  $H_0 : \beta_j = 0$  is based on the following idea. The null hypothesis  $H_0$  says that the explanatory variable  $x_j$  can be dropped from the linear model. Let us call this reduced model  $m$ .
- ▶ Also, let us call the original model  $M$  (this is the full model:  $y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$ ).
- ▶ The model  $m$  is a *submodel* of  $M$ .



## $F$ -test for $H_0 : \beta_j = 0$

- ▶ The following presents another natural test for  $H_0$ . Let the Residual

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ The following presents another natural test for  $H_0$ . Let the Residual Sum of Squares in the model  $m$  be denoted by  $RSS(m)$  and let the RSS in the full model be  $RSS(M)$ .

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ The following presents another natural test for  $H_0$ . Let the Residual Sum of Squares in the model  $m$  be denoted by  $RSS(m)$  and let the RSS in the full model be  $RSS(M)$ .
- ▶ It is always true that  $RSS(M) \leq RSS(m)$ .

## F-test for $H_0 : \beta_j = 0$

- ▶ The following presents another natural test for  $H_0$ . Let the Residual Sum of Squares in the model  $m$  be denoted by  $RSS(m)$  and let the RSS in the full model be  $RSS(M)$ .
- ▶ It is always true that  $RSS(M) \leq RSS(m)$ .
- ▶ Now if  $RSS(M)$  is *much smaller* than  $RSS(m)$ , it means that the explanatory variable  $x_j$  contributes a lot to the regression and hence cannot be dropped i.e., we reject the null hypothesis  $H_0$ .

## F-test for $H_0 : \beta_j = 0$

- ▶ The following presents another natural test for  $H_0$ . Let the Residual Sum of Squares in the model  $m$  be denoted by  $RSS(m)$  and let the RSS in the full model be  $RSS(M)$ .
- ▶ It is always true that  $RSS(M) \leq RSS(m)$ .
- ▶ Now if  $RSS(M)$  is *much smaller* than  $RSS(m)$ , it means that the explanatory variable  $x_j$  contributes a lot to the regression and hence cannot be dropped i.e., we reject the null hypothesis  $H_0$ .
- ▶ On the other hand, if  $RSS(M)$  is *only a little smaller* than  $RSS(m)$ , then  $x_j$  does not really contribute a lot in predicting  $y$  and hence can be dropped i.e., we do not reject  $H_0$ .

## $F$ -test for $H_0 : \beta_j = 0$

- Therefore one can test  $H_0$  via the test statistic:

$$RSS(m) - RSS(M).$$

## F-test for $H_0 : \beta_j = 0$

- ▶ Therefore one can test  $H_0$  via the test statistic:

$$RSS(m) - RSS(M).$$

- ▶ We would reject the null hypothesis if this is large. How large?

## F-test for $H_0 : \beta_j = 0$

- ▶ Therefore one can test  $H_0$  via the test statistic:

$$RSS(m) - RSS(M).$$

- ▶ We would reject the null hypothesis if this is large. How large?
- ▶ To answer this, we need to look at the **null distribution** of  $RSS(m) - RSS(M)$ .



## F-test for $H_0 : \beta_j = 0$

- ▶ Therefore one can test  $H_0$  via the test statistic:

$$RSS(m) - RSS(M).$$

- ▶ We would reject the null hypothesis if this is large. How large?
- ▶ To answer this, we need to look at the **null distribution** of  $RSS(m) - RSS(M)$ .
- ▶ We will show later today that

$$\frac{RSS(m) - RSS(M)}{\sigma^2} \sim \chi_1^2$$

under the null hypothesis.

## F-test for $H_0 : \beta_j = 0$

- ▶ Therefore one can test  $H_0$  via the test statistic:

$$RSS(m) - RSS(M).$$

- ▶ We would reject the null hypothesis if this is large. How large?
- ▶ To answer this, we need to look at the **null distribution** of  $RSS(m) - RSS(M)$ .
- ▶ We will show later today that whether use M or m to estimate sigma square?

$$\frac{RSS(m) - RSS(M)}{\sigma^2} \sim \chi_1^2$$

under the null hypothesis.

- ▶ Since we do not know  $\sigma^2$ , we estimate it by the square of the residual standard error:

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

This one is independent of  $RSS(m) - RSS(M)$

## F-test for $H_0 : \beta_j = 0$

- ▶ Since we do not know  $\sigma^2$ , we estimate it by the square of the residual standard error:

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

to obtain the test statistic:

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$

## F-test for $H_0 : \beta_j = 0$

- ▶ Since we do not know  $\sigma^2$ , we estimate it by the square of the residual standard error:

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

to obtain the test statistic:

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$

- ▶ The numerator and the denominator are independent (to be shown later). This independence will not hold if the denominator were  $RSS(m)/(n - p)$ .

## F-test for $H_0 : \beta_j = 0$

- ▶ Since we do not know  $\sigma^2$ , we estimate it by the square of the residual standard error:

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

to obtain the test statistic:

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$

- ▶ The numerator and the denominator are independent (to be shown later). This independence will not hold if the denominator were  $RSS(m)/(n - p)$ .
- ▶ Thus under the null hypothesis

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - m)} \sim F_{1, n-p-1}.$$

## F-test for $H_0 : \beta_j = 0$

- ▶ Since we do not know  $\sigma^2$ , we estimate it by the square of the residual standard error:

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

to obtain the test statistic:

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$

- ▶ The numerator and the denominator are independent (to be shown later). This independence will not hold if the denominator were  $RSS(m)/(n - p)$ .
- ▶ Thus under the null hypothesis

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - m)} \sim F_{1, n-p-1}.$$

$p$ -value can therefore be got by

$$\mathbb{P} \left( F_{1, n-p-1} > \frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)} \right).$$

## $F$ -test for $H_0 : \beta_j = 0$

- ▶ This is the p-value for the two-sided alternative. For the one-sided alternative, divide by two.

# Equivalence of These Two Tests

- ▶ It turns out that these two tests for testing  $H_0 : \beta_j = 0$  are equivalent in the sense that they give the same  $p$ -value.



# Equivalence of These Two Tests

- ▶ It turns out that these two tests for testing  $H_0 : \beta_j = 0$  are equivalent in the sense that they give the same  $p$ -value.
- ▶ This is because

$$\left( \frac{\hat{\beta}_j}{s.e(\hat{\beta}_j)} \right)^2 = \frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$

This is not very difficult to prove but we shall skip its proof.

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables.

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ .

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ .  
Examples include:

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ . Examples include:

1. For the constraint  $\beta_1 = 0$ , the model  $m$  becomes:

$$y_i = \beta_0 + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ . Examples include:

1. For the constraint  $\beta_1 = 0$ , the model  $m$  becomes:

$$y_i = \beta_0 + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

2. For  $\beta_1 = \beta_2 = 0$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ . Examples include:

1. For the constraint  $\beta_1 = 0$ , the model  $m$  becomes:

$$y_i = \beta_0 + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

2. For  $\beta_1 = \beta_2 = 0$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

3. For  $\beta_1 = \cdots = \beta_p = 0$ , the model  $m$  becomes  $y_i = \beta_0 + e_i$ .



# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ . Examples include:

1. For the constraint  $\beta_1 = 0$ , the model  $m$  becomes:

$$y_i = \beta_0 + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

2. For  $\beta_1 = \beta_2 = 0$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

3. For  $\beta_1 = \cdots = \beta_p = 0$ , the model  $m$  becomes  $y_i = \beta_0 + e_i$ .

4. For  $\beta_1 = \beta_2$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_1(x_{i1} + x_{i2}) + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

# General Linear Hypothesis Testing via the F-test

- ▶ Let  $M$  denote the full regression model:

$$y_i = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip} + e_i$$

which has  $p$  explanatory variables. Let  $r$  denote the rank of  $X$  in this full model.

- ▶ Let  $m$  denote a sub-model of  $M$  that is obtained by a linear constraint on the parameter  $\beta = (\beta_0, \dots, \beta_p)$  of  $M$ . Examples include:

1. For the constraint  $\beta_1 = 0$ , the model  $m$  becomes:

$$y_i = \beta_0 + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

2. For  $\beta_1 = \beta_2 = 0$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

3. For  $\beta_1 = \cdots = \beta_p = 0$ , the model  $m$  becomes  $y_i = \beta_0 + e_i$ .

4. For  $\beta_1 = \beta_2$ , the model  $m$  becomes

$$y_i = \beta_0 + \beta_1 (x_{i1} + x_{i2}) + \beta_3 x_{i3} + \cdots + \beta_p x_{ip} + e_i.$$

5. For  $\beta_1 = 3$ , the model  $m$  becomes

$$y_i = \beta_0 + 3x_{i1} + \beta_2 x_{i2} + \cdots + \beta_p x_{ip} + e_i.$$

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?
- ▶ This test can be carried out by noting that if  $RSS(m) - RSS(M)$  is large, then  $m$  is not a good model for the data and we therefore reject  $H_0$ .

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?
- ▶ This test can be carried out by noting that if  $RSS(m) - RSS(M)$  is large, then  $m$  is not a good model for the data and we therefore reject  $H_0$ . On the other hand, if  $RSS(m) - RSS(M)$  is small, we do not reject  $H_0$ .

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?
- ▶ This test can be carried out by noting that if  $RSS(m) - RSS(M)$  is large, then  $m$  is not a good model for the data and we therefore reject  $H_0$ . On the other hand, if  $RSS(m) - RSS(M)$  is small, we do not reject  $H_0$ .
- ▶ The test is therefore based on  $RSS(m) - RSS(M)$ .

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?
- ▶ This test can be carried out by noting that if  $RSS(m) - RSS(M)$  is large, then  $m$  is not a good model for the data and we therefore reject  $H_0$ . On the other hand, if  $RSS(m) - RSS(M)$  is small, we do not reject  $H_0$ .
- ▶ The test is therefore based on  $RSS(m) - RSS(M)$ .
- ▶ What is the distribution of this quantity under the null hypothesis?

# General Linear Hypothesis Testing via the F-test

- ▶ How do we test the hypothesis  $H_0 : m$  against  $H_1 : M$ ?
- ▶ This test can be carried out by noting that if  $RSS(m) - RSS(M)$  is large, then  $m$  is not a good model for the data and we therefore reject  $H_0$ . On the other hand, if  $RSS(m) - RSS(M)$  is small, we do not reject  $H_0$ .
- ▶ The test is therefore based on  $RSS(m) - RSS(M)$ .
- ▶ What is the distribution of this quantity under the null hypothesis?
- ▶ We will show that

$$\frac{RSS(m) - RSS(M)}{\sigma^2} \sim \chi^2_{p-q}$$

where  $q$  is the number of explanatory variables in  $m$  and  $p$  is the number of explanatory variables in  $M$ . Let us now prove this fact.



## General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively.

## General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

## General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

so that

$$RSS(m) - RSS(M) = Y^T(H(M) - H(m))Y.$$

# General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

so that

$$RSS(m) - RSS(M) = Y^T(H(M) - H(m))Y.$$

- ▶ We need the null distribution of  $RSS(m) - RSS(M)$ .

## General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

so that

$$RSS(m) - RSS(M) = Y^T(H(M) - H(m))Y.$$

- ▶ We need the null distribution of  $RSS(m) - RSS(M)$ . So we shall assume that  $Y = X(m)\beta(m) + e$  (where  $X(m)$  is the  $X$ -matrix in the model  $m$ ). Why?

## General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

so that

$$RSS(m) - RSS(M) = Y^T(H(M) - H(m))Y.$$

- ▶ We need the null distribution of  $RSS(m) - RSS(M)$ . So we shall assume that  $Y = X(m)\beta(m) + e$  (where  $X(m)$  is the  $X$ -matrix in the model  $m$ ). Why? It is important to realize that  $H(m)X(m) = X(m)$  and also  $H(M)X(m) = X(m)$ .

# General Linear Hypothesis Testing via the F-test

- ▶ Let the hat matrices in the two models be denoted by  $H(m)$  and  $H(M)$  respectively. Write

$$RSS(m) = Y^T(I - H(m))Y \quad \text{and} \quad RSS(M) = Y^T(I - H(M))Y$$

so that

$$RSS(m) - RSS(M) = Y^T(H(M) - H(m))Y.$$

- ▶ We need the null distribution of  $RSS(m) - RSS(M)$ . So we shall assume that  $Y = X(m)\beta(m) + e$  (where  $X(m)$  is the  $X$ -matrix in the model  $m$ ). Why? It is important to realize that  $H(m)X(m) = X(m)$  and also  $H(M)X(m) = X(m)$ . So  $\text{Plug in } Y = X(m)\beta(m) + e$

$$RSS(m) - RSS(M) = e^T(H(M) - H(m))e.$$

## General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . Prove it by taking square and see if we can get the same matrix



## General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . To see that  $H(M) - H(m)$  is idempotent notice that

$$(H(M) - H(m))^2 = H(M) + H(m) - 2H(M)H(m). \quad (1)$$

## General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . To see that  $H(M) - H(m)$  is idempotent notice that

$$(H(M) - H(m))^2 = H(M) + H(m) - 2H(M)H(m). \quad (1)$$

- ▶ Now  $H(M)H(m) = H(m)$ .

## General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . To see that  $H(M) - H(m)$  is idempotent notice that

$$(H(M) - H(m))^2 = H(M) + H(m) - 2H(M)H(m). \quad (1)$$

- ▶ Now  $H(M)H(m) = H(m)$ . To see this, it is enough to show that  $H(M)H(m)v = H(m)v$  for every vector  $v$ .

# General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . To see that  $H(M) - H(m)$  is idempotent notice that

Take transpose

$$(H(M) - H(m))^2 = H(M) + H(m) - 2H(M)H(m). \quad (1)$$

- ▶ Now  $H(M)H(m) = H(m)$ . To see this, it is enough to show that  $H(M)H(m)v = H(m)v$  for every vector  $v$ .
- ▶ Now recall that  $H(m)v$  denotes the projection of  $v$  onto the column space of  $X(m)$ . And  $H(M)H(m)v$  projects  $H(m)v$  onto the column space of  $X(M)$  (which equals the original  $X$  matrix).

# General Linear Hypothesis Testing via the F-test

- ▶  $H(M) - H(m)$  is a symmetric and idempotent  $n \times n$  matrix of rank  $p - q$ . To see that  $H(M) - H(m)$  is idempotent notice that

$$(H(M) - H(m))^2 = H(M) + H(m) - 2H(M)H(m). \quad (1)$$

Why  $H(m)H(M) = H(m)^T$ ?

- ▶ Now  $H(M)H(m) = H(m)$ . To see this, it is enough to show that  $H(M)H(m)v = H(m)v$  for every vector  $v$ .
- ▶ Now recall that  $H(m)v$  denotes the projection of  $v$  onto the column space of  $X(m)$ . And  $H(M)H(m)v$  projects  $H(m)v$  onto the column space of  $X(M)$  (which equals the original  $X$  matrix).
- ▶ But because the column space of  $X(m)$  is contained in the column space of  $X(M)$ , it follows that  $H(m)v$  is already contained in the column space of  $X(M)$ . Thus its projection onto the column space of  $X(M)$  equals itself. So  $H(M)H(m)v = H(m)v$ .

## General Linear Hypothesis Testing via the F-test

- ▶ Because  $H(M)H(m) = H(m)$ , it follows from (1) that

$$(H(M) - H(m))^2 = H(M) - H(m).$$

# General Linear Hypothesis Testing via the F-test

- ▶ Because  $H(M)H(m) = H(m)$ , it follows from (1) that

$$(H(M) - H(m))^2 = H(M) - H(m).$$

- ▶ Therefore, under the null hypothesis

$$\frac{RSS(m) - RSS(M)}{\sigma^2} = \left(\frac{e}{\sigma}\right)^T (H(M) - H(m)) \frac{e}{\sigma} \sim \chi_{p-q}^2.$$

# General Linear Hypothesis Testing via the F-test

- ▶ Because  $H(M)H(m) = H(m)$ , it follows from (1) that

$$(H(M) - H(m))^2 = H(M) - H(m).$$

- ▶ Therefore, under the null hypothesis

$\text{trace}(H(M) - H(m)) = \text{rank of it because symmetric and idempotent}$

$$\frac{RSS(m) - RSS(M)}{\sigma^2} = \left(\frac{e}{\sigma}\right)^T (H(M) - H(m)) \frac{e}{\sigma} \sim \chi_{p-q}^2.$$

- ▶ Since we do not know  $\sigma^2$ , we estimate it by

$$\hat{\sigma}^2 = \frac{RSS(M)}{n - p - 1},$$

to obtain the test statistic:

$$\frac{RSS(m) - RSS(M)}{RSS(M)/(n - p - 1)}$$



# General Linear Hypothesis Testing via the F-test

- ▶ The numerator and the denominator are independent because  $RSS(m) - RSS(M) = Y^T (H(M) - H(m)) Y$  and  $RSS(M) = Y^T (I - H(M)) Y$  and the product of the matrices

$$(H(M) - H(m)) (I - H(M)) = H(M) - H(M)^2 - H(m) + H(m)H(M)$$

# General Linear Hypothesis Testing via the F-test

- ▶ The numerator and the denominator are independent because  $RSS(m) - RSS(M) = Y^T (H(M) - H(m)) Y$  and  $RSS(M) = Y^T (I - H(M)) Y$  and the product of the matrices

$$(H(M) - H(m))(I - H(M)) = H(M) - H(M)^2 - H(m) + H(m)H(M)$$

- ▶ Thus under the null hypothesis

$$\frac{(RSS(m) - RSS(M))/(p - q)}{RSS(M)/(n - p - 1)} \sim F_{p-q, n-p-1}.$$

# General Linear Hypothesis Testing via the F-test

- ▶ The numerator and the denominator are independent because  $RSS(m) - RSS(M) = Y^T (H(M) - H(m)) Y$  and  $RSS(M) = Y^T (I - H(M)) Y$  and the product of the matrices

$$(H(M) - H(m)) (I - H(M)) = H(M) - H(M)^2 - H(m) + H(m)H(M)$$

- ▶ Thus under the null hypothesis

$$\frac{(RSS(m) - RSS(M))/(p - q)}{RSS(M)/(n - p - 1)} \sim F_{p-q, n-p-1}.$$

- ▶  $p$ -value can therefore be got by

$$\mathbb{P} \left( F_{p-q, n-p-1} > \frac{(RSS(m) - RSS(M))/(p - q)}{RSS(M)/(n - p - 1)} \right).$$

# General Linear Hypothesis Testing via the F-test

- ▶ If the null hypothesis can be written in terms of a single linear function of  $\beta$ , such as  $H_0 : \beta_1 + 5\beta_3 = 5$ . Then it can also be tested via the  $t$ -test; using the statistic:

$$\frac{\hat{\beta}_1 + 5\hat{\beta}_3 - 5}{s.e(\hat{\beta}_1 + 5\hat{\beta}_3)}$$

which has the  $t$ -distribution with  $n - p - 1$  degrees of freedom under  $H_0$ . This test and the corresponding  $F$ -test will have the same  $p$ -value.

# Testing for all explanatory variables

- ▶ How do we test  $H_0 : \beta_1 = \cdots = \beta_p = 0$  against its complement?

# Testing for all explanatory variables

- ▶ How do we test  $H_0 : \beta_1 = \cdots = \beta_p = 0$  against its complement?
- ▶ Just take  $m$  to be the model  $y_i = \beta_0 + e_i$ .

# Testing for all explanatory variables

- ▶ How do we test  $H_0 : \beta_1 = \dots = \beta_p = 0$  against its complement?
- ▶ Just take  $m$  to be the model  $y_i = \beta_0 + e_i$ . In this case,  $RSS(m) = TSS$  and  $q = 0$  and  $RSS(M) = RSS$ .

# Testing for all explanatory variables

- ▶ How do we test  $H_0 : \beta_1 = \dots = \beta_p = 0$  against its complement?
- ▶ Just take  $m$  to be the model  $y_i = \beta_0 + e_i$ . In this case,  $RSS(m) = TSS$  and  $q = 0$  and  $RSS(M) = RSS$ .
- ▶ Thus the  $p$ -value is

$$\mathbb{P} \left\{ F_{p, n-p-1} > \frac{(TSS - RSS)/p}{RSS/(n - p - 1)} \right\}.$$